Solve with us 10 - Not Graded

This assignment will not be graded and is only for practice.
Key Points: 1Key Points: 1

- Gradient of a function:Gradient of a function: If ff is a scalar valued function defined on a domain DD in R^n Rn, then the gradient ∇f∇f is defined as the vector valued function $(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \ \dots, \frac{\partial f}{\partial x_n})$(∂x1∂f ,∂x2∂f, …, ∂xn∂f).
- If ∇f∇f exists and is continuous in an open ball around the point $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an), then the directional derivative of the function ff at $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an) in the direction of a vector vv is given by the dot product of the gradient of the function at $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an) with the unit vector in the direction of vv i.e.,

$$D_v f(a_1, a_2, \ \dots, a_n) = \nabla f(a_1, a_2, \ \dots, a_n).\frac{v}{||v||}$$Dvf(a1, a2, …, an) = ∇f(a1, a2, …, an).||v||v

- If ∇f∇f exists and is continuous in an open ball around the point $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an), then the gradient is the direction in which the directional derivative at PP is maximized amongst all directions, i.e., the directional derivative at PP is maximized amongst all the directions in the direction of the vector $\frac{\nabla f(a_1, a_2, \ \dots, a_n)}{||\nabla f(a_1, a_2, \ \dots, a_n)||}$||∇f(a1,a2,…,an)||∇f(a1,a2,…,an).
- If ∇f∇f exists and is continuous in an open ball around the point $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an), then the direction opposite to the gradient is the direction in which the directional derivative at PP is minimized amongst all directions, i.e., the directional derivative at PP is minimized amongst all the direction in the direction of the vector $\frac{-\nabla f(a_1, a_2, \ \dots, a_n)}{||\nabla f(a_1, a_2, \ \dots, a_n)||}$||∇f(a1,a2,…,an)||−∇f(a1,a2,…,an).
- If ∇f∇f exists and is continuous in an open ball around the point $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an), then the maximum value amongst all directional derivatives at PP is

$$\nabla f(a_1, a_2, \ \dots, a_n).\frac{\nabla f(a_1, a_2, \ \dots, a_n)}{||\nabla f(a_1, a_2, \ \dots, a_n)||} = \frac{||\nabla f(a_1, a_2, \ \dots, a_n)||^2}{||\nabla f(a_1, a_2, \ \dots, a_n)||} = ||\nabla f(a_1, a_2, \ \dots, a_n)||$$∇f(a1, a2, …, an).||∇f(a1,a2,…,an)||∇f(a1,a2,…,an) = ||∇f(a1,a2,…,an)||||∇f(a1,a2,…,an)||2 = ||∇f(a1, a2, …, an)||

Similarly, If ∇f∇f exists and is continuous in an open ball around the point $(a_1, a_2, \ \dots, a_n)$(a1, a2, …, an), then the minimum value amongst all directional derivatives at PP is

$$\nabla f(a_1, a_2, \ \dots, a_n).\frac{-\nabla f(a_1, a_2, \ \dots, a_n)}{||\nabla f(a_1, a_2, \ \dots, a_n)||} = -\frac{||\nabla f(a_1, a_2, \ \dots, a_n)||^2}{||\nabla f(a_1, a_2, \ \dots, a_n)||} = -||\nabla f(a_1, a_2, \ \dots, a_n)||$$∇f(a1, a2, …, an).||∇f(a1,a2,…,an)||−∇f(a1,a2,…,an) = −||∇f(a1,a2,…,an)||||∇f(a1,a2,…,an)||2 = −||∇f(a1, a2, …, an)||

1 point
Find the gradient of the scalar valued function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$.
f(x, y, z) = 2x3 + xy2 + 6z2 + y2 + xyz.

○
$(6x^2, 2y, 12z)$(6x2, 2y, 12z)

○

$(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)$(6x2 + y2 + yz, 2xy + 2y + xz, 12z + xy)

○

$(6x^2 + y^2, 2xy + 2y, 12z)$(6x2 + y2, 2xy + 2y, 12z)

○

$(6x^2 + yz, 2y + xz, 12z + xy)$(6x2 + yz, 2y + xz, 12z + xy)

No, the answer is incorrect.

Score: 0

Accepted Answers:

$(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)$(6x2 + y2 + yz, 2xy + 2y + xz, 12z + xy)

1 point

Find the directional derivative of $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$
f(x, y, z) = 2x3 + xy2 + 6z2 + y2 + xyz at (2, 1, 2)(2, 1, 2) in the direction of a vector (0, 3, 0)(0, 3, 0).

○

3030

○

00

○

33

○

1010

No, the answer is incorrect.

Score: 0

Feedback:

Accepted Answers:

1010

1 point

Find the direction in which the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$
f(x, y, z) = 2x3 + xy2 + 6z2 + y2 + xyz increases the fastest from the point (1, 1, 1)(1, 1, 1)?

○

(1, 1, 1)(1, 1, 1)

○

$(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$(3

√

1, 3

√

1, 3

√

1)

○

(8, 5, 13)(8, 5, 13)

○

$(\frac{8}{\sqrt{258}}, \frac{5}{\sqrt{258}}, \frac{13}{\sqrt{258}})$(258

√

8, 258

√

5, 258

√

13)

No, the answer is incorrect.
Score: 0
Accepted Answers:
$(\frac{8}{\sqrt{258}}, \frac{5}{\sqrt{258}}, \frac{13}{\sqrt{258}})$(258
√
8, 258
√
5, 258
√
13)
1 point
Find the maximum directional derivative of the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ f(x, y, z) = 2x3 + xy2 + 6z2 + y2 + xyz at the point $P = (1, 2, 3)$P = (1, 2, 3).
○
$\sqrt{1821}$ 1821
√
○
4343
○
$\sqrt{14}$ 14
√
○
155155
No, the answer is incorrect.
Score: 0
Accepted Answers:
$\sqrt{1821}$ 1821
√

1 point
Find the minimum directional derivative of the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ f(x, y, z) = 2x3 + xy2 + 6z2 + y2 + xyz at the point $P = (1, 0, 0)$P = (1, 0, 0).
○
−1−1
○
66
○
−6−6
○
11
No, the answer is incorrect.
Score: 0
Accepted Answers:
−6−6
1 point
Let $f(x, y) = x^2 y^2 + y^3$ f(x, y) = x2y2 + y3. Find the directional derivative of f f at the point (1, 1) (1, 1) in the direction which forms an angle (counterclockwise) of $\frac{\pi}{3}$3π with the positive xx-axis.
○
$\sqrt{29}$ 29

√

○
√29 + √2 29
√
+ 2
√

○
√2 2
√

○
1 + 5√3/2 1 + 25 3
√

No, the answer is incorrect.
Score: 0
Feedback:

Accepted Answers:
1 + 5√3/2 1 + 25 3
√

Key Points: 2Key Points: 2

- Let $f(x, y)$f(x, y) be a function defined on a domain DD in R^2 R2 containing some open ball around the point (a, b)(a, b). Equations of the tangent lines to the function f f at the point (a, b)(a, b) in the direction of unit vector (u_1, u_2)(u1, u2):

  i)Parametric equations:Parametric equations: $x(t) = a + tu_1$ x(t) = a + tu1, $y(t) = b + tu_2$ y(t) = b + tu2, $z(t) = f(a, b) + tf_u(a, b)$z(t) = f(a, b) + tfu(a, b).
  ii) Symmetric equations:Symmetric equations: $\frac{x-a}{u_1} = \frac{y-b}{u_2} = \frac{z-f(a,b)}{f_u(a,b)}$u1x−a = u2y−b = fu (a,b)z−f(a,b).
  iii)Vector form:Vector form: $(x(t), y(t), z(t)) = (a, b, f(a, b)) + t(u_1, u_2, f_u(a, b))$ (x(t), y(t), z(t)) = (a, b, f(a, b)) + t(u1, u2, fu(a, b)).
- The equation of the tangent plane of the graph of a function of two variables $z = f(x, y)$z = f(x, y) at the point (x_0, y_0, z_0)(x0, y0, z0) (where, $z_0 = f(x_0, y_0)$z0 = f(x0, y0)) is given by

  $f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) - 1(z - z_0) = 0$fx(x0, y0)(x − x0) + fy(x0, y0) (y − y0) − 1(z − z0) = 0

  $f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) + f(x_0, y_0) = z$fx(x0, y0)(x − x0) + fy(x0, y0) (y − y0) + f(x0, y0) = z
- Linear approximation of the function $z = f(x, y)$z = f(x, y) at the point (x_0, y_0, z_0)(x0, y0, z0) is given by

$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$L(x, y) = f(x0, y0) + fx (x0, y0)(x − x0) + fy(x0, y0)(y − y0)

- Let $z = f(x\sim) = f(x_1, x_2, \ldots, x_n)$z = f(∼x) = f(x1, x2, … , xn) be a function defined on a domain DD in R^2 R2 containing some open ball around the point $a\sim = (a_1, a_2, \ldots, a_n)\sim$a = (a1, a2, … , an). Equations of the tangent lines to the function ff at the point $a\sim\sim$a in the direction of unit vector $u\sim = (u_1, u_2, \ldots, u_n)\sim$u = (u1, u2, … , un):

  i)Parametric equations:Parametric equations: $x_i(t) = a_i + tu_i$ xi(t) = ai + tui, $z(t) = f(a\sim) + tf_{u\sim}(a\sim)$z(t) = f(∼a) + tf∼u(∼a).

  ii)Vector form:Vector form: $(x\sim(t), z(t)) = (a\sim, f(a\sim)) + t(u\sim, f_{u\sim}(a\sim))$.(∼x(t), z(t)) = (∼a, f(∼a)) + t(∼u, f∼u(∼a)).
- Let $f(x\sim)$f(∼x) be a function defined on domain DD in R^n Rn containing some open ball around the point $a\sim\sim$a. Suppose ∇f∇f exists and is continuous on some open ball around the point $a\sim\sim$a. Equation of tangent plane to ff at $a\sim\sim$a:

  $$z = f(a\sim) + \nabla f(a\sim).(x\sim - a\sim).$$z = f(∼a) + ∇f(∼a).(∼x − ∼a).

  $$z = f(a\sim) + \sum \frac{\partial f}{\partial x_i}(a\sim)(x_i - a_i)$$z = f(∼a) + ∑ ∂xi∂f(∼a)(xi − ai)

1 point
Find the equation of the tangent plane to the graph of the function given by
$z = f(x, y) = \sqrt{r^2 - x^2 - y^2}$z = f(x, y) = r2 − x2 − y2
√
at the point $(0, 0, r)$(0, 0, r).

○
$x + y + z = r$x + y + z = r

○
$2x + 2y + 2z = r$2x + 2y + 2z = r

○
$z = r$z = r

○
$x + y = r$x + y = r

No, the answer is incorrect.
Score: 0
Accepted Answers:
$z = r$z = r
1 point
Find the equation of the tangent plane to the graph of the function given by
$z = f(x, y) = \frac{1}{\sqrt{c}}\sqrt{d - ax^2 - by^2}$z = f(x, y) = c
√
1 d − ax2 − by2

$\sqrt{}$

at the point $(1, 2, f(1, 2))$(1, 2, f(1, 2)).

○

$ax + 2by + \sqrt{c(d - a - 4b)}\ z$ = 0ax + 2by + c(d − a − 4b)

$\sqrt{}$

z = 0

○

$ax + 2by + \sqrt{c(d - a - 4b)}\ z = d$ax + 2by + c(d − a − 4b)

$\sqrt{}$

z = d

○

$2ax + 4by + \sqrt{c(d - a - 4b)}\ z = d$2ax + 4by + c(d − a − 4b)

$\sqrt{}$

z = d

○

$ax + 2by + \sqrt{c(d + a + 4b)}\ z = d$ax + 2by + c(d + a + 4b)

$\sqrt{}$

z = d

No, the answer is incorrect.

Score: 0

Accepted Answers:

$ax + 2by + \sqrt{c(d - a - 4b)}\ z = d$ax + 2by + c(d − a − 4b)

$\sqrt{}$

z = d

1 point

Find the equation of the tangent plane to the graph of the function $z = f(x, y) = x^2 y$z = f(x, y) = x2y at the point (1, 1, 1)(1, 1, 1).

○

$2x + y - z$ = 22x + y − z = 2

○

$2x + y + z$ = 22x + y + z = 2

○

$2x + y - z$ = 02x + y − z = 0

○

$2x + y$ = 22x + y = 2

No, the answer is incorrect.

Score: 0

Accepted Answers:

$2x + y - z$ = 22x + y − z = 2

Key Points: 3Key Points: 3

- Definition:Definition: Consider a scalar valued multivariable function $f: R^n \rightarrow R$f : Rn → R. A point $(a_1, a_2, \ldots, a_n) \in R^n$(a1, a2, … , an) ∈ Rn is said to be a critical point of the function ff, if $\nabla f(a_1, a_2, \ldots, a_n) = 0$∇f(a1, a2, … , an) = 0 or at least one of $f_{x_i}(a_1, a_2, \ldots, a_n)$fxi(a1, a2, … , an) for $i = 1, 2, \ldots, n$i = 1, 2, … , n does not exist.
- Steps to find critical points of a function,

i) Find the partial derivatives f_{x_i} fxi for $i = 1, 2, \ldots, n$i = 1, 2, … , n. If one of $f_{x_i}(a_1, a_2, \ldots, a_n)$fxi(a1, a2, … , an) for $i = 1, 2, \ldots, n$i = 1, 2, … , n does not exist then $(a_1, a_2, \ldots, a_n)$(a1, a2, … , an) is a critical point.

ii) Otherwise, find the all possible point $(a_1, a_2, \ldots, a_n) \in R^n$ (a1, a2, … , an) ∈ Rn such that $f_{x_i}(a_1, a_2, \ldots, a_n) = 0$fxi(a1, a2, … , an) = 0 for all $i = 1, 2, \ldots n$i = 1, 2, … n, simultaneously.

- A function $f(x_1, x_2, \ldots, x_n)$f(x1, x2, … , xn) has a local minimum at the point $(a_1, a_2, \ldots, a_n)$(a1, a2, … , an) if $f(x_1, x_2, \ldots, x_n) \geq f(a_1, a_2, \ldots, a_n)$f(x1, x2, … , xn) ≥ f(a1, a2, … , an) for all points $(x_1, x_2, \ldots, x_n)$(x1, x2, … , xn) in some open ball around $(a_1, a_2, \ldots, a_n)$(a1, a2, … , an).
- A function $f(x_1, x_2, \ldots, x_n)$f(x1, x2, … , xn) has a local maximum at the point $(a_1, a_2, \ldots, a_n)$(a1, a2, … , an) if $f(x_1, x_2, \ldots, x_n) \leq f(a_1, a_2, \ldots, a_n)$f(x1, x2, … , xn) ≤ f(a1, a2, … , an) for all points $(x_1, x_2, \ldots, x_n)$(x1, x2, … , xn) in some open ball around $(a_1, a_2, \ldots, a_n)$(a1, a2, … , an).
- Local extrema are critical points.
- Not all critical points are local extrema. Those critical points at which the gradient exists and is 0 but are not local extrema, are called saddle points.
- A global (or absolute) maximum/minimum is a point for which the function value is maximum/minimum amongst all points in the domain. It always exists when the domain is closed and bounded and can be found by comparing the function values at all critical points in the domain as well as on successive boundaries.

1 point

Let $f(x, y) = \frac{x^3}{3} + y^2 + 2xy - 6x - 3y + 4$f(x, y) = 3x3 + y2 + 2xy − 6x − 3y + 4. Which of the following options are true about ff?

☐ $f_y = 0$fy = 0 is a straight line passing through the origin.

☐ $(-1, \frac{5}{2})$(−1, 25) and $(3, -\frac{3}{2})$(3, −23) are critical points.

☐ $f_x(-1, \frac{5}{2}) = 0$fx(−1, 25) = 0

☐ $f_y(3, -\frac{3}{2}) = 2$fy(3, −23) = 2

No, the answer is incorrect.
Score: 0
Accepted Answers:
$(-1, \frac{5}{2})$(−1, 25) and $(3, -\frac{3}{2})$(3, −23) are critical points.
$f_x(-1, \frac{5}{2}) = 0$fx(−1, 25) = 0

1 point

Choose the correct statement(s) about the maxima/minima of $f(x, y) = x^4 + y^4$ f(x, y) = x4 + y4.

☐ There are no critical points.
☐ There is exactly one critical point.
☐ (0, 0)(0, 0) is a point of local minima.

No, the answer is incorrect.
Score: 0
Feedback:

Accepted Answers:
There is exactly one critical point.
(0, 0)(0, 0) is a point of local minima.

1 point

Suppose $f(x, y) = x^2 + 2y^2 + 3xy - 5y$f (x, y) = x2 + 2y2 + 3xy − 5y be a scalar valued function. Which of the following is (are) critical point(s) of $f(x, y)$f(x, y)?

○
$(-10, 15)$(−10, 15)

○
$(0, 0)$(0, 0)

○
$(15, -10)$(15, −10)

○
$(15, 10)$(15, 10)

No, the answer is incorrect.
Score: 0
Feedback:
$f_x(x,y)=2x+3y$ and $f_y(x,y)=3x+4y-5$
Accepted Answers:
$(15, -10)$(15, −10)

1 point
Which of the following statements are correct?

☐
The minimum value of the function $f(x, y) = x^2 + y^2 + 8$f(x, y) = x2 + y2 + 8 is 00.

☐
The minimum value of the function $f(x, y) = x^2 + y^2 + 8 - 4x - 4y$f(x, y) = x2 + y2 + 8 − 4x − 4y is 00.

☐
The maximum value of the function $f(x, y) = -x^2 - y^2$ f(x, y) = −x2 − y2 is 00.

☐
The maximum value of the function $f(x, y) = x^2 + y^2 - 2x + 1$f(x, y) = x2 + y2 − 2x + 1 is 00.

No, the answer is incorrect.
Score: 0
Accepted Answers:
The minimum value of the function $f(x, y) = x^2 + y^2 + 8 - 4x - 4y$f(x, y) = x2 + y2 + 8 − 4x − 4y is 00.
The maximum value of the function $f(x, y) = -x^2 - y^2$ f(x, y) = −x2 − y2 is 00.

1 point
Find the point on the plane $2x - y + 2z = 16$2x − y + 2z = 16 closest to the origin.

○
$(\frac{32}{9}, -\frac{16}{9}, \frac{32}{9})$(932, −916, 932)

○
$(1, -1, \frac{13}{2})$(1, −1, 213)

○
$(\frac{32}{9}, 1, \frac{89}{18})$(932, 1, 1889)

○
$(1, 0, -1)$(1, 0, −1)

No, the answer is incorrect.
Score: 0
Feedback:

Accepted Answers: