



Reflect with us - Week 8



This assignment will not be graded and is only for practice.

1 point

Choose the set of correct options.

☐

Suppose $\beta = \{v_1, v_2, \dots, v_n\}$ is an orthogonal basis of an inner product space V . If there exists some $v \in V$, such that $\langle v, v_i \rangle = 0$ for all $i = 1, 2, \dots, n$, then $v = 0$.

☐

There exists an orthonormal basis for $\mathbb{R}^n$ with the standard inner product.

☐

If P_W denotes the linear transformation which projects the vectors of an inner product space V to a subspace W of V , then $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$, where 0 denotes the zero vector of V .

☐

$\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ cannot represent a matrix corresponding to some projection.

No, the answer is incorrect.

Score: 0

Accepted Answers:

Suppose $\beta = \{v_1, v_2, \dots, v_n\}$ is an orthogonal basis of an inner product space V . If there exists some $v \in V$, such that $\langle v, v_i \rangle = 0$ for all $i = 1, 2, \dots, n$, then $v = 0$.

There exists an orthonormal basis for $\mathbb{R}^n$ with the standard inner product.

If P_W denotes the linear transformation which projects the vectors of an inner product space V to a subspace W of V , then $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$, where 0 denotes the zero vector of V .

$\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ cannot represent a matrix corresponding to some projection.

Solution:

Option 1: Option 1:

$\beta = \{v_1, v_2, \dots, v_n\}$ is an orthonormal basis of the inner product space V .

Step 1: As $v \in V$, we can express v as a linear combination of the vectors

$v_1, v_2, \dots, v_n$ as follows:

$v = \alpha_1 v_1 + \dots + \alpha_n v_n$, where $\alpha_1, \dots, \alpha_n \in \mathbb{R}$

Taking inner product of v with v_1 , we have $\langle v, v_1 \rangle = \langle \alpha_1 v_1 + \dots + \alpha_n v_n, v_1 \rangle = \alpha_1 \langle v_1, v_1 \rangle + \dots + \alpha_n \langle v_n, v_1 \rangle$

Recall two properties of inner products on V :

i) $\langle u_1 + u_2, u_3 \rangle = \langle u_1, u_3 \rangle + \langle u_2, u_3 \rangle$ for any vectors

$u_1, u_2, u_3 \in V$.

ii) $\langle cu_1, u_2 \rangle = c \langle u_1, u_2 \rangle$ for any vectors $u_1, u_2 \in V$ and $c \in \mathbb{R}$.

Using these properties we have,



$$\langle v, v_1 \rangle = \alpha_1 \langle v_1, v_1 \rangle + \alpha_2 \langle v_2, v_1 \rangle + \dots + \alpha_n \langle v_n, v_1 \rangle \quad \langle v, v_1 \rangle = \alpha_1 \langle v_1, v_1 \rangle + \alpha_2 \langle v_2, v_1 \rangle + \dots + \alpha_n \langle v_n, v_1 \rangle.$$

What is the value of $\langle v_1, v_1 \rangle \langle v_1, v_1 \rangle$?

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 1

1 point

What is the value of $\langle v_2, v_1 \rangle \langle v_2, v_1 \rangle$?

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 0

1 point

What is the value of $\langle v_i, v_1 \rangle \langle v_i, v_1 \rangle$, for any $i \in \{2, 3, \dots, n\}$?

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 0

1 point

Feedback: Recall the definition of an orthonormal basis and also observe that β is an orthonormal basis.

If $\langle v, v_1 \rangle = 0$, then what is the value of α_1 ?

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 0

1 point

Feedback: Observe that $\langle v, v_1 \rangle = \alpha_1 \langle v, v_1 \rangle = \alpha_1$

Similarly we can conclude that, $\alpha_i = 0$, for each $i = 1, 2, \dots, n$

Hence, $v = 0v = 0$.

Option 2: Let $\gamma = \{v_1, v_2, \dots, v_n\}$ be a given ordered basis of a vector space. The steps in the process are described below to yield an orthonormal basis $\beta = \{u_1, u_2, \dots, u_n\}$.

Step-1 $w_1 = v_1$, and $u_1 = \frac{w_1}{\|w_1\|}$

Step-2 $w_2 = v_2 - \langle v_2, u_1 \rangle u_1$ and $u_2 = \frac{w_2}{\|w_2\|}$

Step-3 $w_3 = v_3 - \langle v_3, u_1 \rangle u_1 - \langle v_3, u_2 \rangle u_2$ and $u_3 = \frac{w_3}{\|w_3\|}$

Step-i $w_i = v_i - \sum_{j=1}^{i-1} \langle v_i, u_j \rangle u_j$ and $u_i = \frac{w_i}{\|w_i\|}$.

Step-n $w_n = v_n - \sum_{j=1}^{n-1} \langle v_n, u_j \rangle u_j$ and $u_n = \frac{w_n}{\|w_n\|}$.

Hence, Option 2 is correct.

Option 3: The projection of a vector to a subspace: Let V be an inner product space, $v \in V$ and $W \subseteq V$ be a subspace. Then the projection of v onto W is the vector in W , denoted by $P_W(v)$, computed as follows:

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for W . Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for W .

Then $P_W(v) = \sum_{i=1}^n \langle v, v_i \rangle v_i$

Step 1:

$$\begin{aligned} P_W^2(v) &= P_W(P_W(v)) \\ &= P_W\left(\sum_{i=1}^n \langle v, v_i \rangle v_i\right) \\ &= \sum_{i=1}^n P_W(\langle v, v_i \rangle v_i) \\ &= \sum_{i=1}^n \langle v, v_i \rangle P_W(v_i) \end{aligned}$$

$$(\langle v, v_i \rangle v_i) = \sum_{i=1}^n \langle v, v_i \rangle P_W(v_i)$$

1 point

At this point can you identify $P_W(v_i)$?

☐

v_i

☐ 0

No, the answer is incorrect.

Score: 0

Accepted Answers:

v_i

Feedback: As v_i is itself in W , the projection of it onto W is nothing but the vector v_i itself. On applying the definition, we get, $P_W(v_i) = \sum_{j=1}^n \langle v_i, v_j \rangle v_j = v_i$ as $\langle v_i, v_j \rangle = 0$ when $i \neq j$ and $\langle v_i, v_i \rangle = 1$.

Hence we have

$$\begin{aligned} P_W^2(v) &= \sum_{i=1}^n \langle v, v_i \rangle P_W(v_i) = \sum_{i=1}^n \langle v, v_i \rangle v_i = P_W(v) \\ P_W(v_i) &= \sum_{i=1}^n \langle v, v_i \rangle v_i = P_W(v) \end{aligned}$$

So we can conclude $P_W^2 = P_W$.

Step 2: Let $v \in \text{range}(P_W) \cap \text{null space}(P_W)$. So $v \in \text{range}(P_W)$ and $v \in \text{null space}(P_W)$.

As $v \in \text{range}(P_W)$ there exists $v' \in V$ such that $P_W(v') = v$, and as $v \in \text{null space}(P_W)$, we have $P_W(v) = 0$.

$$\begin{aligned}
 P_W(v') &= v \\
 P_W^2(v') &= P_W(P_W(v')) \\
 P_W(v') &= P_W(v) \text{ [as, } P_W^2 = P_W \text{]} \\
 P_W(v') &= 0 \\
 v &= 0
 \end{aligned}$$

$$(v')) = P_W(v) \text{ [as, } P_W^2 = P_W \text{]} = 0 = 0$$

Hence $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$ $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$.

Option 4: Option 4: If a matrix A represents a projection, then A^2 must be A .

Let $A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ $A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$.

1 point

Is $A^2 = AA = A$?

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Accepted Answers:

No

Hence, $\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ cannot represent a matrix corresponding to some projection.

[Check Answers](#)

Your score is: 0/7