This assignment will not be graded and is only for practice.

Key points: 1

Definition of linear dependence: A set of vectors $v_1, v_2, ...v_n \in V$ is said to be linearly dependent if there exist scalars $a_1, a_2, ...a_n \in \mathrm{R}$, not all zero such that $a_1v_1 + a_2v_2 + ... + a_nv_n = 0$.
To check the vectors $v_1, v_2, ...v_n \in \mathrm{R}^m$ (with usual addition and scalar multiplication) are linearly dependent, we have to verify that the homogeneous system of linear equations $Ax = 0$ has infinitely many solutions, where the jth column of A is v_j.
Definition of linear independence: A set of vectors $v_1, v_2, ...v_n \in V$ is said to be linearly independent if $a_1v_1 + a_2v_2 + ... + a_nv_n = 0$ implies that $a_i = 0$ for $i = 1, 2, ...n$.
To check the vectors $v_1, v_2, ...v_n \in \mathrm{R}^m$ (with usual addition and scalar multiplication) are linearly independent, we have to verify that the homogeneous system of linear equations $Ax = 0$ has only the trivial solution (i.e., $x = 0$), where the jth column of A is v_j.
If S is a set containing zero vector, then the set S is a linearly dependent set.
Let S be a set of vectors from a vector space V. If an element of S is a scalar multiple of another element from the set S, then S is a linearly dependent set.

1) Vectors (2,1) and (6, 3) from R^2 are ***1 point***

- ◯ linearly independent.
- ◯ linearly dependent.
- ◯ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

linearly dependent.

2) Suppose the set $\{(3, 7, 4), v\}$ is a linearly dependent set in the vector space R^3 with usual addition and scalar multiplication. Which of the following vectors are possible candidates for v? ***1 point***

- ☐ $(0, 0, 0)$
- ☐ $(5, \frac{35}{3}, \frac{20}{3})$
- ☐ $(1, 0, 0)$
- ☐ $(6, 14, -8)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$(0, 0, 0)$
$(5, \frac{35}{3}, \frac{20}{3})$

3) The set of vectors $\{(1, 1, -1), (-1, 1, 1), \text{ and } (0, \frac{1}{2}, 0)\}$ in R^3 is **1 point**

- ◯ linearly independent.
- ◯ linearly dependent.
- ◯ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

linearly dependent.

4) $M_{2\times2}(\mathrm{R})$ denotes the vector space consisting of real square matrices of order 2 with usual matrix addition and scalar multiplication. Consider the following matrices: **1 point**

$$M_1 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, M_2 = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}, M_3 = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$$

Which of the following options is(are) true?

[**Hint:** Form a system of linear equations using matrix equation $x\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} + y\begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = 0$]

- ☐ $\{M_1, M_2\}$ is a linearly independent set.
- ☐ $\{M_1, M_2, M_3\}$ is a linearly dependent set.
- ☐ $\{M_2, M_3\}$ is a linearly dependent set.
- ☐ $\{M_1, M_3\}$ is a linearly independent set.

No, the answer is incorrect.

Score: 0

Accepted Answers:

$\{M_1, M_2\}$ is a linearly independent set.
$\{M_1, M_3\}$ is a linearly independent set.

Key points: 2

Given two vectors $(x_1, y_1),$ and $(x_2, y_2) \in \mathrm{R}^2$ construct $A = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix}$. Then these vectors are linearly dependent if and only if $det(A) = 0$ and linearly independent if and only if $det(A) = 0$.

Given three vectors $(x_1, y_1, z_1), (x_2, y_2, z_2)$ and $(x_3, y_3, z_3) \in \mathrm{R}^3$ construct $A = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$. Then these vectors are linearly dependent if and only if $det(A) = 0$ and linearly independent if and only if $det(A) = 0$

5) Vectors (2,3) and (4, 5) from R^2 are ***1 point***

[**Hint:** Form the matrix using the vectors and calculate the determinant of the matrix.]

- ◯ linearly independent.
- ◯ linearly dependent.
- ◯ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

linearly independent.

6) Consider a set of vectors $S = \{(1, 2), (-1, 2), (3, 1), (-3, 1)\} \subset \mathrm{R}^2$. Choose the set of correct options. ***1 point***

[**Hint:** Form the matrix using the vectors as the columns for each option and calculate the determinant of the matrix.]

- ☐ $\{(1, 2), (-1, 2)\}$ is a linearly dependent set.

- [ ] $\{(-1, 2), (-3, 1)\}$ is a linearly independent set.
- [ ] $\{(-3, 1), (-1, 2)\}$ is a linearly independent set.
- [ ] $\{(3, 1), (-3, 1)\}$ is a linearly dependent set.
- [ ] $\{(3, 1), (1, 2)\}$ is a linearly independent set.

No, the answer is incorrect.

Score: 0

Accepted Answers:

$\{(-1, 2), (-3, 1)\}$ is a linearly independent set.
$\{(-3, 1), (-1, 2)\}$ is a linearly independent set.
$\{(3, 1), (1, 2)\}$ is a linearly independent set.

7) The set of vectors $\{(1, 1, 0), (0, 1, 1), (1, 0, 1)\}$ is ***1 point***

[**Hint:** Form the matrix using the vectors as the columns and calculate the determinant of the matrix.]

- ◯ linearly dependent.
- ◯ linearly independent.
- ◯ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

linearly independent.

8) Suppose $\{(2, 3, 0), (0, 1, 0), v\}$ is a linearly dependent set in the vector space R^3 with usual addition and scalar multiplication. Which of the following vectors are possible candidates for v? ***1 point***

- [ ] $(2, 3, 1)$
- [ ] $(1, 2, 0)$
- [ ] $(1, 3, 0)$
- [ ] $(0, 1, 1)$

☐ $(\pi, e, 0)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$(1, 2, 0)$
$(1, 3, 0)$
$(\pi, e, 0)$

9) Consider three vectors $v_1 = (1, 0, 0), v_2 = (1, 1, 0), v_3 = (1, 1, 1)$ in the vector space R^3, with usual addition and scalar multiplication. Which of the following option is true? ***1 point***

[**Hint:** Find out the vectors of the given sets in the options and form the matrices using the vectors as the columns and calculate the determinant of the matrices.]

○ $\{v_1, v_2, v_3\}$ is a linearly dependent set.

○ $\{v_1 + v_2, v_1 + v_3, v_3\}$ is a linearly dependent set.

○ $\{v_2 - v_1, v_1, v_2\}$ is a linearly dependent set.

No, the answer is incorrect.

Score: 0

Accepted Answers:

$\{v_2 - v_1, v_1, v_2\}$ is a linearly dependent set.

Key points: 3

Any set of n vectors in R^2 with $n \geq 3$ is linearly dependent.
Any set of n vectors in R^3 with $n \geq 4$ is linearly dependent.
Any set of $n + 1$ vectors in R^n is linearly dependent.
Any set containing a linearly dependent set is linearly dependent.
If a set is linearly independent, then every subset of it, is linearly independent.

10) The subset $S = \{(3, 1), (1, 2), (-3, 1)\}$ of R^2 is ***1 point***

[**Hint:** Any set of n vectors in R^2 with $n \geq 3$ is linearly dependent.]

- ◯ linearly dependent.
- ◯ linearly independent.
- ◯ None of the above.

No, the answer is incorrect.
Score: 0

Accepted Answers:

linearly dependent.

11) Set $S = \{(1, 1, -1), (-1, 1, 1), (0, \frac{1}{2}, 0), (0, 1, -2), (1, 0, -2)\}$ subset of R^3 is ***1 point***

[**Hint:** Any set of n vectors in R^3 with $n \geq 4$ are linearly dependent.]

- ◯ linearly dependent.
- ◯ linearly independent.
- ◯ None of the above.

No, the answer is incorrect.
Score: 0

Accepted Answers:

linearly dependent.

12) Let S be a linearly dependent subset of R^3. Which of the following options are true? ***1 point***

- ☐ $S \cup \{(1, 0, 0)\}$ must be linearly dependent.
- ☐ $S \cup \{v\}$ must be linearly dependent for any $v \in \mathrm{R}^3$.
- ☐ $S \setminus \{v\}$ must be linearly dependent for any $v \in S$.
- ☐ There may exist some $v \in S$, such that $S \setminus \{v\}$ is still linearly dependent.

No, the answer is incorrect.
Score: 0

Accepted Answers:

$S \cup \{(1, 0, 0)\}$ must be linearly dependent.
$S \cup \{v\}$ must be linearly dependent for any $v \in \mathrm{R}^3$.
There may exist some $v \in S$, such that $S \setminus \{v\}$ is still linearly dependent.

Your score is: 0/12