



## AQ9.2: Activity Questions 2 - Not Graded



This assignment will not be graded and is only for practice.

### Partial Derivatives:

#### Level 1:

What is the rate of change of  $f(x, y, z) = 5x^2 - 6xy + 3z^2 - 2yz$  at  $(1, 2, -1)$  with respect to  $z$ ?

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:

(Type: Numeric) -10

1 point

1 point

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined as  $f(x, y) = x^3 + 3xy$ . Identify the correct option with respect to the partial derivatives of  $f$ .

☐

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} \quad \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}$$

☐

$$\frac{\partial f}{\partial x}(0, 0) = \frac{\partial f}{\partial y}(0, 0) \quad \frac{\partial^2 f}{\partial x \partial y}(0, 0) = \frac{\partial^2 f}{\partial y \partial x}(0, 0)$$

☐

$$\frac{\partial f}{\partial x}(1, 1) = \frac{\partial f}{\partial y}(1, 1) \quad \frac{\partial^2 f}{\partial x \partial y}(1, 1) = \frac{\partial^2 f}{\partial y \partial x}(1, 1)$$

☐

$$\frac{\partial f}{\partial x}(1, 0) = \frac{\partial f}{\partial y}(1, 0) \quad \frac{\partial^2 f}{\partial x \partial y}(1, 0) = \frac{\partial^2 f}{\partial y \partial x}(1, 0)$$

☐

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} + 3 \quad \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x} + 3$$

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:

$$\frac{\partial f}{\partial x}(0, 0) = \frac{\partial f}{\partial y}(0, 0) \quad \frac{\partial^2 f}{\partial x \partial y}(0, 0) = \frac{\partial^2 f}{\partial y \partial x}(0, 0)$$

$$\frac{\partial f}{\partial x}(1, 0) = \frac{\partial f}{\partial y}(1, 0) \quad \frac{\partial^2 f}{\partial x \partial y}(1, 0) = \frac{\partial^2 f}{\partial y \partial x}(1, 0)$$

1 point

Let  $f(x, y) = \frac{xy}{x^2 + y}$ . Identify the correct options with respect to the partial derivatives of  $f$ .

☐

$$\frac{\partial f}{\partial x} \text{ and } \frac{\partial f}{\partial y} \text{ are not defined at } (0, 0)$$

☐

$$\frac{\partial f}{\partial x}(1, 1) = 0, \frac{\partial f}{\partial y}(1, 1) = 1/4 \quad \frac{\partial^2 f}{\partial x \partial y}(1, 1) = 0, \frac{\partial^2 f}{\partial y \partial x}(1, 1) = 1/4$$

☐

$$\frac{\partial f}{\partial x}(1, 1) = 0, \frac{\partial f}{\partial y}(1, 1) = 0 \quad \frac{\partial^2 f}{\partial x \partial y}(1, 1) = 0, \frac{\partial^2 f}{\partial y \partial x}(1, 1) = 0$$

☐

$$\frac{\partial f}{\partial x}(1, 0) = 0, \frac{\partial f}{\partial y}(1, 0) = 1 \quad \frac{\partial^2 f}{\partial x \partial y}(1, 0) = 0, \frac{\partial^2 f}{\partial y \partial x}(1, 0) = 1$$

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:



$\frac{\partial f}{\partial x} \partial x \partial f$  and  $\frac{\partial f}{\partial y} \partial y \partial f$  are not defined at  $(0,0)$

$$\frac{\partial f}{\partial x}(1,1) = 0, \frac{\partial f}{\partial y}(1,1) = 1/4, \partial x \partial f(1,1) = 0, \partial y \partial f(1,1) = 1/4$$

$$\frac{\partial f}{\partial x}(1,0) = 0, \frac{\partial f}{\partial y}(1,0) = 1, \partial x \partial f(1,0) = 0, \partial y \partial f(1,0) = 1$$

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined as  $f(x, y) = x^2 + 4xy - y^2$ . Suppose  $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} \partial x \partial f = \partial y \partial f$  at  $(12, k)$ . Find the value of  $k$

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

(Type: Numeric) 4

1 point

**Level 2:**

1 point

Suppose  $f(x, y, z) = xy + yz + zx - 3xyz$  is a multivariable function defined on domain  $D \subset \mathbb{R}^3$ . Which of the following is(are) correct?

☐

$f$  is a scalar valued multivariable function.

☐

The rate of change of the function  $f$  at  $(0,1,1)$  with respect to  $x$  is 1.

☐

The rate of change of the function  $f$  at  $(2,2,2)$  with respect to  $x$  is same as the rate of change of the function  $f$  at  $(2,2,2)$  with respect to  $z$ .

☐

The rate of change of the function  $f$  at  $(1,3,5)$  with respect to  $y$  is -9.

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$f$  is a scalar valued multivariable function.

The rate of change of the function  $f$  at  $(2,2,2)$  with respect to  $x$  is same as the rate of change of the function  $f$  at  $(2,2,2)$  with respect to  $z$ .

The rate of change of the function  $f$  at  $(1,3,5)$  with respect to  $y$  is -9.

1 point

Suppose  $f: D \rightarrow \mathbb{R}$  is a multivariable function defined on domain  $D \subset \mathbb{R}^2$  and also satisfying the following conditions:

$$(i) \frac{\partial f}{\partial x} = 6x + 4xy + 3y, \partial x \partial f = 6x + 4xy + 3y$$

$$(ii) \frac{\partial f}{\partial y} = 2x^2 + 3x - 3y, \partial y \partial f = 2x^2 + 3x - 3y$$

Which of the following functions can not be  $f$ ?

☐

$$f(x, y) = 3x^2 + 2x^2y + 3xy - \frac{3}{2}y^2, f(x, y) = 3x^2 + 2x^2y + 3xy - 23y^2$$

☐

$$f(x, y) = 3x^2 + 2x^2y - 3xy - \frac{3}{2}y^2, f(x, y) = 3x^2 + 2x^2y - 3xy - 23y^2$$

☐

$$f(x, y) = 3x^2 - 3xy - 2x^2y + 3y^2, f(x, y) = 3x^2 - 3xy - 2x^2y + 3y^2$$

☐

$$f(x, y) = 3x^2 + 3y^2 + 3xy, f(x, y) = 3x^2 + 3y^2 + 3xy$$

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$$f(x, y) = 3x^2 + 2x^2y - 3xy - \frac{3}{2}y^2 \quad f(x, y) = 3x^2 + 2x^2y - 3xy - 23y^2$$

$$f(x, y) = 3x^2 - 3xy - 2x^2y + 3y^2 \quad f(x, y) = 3x^2 - 3xy - 2x^2y + 3y^2$$

$$f(x, y) = 3x^2 + 3y^2 + 3xy \quad f(x, y) = 3x^2 + 3y^2 + 3xy$$

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be given by  $f(x, y) = (P(x, y), Q(x, y))$  where  $P(x, y) = 3x^2y$ ,  $Q(x, y) = 5x + y^3$ . Consider a matrix

$A = \begin{bmatrix} \frac{\partial P}{\partial x} & \frac{\partial P}{\partial y} \\ \frac{\partial Q}{\partial x} & \frac{\partial Q}{\partial y} \end{bmatrix}$ . If the determinant of  $A$  is  $axy^3 - bx^2$  where

$a, b \in \mathbb{R}$ , then find the value of  $a - b$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

(Type: Numeric) 3

1 point

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be given by  $f(x, y) = (P(x, y), Q(x, y))$  where  $P(x, y) = 6x + y^2$ ,  $Q(x, y) = 3x^2 + 2y$ . Consider a

matrix  $A = \begin{bmatrix} \frac{\partial P}{\partial x} & \frac{\partial P}{\partial y} \\ \frac{\partial Q}{\partial x} & \frac{\partial Q}{\partial y} \end{bmatrix}$ . If  $\det(A) = 0$ , then find the value of  $xy$

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

(Type: Numeric) 1

1 point

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined as  $f(x, y) = e^{4(x+y)} \sin(7x - 4y)$

If  $4\frac{\partial f}{\partial x} + 7\frac{\partial f}{\partial y} = k \cdot f(x, y)$  where  $k \in \mathbb{R}$ , then find the value of  $k$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

(Type: Numeric) 44

1 point

[Check Answers](#)

Your score is: 0/9