

This assignment will not be graded and is only for practice.

Level 1

1) Consider the following expression: $v_1 = (1, 0)$, $v_2 = (0, 0)$, $v_3 = (0, 1)$, $v_5 = 1$, $v_6 = 5$, $v_7 = (1, 0, 1)$, $v_8 = (0, 0, 1)$, $v_9 = (0, 0, 0)$, $v_{10} = (0, 1, 0)$. Choose the set of correct options. **1 point**

- ☐ v_1, v_7 represent vectors in $\mathbb{R}^2$.
- ☐ v_2, v_5 represent vectors in $\mathbb{R}^1$.
- ☐ v_2, v_5, v_9 represent vectors in $\mathbb{R}^3$.
- ☐ v_1, v_2, v_3 represent vectors in $\mathbb{R}^2$.
- ☐ v_5, v_6 represent vectors in $\mathbb{R}^1$.
- ☐ v_1, v_6 represent vectors in $\mathbb{R}^1$.
- ☐ v_8, v_9, v_{10} represent vectors in $\mathbb{R}^3$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

v_1, v_2, v_3 represent vectors in $\mathbb{R}^2$.
 v_5, v_6 represent vectors in $\mathbb{R}^1$.
 v_8, v_9, v_{10} represent vectors in $\mathbb{R}^3$.

2) Let v, v_1, v_2 represent vectors in a vector space V over $\mathbb{R}$. Which of the following expressions make sense and represent vectors in V ? **1 point**

- ☐ $cv \in V$, where $c \in \mathbb{R}$.
- ☐ $\frac{v}{2}$.
- ☐ v^2 .
- ☐ $v_1 - v_2$.
- ☐ $v_1^2 - v_2^2$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

$cv \in V$, where $c \in \mathbb{R}$.

$\frac{v}{2}$.

$v_1 - v_2$.

3) Consider the set $V = \{(z - x, -y) \mid x, y, z \in \mathbb{R}\} \subseteq \mathbb{R}^2$ with the usual addition and scalar multiplication as in $\mathbb{R}^2$ and associated statements given below. Choose the correct options. **1 point**

- ☐ V is closed under addition i.e $(v_1, v_2 \in V \implies v_1 + v_2 \in V)$.
- ☐ V is closed under scalar multiplication i.e $(c \in \mathbb{R}, v \in V \implies cv \in V)$.
- ☐ V has a zero element with respect to addition. i.e., there exists some element 0 such that $v + 0 = v$, for all $v \in V$.
- ☐ V is not a vector space.
- ☐ $(a + b)v = av + bv$ where $a, b \in \mathbb{R}$ and $v \in V$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

V is closed under addition i.e $(v_1, v_2 \in V \implies v_1 + v_2 \in V)$.

V is closed under scalar multiplication i.e $(c \in \mathbb{R}, v \in V \implies cv \in V)$.

V has a zero element with respect to addition. i.e., there exists some element 0 such that $v + 0 = v$, for all $v \in V$.

$(a + b)v = av + bv$ where $a, b \in \mathbb{R}$ and $v \in V$.

4) Consider a set $V = \{(1, x) \mid x \in \mathbb{R}\} \subseteq \mathbb{R}^2$ with the usual addition and scalar multiplication as in $\mathbb{R}^2$ and associated statements given below. Choose the correct options. **1 point**

- ☐ V is closed under addition.
- ☐ V has no zero element with respect to addition. i.e., there does not exists any element 0 such that $v + 0 = v$, for all $v \in V$.
- ☐ $(a + b)v = av + bv$ where $a, b \in \mathbb{R}$ and $v \in V$.

☐ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

V has no zero element with respect to addition. i.e., there does not exist any element 0 such that $v + 0 = v$, for all $v \in V$.

$(a + b)v = av + bv$ where $a, b \in \mathbb{R}$ and $v \in V$.

5) Which of the following sets with the given addition and scalar multiplication (scalars are real **1 point** numbers in every case) form vector spaces?

$$V_1 = \{(x, y, z) \mid x, y, z \in \mathbb{R}\}$$

$$\text{Addition: } (x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, y_1 + 1, z_2 + 1);$$

$$(x_1, y_1, z_1), (x_2, y_2, z_2) \in V_1$$

$$\text{Scalar multiplication: } c(x, y, z) = (x, y, cz); (x, y, z) \in V_1, c \in \mathbb{R}$$

$$V_2 = \{(x, y, z) \mid x, y, z \in \mathbb{R}\}$$

$$\text{Addition: } (x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1, y_1, z_1);$$

$$(x_1, y_1, z_1), (x_2, y_2, z_2) \in V_2$$

$$\text{Scalar multiplication: } c(x, y, z) = (x, cy, z); (x, y, z) \in V_2, c \in \mathbb{R}$$

$$V_3 = \{(x, y, z) \mid x, y, z \in \mathbb{R}\}$$

$$\text{Addition: } (x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, 0, 0);$$

$$(x_1, y_1, z_1), (x_2, y_2, z_2) \in V_3$$

$$\text{Scalar multiplication: } c(x, y, z) = (cx, y, z); (x, y, z) \in V_3, c \in \mathbb{R}$$

Choose the correct options.

☐ V_3 is not vector space.

☐ V_2 is vector spaces.

☐ All are vector spaces.

☐ V_1 is not vector space.

No, the answer is incorrect.

Score: 0

Accepted Answers:

V_3 is not vector space.

V_1 is not vector space.

6) If (a, b, c) is the inverse element of $(1, 2, 5) \in \mathbb{R}^3$ of the vector space $\mathbb{R}^3$, then find the value of $a - b + c$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) -4

1 point

Level 2

7) Consider the following sets:

1 point

$V_1 = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is a symmetric matrix, i.e., } A = A^T\}$

$V_2 = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is a scalar matrix}\}$

$V_3 = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is a diagonal matrix}\}$

$V_4 = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is an upper triangular matrix}\}$

$V_5 = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is a lower triangular matrix}\}$

Choose the set of correct options.

- ☐ Only V_1 is a subspace of $M_{2 \times 2}(\mathbb{R})$.
- ☐ Only V_4 is a subspace of $M_{2 \times 2}(\mathbb{R})$.
- ☐ Both V_2 and V_3 are subspaces of $M_{2 \times 2}(\mathbb{R})$
- ☐ All are subspaces of $M_{2 \times 2}(\mathbb{R})$

No, the answer is incorrect.

Score: 0

Accepted Answers:

Both V_2 and V_3 are subspaces of $M_{2 \times 2}(\mathbb{R})$

All are subspaces of $M_{2 \times 2}(\mathbb{R})$

8) Choose the set of correct options.

1 point

- ☐ A vector in $\mathbb{R}^3$ can be thought of as matrix of order 1×3 or 3×1 .
- ☐ Intersection of any two subspaces of a vector space V is not a subspace.
- ☐ Intersection of any two subsets of a vector space V is a subspace.
- ☐ The empty subset of a vector space V is a subspace of V .
- ☐ $V = \{(x, 0, 0) \mid x \in \mathbb{R}\}$ is a subspace of $\mathbb{R}^3$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

A vector in $\mathbb{R}^3$ can be thought of as matrix of order 1×3 or 3×1 . $V = \{(x, 0, 0) \mid x \in \mathbb{R}\}$ is a subspace of $\mathbb{R}^3$.9) Consider a set $V = \{(x, y) \mid x, y \in \mathbb{R}\} \subseteq \mathbb{R}^2$ with the usual addition as in $\mathbb{R}^2$ and scalar multiplication is defined as**1 point**

$$c(x, y) = \begin{cases} (0, 0) & c = 0 \\ (\frac{cx}{2}, \frac{y}{c}) & c \neq 0 \end{cases} \quad (x, y) \in V, c \in \mathbb{R}$$

Consider the statements given below.

P: V is closed under addition.**Q:** V has zero element with respect to addition. i.e., there exists some element 0 such that $v + 0 = v$, for all $v \in V$.**R:** $1 \cdot v = v$ where $1 \in \mathbb{R}$ and $v \in V$.**S:** $a(v_1 + v_2) = av_1 + av_2$ where $v_1, v_2 \in V$ and $a \in \mathbb{R}$.**T:** $(a + b)v = av + bv$ where $a, b \in \mathbb{R}$ and $v \in V$.

Choose the set of correct options.

- ☐ Only P is true.
- ☐ Only Q is true.
- ☐ Both P and Q are true.

- ☐ P,Q and S are true.
- ☐ Both R and T are not true.

No, the answer is incorrect.

Score: 0

Accepted Answers:

Both P and Q are true.

P,Q and S are true.

Both R and T are not true.

10) Consider the following statements

Statement P: $W = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A^2 = A\}$, a subset of $M_{2 \times 2}(\mathbb{R})$ with usual addition and scalar multiplication as in $M_{2 \times 2}(\mathbb{R})$, is a subspace of $M_{2 \times 2}(\mathbb{R})$.

Statement Q: $W = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } \det(A) = 0\}$, a subset of $M_{2 \times 2}(\mathbb{R})$ with usual addition and scalar multiplication as in $M_{2 \times 2}(\mathbb{R})$, is a subspace of $M_{2 \times 2}(\mathbb{R})$.

Statement R: Consider a system of linear equations $Ax^T = b$, where A is invertible square matrix of order 2, $x = (x_1, x_2)$ and $b = 0$. Let $W = \{x = (x_1, x_2) \mid x \in \mathbb{R}^2 \text{ and } Ax^T = b\}$ is a subset of $\mathbb{R}^2$ with usual addition and scalar multiplication as in $\mathbb{R}^2$, is a subspace of $\mathbb{R}^2$.

Statement S: Consider a system of linear equations $Ax^T = 0$, where A is invertible square matrix of order 2 and $x = (x_1, x_2)$. Let $W = \{x = (x_1, x_2) \mid x \in \mathbb{R}^2 \text{ and } Ax^T = 0\}$ is a subset of $\mathbb{R}^2$ with usual addition and scalar multiplication as in $\mathbb{R}^2$, is a subspace of $\mathbb{R}^2$.

Find the number of correct statements

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 1

1 point

Your score is: 0/10