

The due date for submitting this assignment has passed.

Due on 2026-04-08, 23:59 IST.

You may submit any number of times before the due date. The final submission will be considered for grading.

You have last submitted on: 2026-04-08, 08:45 IST

Note: This assignment will be evaluated after the deadline passes. You will get your score 48 hrs after the deadline. Until then the score will be shown as Zero.

Multiple Select Questions (MSQ):

1) An inner product on $\mathbb{R}^3$ is defined as:

1 point

$$\langle \cdot, \cdot \rangle: \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$$

$$\langle (x_1, x_2, x_3), (y_1, y_2, y_3) \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3.$$

Match the sets of vectors in column A with their properties of orthogonality or orthonormality in column B with respect to the above inner product.

	Set of vectors (Column A)		Properties (Column B)
a)	$\{(2, 3, 4), (-1, 2, -1)\}$	i)	Forms a basis but not orthogonal
b)	$\{\frac{1}{\sqrt{2}}(1, 0, -1), \frac{1}{\sqrt{2}}(-1, 0, -1)\}$	ii)	Forms an orthogonal basis
c)	$\{(2, 3, 4), (-1, 2, -1), (0, 4, -3)\}$	iii)	Orthogonal but not orthonormal, and does not form a basis of $\mathbb{R}^3$
d)	$\{(2, 3, 4), (-1, 2, -1), (11, 2, -7)\}$	iv)	Orthonormal, but does not form a basis of $\mathbb{R}^3$

Table : M2W8G1

Choose the set of correct options.

- ☐ a $\rightarrow$ iv)
- ☒ a $\rightarrow$ iii)
- ☒ b $\rightarrow$ iv)
- ☐ b $\rightarrow$ iii)

☐ c $\rightarrow$ ii)☒ c $\rightarrow$ i)☐ d $\rightarrow$ i)☒ d $\rightarrow$ ii)

Yes, the answer is correct.

Score: 1

Accepted Answers:

a $\rightarrow$ iii)

b $\rightarrow$ iv)

c $\rightarrow$ i)

d $\rightarrow$ ii)

2) Choose the set of correct options.

1 point

☒ Suppose $\beta = \{v_1, v_2, \dots, v_n\}$ is an orthogonal basis of an inner product space V . If there exists some $v \in V$, such that $\langle v, v_i \rangle = 0$ for all $i = 1, 2, \dots, n$, then $v = 0$.

☒ There exists an orthonormal basis for $\mathbb{R}^n$ with the standard inner product.

☒ If P_W denotes the linear transformation which projects the vectors of an inner product space V to a subspace W of V , then $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$, where 0 denotes the zero vector of V .

☒ $\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ cannot represent a matrix corresponding to some projection.

Yes, the answer is correct.

Score: 1

Accepted Answers:

Suppose $\beta = \{v_1, v_2, \dots, v_n\}$ is an orthogonal basis of an inner product space V . If there exists some $v \in V$, such that $\langle v, v_i \rangle = 0$ for all $i = 1, 2, \dots, n$, then $v = 0$.

There exists an orthonormal basis for $\mathbb{R}^n$ with the standard inner product.

If P_W denotes the linear transformation which projects the vectors of an inner product space V to a subspace W of V , then $\text{range}(P_W) \cap \text{null space}(P_W) = \{0\}$, where 0 denotes the

zero vector of V .

$\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ cannot represent a matrix corresponding to some projection.

Numerical Answer Type (NAT):

3) If A is an orthogonal matrix of order 5, then find nullity of the matrix A .

0

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 0

1 point

4) Let $v \in \mathbb{R}^3$ be a vector such that $\|v\| = 5$. If u is the vector obtained from v after the anti-clockwise rotation of XY-plane with angle 70° about the Z-axis, then find the length of the vector u .

5

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 5

1 point

5) Let $v = (1, 2, 2)$ be a vector in $\mathbb{R}^3$. If (a, b, c) is the vector obtained from v after the anti-clockwise rotation of YZ-plane with angle 60° about the X-axis, then find the value of $a + b + c$.

3

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 3

1 point

6) Consider a vector space $M_{2 \times 2}(\mathbb{R})$ and a norm on the vector space defined as

$$\|A\| = \max\{|a_{11}| + |a_{21}|, |a_{12}| + |a_{22}|\}, \text{ where } A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in M_{2 \times 2}(\mathbb{R}).$$

Let $B = \begin{bmatrix} x & \sqrt{2}x \\ -\sqrt{2}y & y \end{bmatrix}$ be an orthogonal matrix i.e. $BB^T = B^TB = I$ and assume $x, y > 0$.

Then find the norm of the matrix $C = \begin{bmatrix} \sqrt{3}x & \sqrt{3}x \\ \sqrt{3}y & \sqrt{3}y \end{bmatrix}$ (i.e., $\|C\|$)

2

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 2

1 point

7) Let $V = \mathbb{R}^2$ be the inner product space with usual inner product and a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined as $T(x, y) = (\frac{a}{\sqrt{a^2+2^2}}x + \frac{3}{\sqrt{b^2+3^2}}y, \frac{2}{\sqrt{a^2+2^2}}x + \frac{b}{\sqrt{b^2+3^2}}y)$. If T is an orthogonal linear transformation, then find the value of $3a + 2b$.

0

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 0

1 point

Comprehension Type Question:

With a particular frame of reference (in $\mathbb{R}^3$), position of a target is given as the vector $(3, 4, 5)$. Three shooters S_1, S_2 , and S_3 are moving along the lines $x = y, x = -y$, and $x = 2y$, on the XY -plane (i.e., $z = 0$) to shoot the target. Suppose that, there is another shooter S_4 , who is moving on the plane $x + y + z = 0$. Suppose all of them shoot the target so that the target is at the closest distance from the respective path or plane on which they are travelling. Answer questions 10, 11 and 12 using the given information.

8) Choose the set of correct options.

1 point

- ☐ S_1 will shoot the target from the point $(\frac{7}{2}, -\frac{7}{2}, 0)$.
- ☒ S_1 will shoot the target from the point $(\frac{7}{2}, \frac{7}{2}, 0)$.
- ☐ S_1 will shoot the target from the point $(1, 1, 0)$.
- ☐ S_2 will shoot the target from the point $(-\frac{1}{2}, -\frac{1}{2}, 0)$.
- ☐ S_2 will shoot the target from the point $(1, -1, 0)$.
- ☒ S_2 will shoot the target from the point $(-\frac{1}{2}, \frac{1}{2}, 0)$.
- ☒ S_3 will shoot the target from the point $(4, 2, 0)$.
- ☐ S_3 will shoot the target from the point $(2, 1, 0)$.
- ☐ S_3 will shoot the target from the point $(0, 0, 0)$.

Yes, the answer is correct.

Score: 1

Accepted Answers:

S_1 will shoot the target from the point $(\frac{7}{2}, \frac{7}{2}, 0)$.
 S_2 will shoot the target from the point $(-\frac{1}{2}, \frac{1}{2}, 0)$.
 S_3 will shoot the target from the point $(4, 2, 0)$.

9) If (a, b, c) is the point from which the shooter S_4 will shoot the target, then find the value of $a + 2b + 3c$.

2

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 2

1 point

10) Let d_i be the distance of the target from the point where the shooter S_i shoots the target, for $i = 1, 2, 3, 4$ and let d be the minimum amongst the d_i . Find the value of d^2 .

25.5

Yes, the answer is correct.

Score: 1

Accepted Answers:

(Type: Numeric) 25.5

1 point