



Solve with us 4 - Not Graded



This assignment will not be graded and is only for practice.

Key points: 1

- The span of a set S (of vectors) is defined as the set of all finite linear combinations of elements (vectors) of S and denoted by $\text{Span}(S)$ i.e.,

$$\text{Span}(S) = \{ \sum_{i=1}^n a_i v_i \in V \mid a_1, a_2, \dots, a_n \in R \text{ and } v_1, v_2, \dots, v_n \in S \}$$

$$\text{Span}(S) = \{ \sum_{i=1}^n a_i v_i \in V \mid a_1, a_2, \dots, a_n \in R \text{ and } v_1, v_2, \dots, v_n \in S \}.$$
- Let V be a vector space. A set $S \subseteq V$ is spanning set for V if $\text{Span}(S) = V$.
- Algorithm to obtain a spanning set for a non-zero vector space V :
 Algorithm to obtain a spanning set for a non-zero vector space V :
 - Start with the empty set $S_0 = \emptyset$.
 - Add any non-zero vector $v_1 \in V$ to S_0 and define it as S_1 . So $S_1 = S_0 \cup \{v_1\}$.
 - If $\text{Span}(S_1) = V$, then we are done. Otherwise choose $v_2 \in V \setminus \text{span}(S_1)$ and add it to the set and call it S_2 . So $S_2 = S_1 \cup \{v_2\}$.
 - If $\text{Span}(S_2) = V$, then we are done. Otherwise, choose $v_3 \in V \setminus \text{Span}(S_2)$, and construct $S_3 = S_2 \cup \{v_3\}$.
 - Repeat the process until $\text{Span}(S_n) = V$ for some n .

1 point

Let $S = \{(1, 2), (2, 3), (1, -1)\} \subseteq R^2$. Which of the following is (are) elements of $\text{Span}(S)$?

☐ $5(1, 2) + 8(1, -1)$

☐ $(0, 0)$



$2(1, 2) - 3(2, 3) + 5(1, -1)$



$(2, 1) - 2(3, 2) - 4(-1, 1)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$5(1, 2) + 8(1, -1)$

$(0, 0)$

$2(1, 2) - 3(2, 3) + 5(1, -1)$

$(2, 1) - 2(3, 2) - 4(-1, 1)$

1 point

Let $S = \{(1, 1, -1), (-1, 1, 1)\} \subset R^3$. Which of the following is (are) elements of $\text{Span}(S)$?

☐ $2(1, 1, -1) + (-1, 1, 1)$

☐ $(8, 8, -8)$

☐ $(0, 2, 0)$

☐ $(1, 1, -2)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$2(1, 1, -1) + (-1, 1, 1)$



(8, 8, -8)

(0, 2, 0)

1 point

Let $S = \{(-1, 1, 1), (0, \frac{1}{2}, 0), (0, 1, -2)\} \subset R^3 \subset R^3$. Which of the following option(s) is (are) true?

[Hint: Form the matrix using the vectors as the columns and calculate the determinant of the matrix.]

☐

S is a linearly dependent set.

☐

S is a linearly independent set.

☐

S is a basis of the vector space R^3 .

☐ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

S is a linearly independent set.

S is a basis of the vector space R^3 .

1 point

Consider a set of vectors $S = \{(1, 2), (-1, 2), (3, 1), (-3, 1)\} \subset R^2$. Choose the set of correct options.

☐

The set $\{(1, 2), (-1, 2), (3, 1)\}$ is a basis of R^2 .

☐

The set $\{(-1, 2), (3, 1)\}$ is a basis of R^2 .

☐

The set $\{(3, 1), (-3, 1)\}$ is a basis of R^2 .

☐ None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

The set $\{(-1, 2), (3, 1)\}$ is a basis of R^2 .

The set $\{(3, 1), (-3, 1)\}$ is a basis of R^2 .

Key points: 2

- The following conditions are equivalent for a set to be a basis of a vector space:
 - Maximal linearly independent set : A linearly independent set such that if any vector is appended to it, the new set is no longer linearly independent.
 - Minimal spanning set : A spanning set such that if any vector is removed from the set, then the set will not be a spanning set.

1 point

Which of the following sets are maximal linearly independent sets in R^2 ?

☐

$\{(1, 0)\}$

☐

$\{(1, 2), (2, 4)\}$

☐ $\{(1, 2), (1, 1)\} \{(1, 2), (1, 1)\}$ ☐ $\{(1, 2), (1, 1), (2, 1)\} \{(1, 2), (1, 1), (2, 1)\}$ ☐ $\{(1, 0), (0, 1)\} \{(1, 0), (0, 1)\}$

No, the answer is incorrect.

Score: 0

Accepted Answers:

 $\{(1, 2), (1, 1)\} \{(1, 2), (1, 1)\}$ $\{(1, 0), (0, 1)\} \{(1, 0), (0, 1)\}$

1 point

Which of the following sets are minimal spanning sets of R^3 ?

☐ $\{(1, 0, 0), (0, 1, 0)\} \{(1, 0, 0), (0, 1, 0)\}$ ☐ $\{(1, 0, 0), (1, 1, 0), (1, 1, 1)\} \{(1, 0, 0), (1, 1, 0), (1, 1, 1)\}$ ☐ $\{(1, 0, 1), (0, 1, 1), (1, 1, 0)\} \{(1, 0, 1), (0, 1, 1), (1, 1, 0)\}$ ☐ $\{(1, 0, 1), (0, 1, 1), (1, 1, 2)\} \{(1, 0, 1), (0, 1, 1), (1, 1, 2)\}$ ☐ $\{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\} \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\}$

No, the answer is incorrect.

Score: 0

Accepted Answers:

 $\{(1, 0, 0), (1, 1, 0), (1, 1, 1)\} \{(1, 0, 0), (1, 1, 0), (1, 1, 1)\}$ $\{(1, 0, 1), (0, 1, 1), (1, 1, 0)\} \{(1, 0, 1), (0, 1, 1), (1, 1, 0)\}$

Key Points: 3**Key Points: 3**

- **Basis:** A basis of a vector space V is a linearly independent subset of V that spans V .

- Basis of a vector space:
 - Maximal linearly independent set.
 - Minimal spanning set.

- How to find a basis of a subspace of R^n ?

- Suppose a subspace V of R^n is defined by some constraints, as follows:

$$V = \{(x_1, x_2, \dots, x_n) \mid f_1(x_1, x_2, \dots, x_n) = 0, f_2(x_1, x_2, \dots, x_n) = 0, \dots, f_m(x_1, x_2, \dots, x_n) = 0\}$$

$$V = \{(x_1, x_2, \dots, x_n) \mid f_1(x_1, x_2, \dots, x_n) = 0, f_2(x_1, x_2, \dots, x_n) = 0, \dots, f_m(x_1, x_2, \dots, x_n) = 0\} \subseteq R^n$$

Where f_i fi's are linear equations in general.

- **Step 1:** By solving f_i fi's we have to try to find out the dependent and independent variables. Suppose there are k independent variables and there are $k - n$ dependent variables.

- **Step 2:** Choose the first independent variable, let it be x_1 and assign the value 1

to that, and assign 0 to all the other independent variables.

- **Step 3:** Find out the values of all the dependent variables with respect to the assignment given in Step 2 and call that vector to be v_1 .

- **Step 4:** Carry out this process for each independent variable, to get the vectors $v_2, \dots, v_k$.

The set of vectors $\{v_1, v_2, \dots, v_k\}$ is a linearly independent set and this is a maximal linearly independent set in V , hence forms a basis of V . Hence, dimension of V is k in this case.

1 point

Let $S = \{(-1, 1, 1), (0, \frac{1}{2}, 1), (0, 1, -2)\} \subset R^3$. Which of the following option(s) is (are) true?

Hint: Form the matrix using the vectors as the columns and calculate the determinant of the matrix.

☐

S is a linearly dependent set.

☐

S is a linearly independent set.

☐

S is a basis of the vector space R^3 .

☐

None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

S is a linearly independent set.

S is a basis of the vector space R^3 .

1 point

Consider a set of vectors $S = \{(1, 2), (-1, 2), (3, 1), (-3, 1)\}$ in R^2 . Choose the set of correct options.

☐

The set $\{(1, 2), (-1, 2), (3, 1)\}$ is a basis of R^2 .

☐

The set $\{(-1, 2), (3, 1)\}$ is a basis of R^2 .

☐

The set $\{(3, 1), (-3, 1)\}$ is a basis of R^2 .

☐

None of the above.

No, the answer is incorrect.

Score: 0

Accepted Answers:

The set $\{(-1, 2), (3, 1)\}$ is a basis of R^2 .

The set $\{(3, 1), (-3, 1)\}$ is a basis of R^2 .

1 point

Let V be a vector space which is defined as follows:

$V = \{(x, y, z) \mid 2x + 3y = 0 \text{ and } y + 5z = 0\} \subseteq R^3$ with usual addition and scalar multiplication. Which of the following sets form bases of V ? (More than one option may be correct)

[Hint: Two constraints are given here, $f_1(x, y, z) = 2x + 3y = 0$ and $f_2(x, y, z) = y + 5z = 0$. Observe that z is the independent variable and x and y are the dependent variables.]

☐
 $\{(2, 3, 0), (0, 1, 5)\} \{(2, 3, 0), (0, 1, 5)\}$
☐
 $\{(3, -2, 0), (0, 5, -1)\} \{(3, -2, 0), (0, 5, -1)\}$
☐
 $\{(\frac{15}{2}, -5, 1)\} \{(215, -5, 1)\}$
☐
 $\{(15, -10, 2)\} \{(15, -10, 2)\}$

No, the answer is incorrect.

Score: 0

Accepted Answers:

 $\{(\frac{15}{2}, -5, 1)\} \{(215, -5, 1)\}$
 $\{(15, -10, 2)\} \{(15, -10, 2)\}$

1 point

Let V be a vector space which is defined as follows: $V = \{(x, y, z, w) \mid x + z = y + w\} \subseteq \mathbb{R}^4$
 $V = \{(x, y, z, w) \mid x + z = y + w\} \subseteq \mathbb{R}^4$ with usual addition and scalar multiplication. Which of the following set forms a basis of V ?

[Hint: The constraint given here is $x + z = y + w$, which can be written as $x - y + z - w = 0$. As there are 4 variables and 1 equation, the number of independent variables is 3, where as the number of dependent variable is 1. If you check the first vector of each option, only Option 2 and Option 4 satisfying the constraint, and in that vector $x = y = 1, x = y = 1$. Start with taking y as the dependent variable, and apply the method mentioned above.]

☐
 $\{(1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 0, 1)\} \{(1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 0, 1)\}.$
☐
 $\{(1, 1, 0, 0), (0, 1, -1, 0), (0, -1, 0, 1)\} \{(1, 1, 0, 0), (0, 1, -1, 0), (0, -1, 0, 1)\}.$
☐
 $\{(1, -1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\} \{(1, -1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\}.$
☐
 $\{(1, 1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\} \{(1, 1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\}.$

No, the answer is incorrect.

Score: 0

Accepted Answers:

 $\{(1, 1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\} \{(1, 1, 0, 0), (0, 1, 1, 0), (0, -1, 0, 1)\}.$

Find the dimension of the vector space

 $V = \{(x, y, z, w) \mid x + y = z + w, x + w = y + z, \text{ and } x, y, z, w \in \mathbb{R}\}$
 $\{(x, y, z, w) \mid x + y = z + w, x + w = y + z, \text{ and } x, y, z, w \in \mathbb{R}\}.$ [Hint: Two constraints are given here]

 $f_1(x, y, z, w) = x + y - z - w = 0$ and $f_2(x, y, z, w) = x + y - z - w = 0$
 $f_2(x, y, z, w) = x - y - z + w = 0$ and $f_2(x, y, z, w) = x - y - z + w = 0$. Try to find out the number of independent variables.]

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 2

1 point

Key Points: 4

- How to find a basis of a subspace V of $M_{m \times n}(R)$:

- **Step 1:** Consider each element of a $m \times n$ matrix as a variable. So there are mn number of variables.

- **Step 2:** Using the constraints given as the definition of the subspace separate out the independent and the dependent variables. Suppose there are k independent variables and $mn - k$ dependent variables.

- **Step 3:** Define a matrix M_1 by assigning 1 to one independent variable and 0 to all the other independent variables, together with finding out the values of all the dependent variables.

- **Step 4:** Repeat this process for all the independent variables to get the vectors $M_2, M_3, \dots, M_k$.

The set of matrices $\{M_1, M_2, \dots, M_k\}$ forms a basis of V . Hence the dimension of V is k in this case.

1 point

Let $V = \{A \mid A \in M_{3 \times 3}(R), A \text{ is a diagonal matrix}\}$. Which of the following sets form a basis of V ?

[Hint: $a_{ij} = 0$ if $i \neq j$. Thus there are 6 constraints and these are the dependent variables while the diagonal entries are the 3 independent elements.]

☐

$$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$$

☐

$$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$$

☐

$$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right\}$$

☐

$$\left\{ \begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \right\}$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$\left\{ \begin{bmatrix} 100 \\ 000 \\ 000 \end{bmatrix}, \begin{bmatrix} 000 \\ 010 \\ 000 \end{bmatrix}, \begin{bmatrix} 000 \\ 000 \\ 001 \end{bmatrix} \right\} = \left\{ \begin{bmatrix} 100000000 \\ 000010000 \\ 000000001 \end{bmatrix} \right\}$$

Let $V = \{A \mid A \in M_{3 \times 3}(R), \text{sum of the diagonal elements are } 0\}$. What will be the dimension of V ?

[Hint: The general form a diagonal matrix of order 3 is $\begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_3 \end{bmatrix}$. Hence there are

three variables, a_1, a_2, a_3 . Now there is one more constraint, which says that the sum of the diagonal elements is 0. Hence we have $a_1 + a_2 + a_3 = 0$. So we can choose a_3 to be the dependent variable and a_1 and a_2 to be the independent variables, as $a_3 = -a_1 - a_2$.]

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 2

1 point

Key points: 5 Key points: 5

- Rank of a matrix:
 - Maximum number of linearly independent column vectors (or row vectors) in a matrix.

NOTE: Rank of a matrix is always less than or equal to the number of columns (or rows).

- How to find rank of a matrix A:
 - Reduce the matrix to row echelon form.
 - Find the number of non zero rows in the reduced matrix which will be the rank of the matrix A.

1 point

Consider the following two matrices:

$$P = \begin{bmatrix} 1 & 0 & 3 \\ 2 & -2 & 0 \\ 0 & 1 & -1 \\ 0 & 2 & 1 \end{bmatrix}; Q = \begin{bmatrix} 1 & 1 & -20 \\ 1 & 0 & 2 \\ 0 & -1 & 0 & 4 \end{bmatrix}$$

Which of the following statements is (are) TRUE? [Hint: Try to reduce the matrices to row echelon form and find the number of non zero rows]

- ☐ Rank(P) = 4
- ☐ Rank(P) = Rank(Q)
- ☐ Rank(QP) = 4
- ☐ Rank(PQ) = Rank(QP)

No, the answer is incorrect.

Score: 0

Accepted Answers:

$\text{Rank}(P) = \text{Rank}(Q)$

$\text{Rank}(PQ) = \text{Rank}(QP)$

[Check Answers](#)