



L10.4: Critical points for multivariable functions



Transcript Language

English



Hello, and welcome to the Maths 2 component of the online B.Sc. program on data science

and programming.

This video is about critical points for multivariable functions.

So, we have studied the notion of critical points for functions of single variable.

Let us recall what that was.

So, for functions of one variable, a point a is called a critical point of a function



f_x , if either f is not differentiable at that point or f prime at a is 0.

So, if f is differentiable at a point of local extremum, it satisfies that f prime a is 0

and so every point of local extremum is a critical point.

This is exactly what we used in order to find the points of local extremum.

So, we set f prime of the function to 0, calculated what are all the points, what are the critical

points, and in amongst them, we did something further in order to determine what kinds of

points they were.

So, here is an example.

We had seen the same picture when we did that as well.

So, we can see here that in this picture, we have several points of local extremum.

For example, there is a point over here, which is a point of local extremum, close to minus

1.5.

There is a point close to 2.

So, these are both local minimums, minima.

And then there is a point between 1 and 1.5, which is a local maximum.

And unfortunately, we can also see that there is another point, which is point 0, which

is what we had called saddle point.

So, not every critical point is a point of local extremum and this is why we needed refined

tests or more other ways of handling critical points in order to determine which of these

are actually local maxima or local minima, because some of them can also be saddle points.

So, what is the saddle point?

It is a critical point which is not a point of local extremum.

So, here, as an example, for 0, what happens is, f' is 0 because you can see that

the tangent line is actually the x axis.

But it is clearly not a point of local extremum, because on the left, you have points for which

the values are, f of those values are less than 0 and on the right, they are bigger than

0, and at 0 it is equal to 0.

So, this was a summary of what we did for functions of one variable.

Of course, we had refined tests.

For example, we went beyond this and after critical points we also looked at the second

derivative and based on that we could classify them in some cases.

So, for multivariable functions, we want to do something similar.

So, let us start reviewing what are points of local extrema for multivariable functions?

So, let f of x be a function defined on a domain D in $\mathbb{R}^n$, and suppose you have a point

a belonging to this domain D . The point a is a local maximum or a point of local maximum

of f if for some open ball B containing this point a for all points in this open ball B

which belong to the domain f evaluated at those points is less than or equal to f evaluated

at a .

So, you want to construct a ball so that within this ball for all those points on which f

is defined, f of a is the largest.

So, this need not be a ball centered at a , it could be a ball which is containing a .

So, but of course, if you can create such a ball you can always create one centered

at a .

Similarly, the point a is a local minimum.

If the same thing happens except that now f of a is the smallest value within that set.

So, remember that for one variable we had open intervals.

We said, if on a small open neighborhood, the point has the least or the largest value.

So, here instead of intervals we have the open ball.

So, in $\mathbb{R}^2$ you will have a disc.

In $\mathbb{R}^3$ you will have a, you will actually have a ball, meaning a sphere.

And in $\mathbb{R}^n$, of course, it is the definition, meaning all those points which are at a fixed

distance from or distance less than a particular number around one point.

So, such a thing so, the local maxima are indeed, locally maximum values.

The local minima are indeed locally minimum values.

And a local extremum is either a local maximum or a local minimum.

So, these are the definitions as far as multivariable functions are concerned, exactly mirroring

what happens for one variable functions.

So, for one variable functions, we just saw that the derivative is always 0 at a local

extremum if it exists.

And the idea there was that if you are at local extremum, then the tangent is parallel

to the x-axis.

So, for the local extrema in the multivariable situation, you would expect something similar.

So, the notion of tangent plane or tangent hyperplane is as we saw what replaces tangent

line.

And as we also observed, the tangent hyperplane or the tangent plane is the equation of that

is governed by the gradient function.

And we have also seen, when we did the behavior of gradients, that gradients mirror, the properties

of gradients mirror that of derivatives for one variable functions.

So, it is natural to think that the role of f' will be played by the gradient.

So, let us look at an example first to observe what happens to the tangent plane and what

happens to the gradient.

So, let us look at the function $\sin(x, y)$.

So, for this function, so here is the graph of that function.

We have seen this graph several times before.

And the reason I have chosen this function is because for this function, it is very clear

what are the local extrema.

So, the sine value takes maximum value 1 and minimum value minus 1.

And you can see that when you are at 1, so there is this ridge over here.

So, all those values are the values 1, so those are going to be local maxima.

And in fact, they will be global maxima, meaning they are maximum amongst all values and the

function takes.

So, in particular, their local maxima, and let us see what happens to the tangent plane

at these points.

Let us choose a point at which we know this local maximum occurs.

For example, if you look at the function sine of x, y , we know that when x, y is $\pi/2$,

then it takes value 1.

So, if I take x to be $\pi/2$ and y to be 1, then that is going to be a point where

you will have a local maximum, and the function value there is 1.

So here is that point that is the point $a, \pi/2$ comma 1 comma 1.

And let us ask what happens to the tangent plane at that point.

So, for that, we will see how all the tangent lines behave.

So, if you take a tangent line at that point, let us see how that behaves.

So, here is a tangent line in the direction of the unit vector u_1 comma u_2 .

I hope you can see that it is indeed a tangent line.

And, as I vary, u_1 and u_2 this tangent line is going to vary.

So, here is how it varies.

So, it is indeed tangent to the graph.

I hope that is apparent from the picture.

And if I remove this graph, so if I remove, let us see it like this, you can see it is

tangent to the graph.

It is touching, it is satisfying the idea that it touches the function at that point.

And if I remove the graph now, you can see it varies in the plane.

And what is that plane, that plane is exactly parallel to the x, y plane.

So, that plane is, in fact, in this case, it is a plane z is equal to 1.

So, we can play it like this.

Now, they, from this perspective, they are all lines, and you can see that this line

is varying on that plane.

So, it is on this plane.

So, what this picture tells us is that as in the one dimensional case where we had the

situation that the local extrema, the tangent lines, if they exist were parallel to the

x axis, so for the two variable case, they will be parallel to the x, y plane, and in

general, you expect them to be parallel to the x_1, x_2, x_n plane.

So, let us study this in terms of the gradient, because after all the gradient determines

the tangent plane or the tangent hyperplane.

So, let us consider the gradient vector at points of local extrema.

So, let f of x be a function defined on a domain D in $\mathbb{R}^n$ containing some open ball around

a point a of local extremum.

Restrict F to a line L passing through a and view it as a function of one variable on L .

Then a is a local extremum for the restricted function on L and hence the directional derivative

of f in the direction of the line L at a is 0.

This is exactly what we saw in that example for sine x, y when we saw the directional,

the tangents along each line in various directions.

So, they were all, the directional derivative was 0.

And as a result, they had no inclination with respect to that plane.

So, they were all flat.

So, they were parallel to the x, y plane.

So, in particular, those partial derivatives which exist at a must be 0.

So, if all the directional derivatives which exist must be 0, in particular, the partial

derivatives, assuming they exist, must be 0.

And so the conclusion is, if gradient f exists for a local extremum a , then gradient f of

a is 0.

I will repeat that, if the gradient exists at that point, then the gradient must be 0.

And when we say 0 here, we mean the 0 vector.

So, this 0 here is the 0 vector.

So, this is the vector.

So, this gives us one way of understanding how to obtain local extrema.

So, what we can try and do is set the gradient to 0 and then look at all those points which

we obtain.

And then out of them, some of them may be local extremum.

This is exactly the strategy followed for the one variable case and such points are

called critical points.

So, point a is called a critical point of a function f of several variables, if either

the gradient does not exist or the gradient exists and equals 0 .

Again, 0 here means the 0 vector.

So, let us look at an example.

So, what are the critical points of this function f of x, y is x^2 plus $6xy$ plus $4y^2$

minus $2x$ minus $4y$.

Well, so this is a very nice.

It is a polynomial function.

So, here, there is no issue about gradient not existing.

So, let us compute what are the partial derivatives.

So, $\frac{\partial f}{\partial x}$ is $2x + 6y - 2$, $\frac{\partial f}{\partial y}$ is $6x + 8y - 4$.

So, for the critical points, I set the gradient function to 0 .

So, the gradient function is the, at x, y is the tuple, $2x + 6y - 2$ comma $6x +$

$8y - 4$, set it to 0 , so set gradient of f to 0 and by 0 we mean the 0 vector.

So, that is $2x$ plus $6y$ plus 2 comma $6x$ plus $8y$ minus 4 is 0 comma 0 .

And now we have to solve these.

So, the solutions for these will give us all the critical points.

So, how do we solve this?

Well, this is a very nice system of linear equations.

And we actually know how to solve this.

So, let us use Gaussian elimination to solve these system, this system of equations.

So, if we do that, we get 2 , 6 , minus 2 and then 6 , 8 , 4 .

Let us divide the first row by 2 .

So, we get 1 , 3 , minus 1 .

Let us use the 1 in the 11 place knock out the 6 .

So, that gives us 1 , 3 , minus 1 , 0 and then 8 , minus 6 times 3 , so 8 minus 18 which is

minus 10 and then 4 minus 6 times minus 1 , so 4 plus 6 is 10 .

So, if I divide the second row by 10 or minus 10 , I get 1 , 3 , minus 1 , 0 , 1 , minus and now

finally I can, final step I can do $R1$ minus $3R2$, so that gives me minus 1 , minus 3 times

minus 1, so minus 1 plus 3 which is 2.

So, this means x is 2, y is minus 1 and that is exactly the critical point.

So, there is only one critical point and it is 2 comma minus 1.

So, I hope it is clear how I am computing these.

So, you set your gradient vector to 0 and then you will get a bunch of equations to

solve.

Need not be linear this time we were lucky.

And then you have to solve them and the solutions will give you the critical points.

Now, there could be lots of critical points.

There is no reason for them to be for only one or two critical points to be there.

For example, we saw the function sine of x , y .

And if you compute the critical points there, we know what the gradient is, I will suggest

you set it to 0 and see what happens.

So, you will see that you get a bunch of equations.

So, one solution is going to be 0, 0, but then there will be tons of solutions corresponding

to cosine of x , y is 0, which we know is an infinite set.

So, this is how we get critical points.

Now, sometimes it may happen that the local extrema are not the critical points, meaning

there are more critical points than just the local extrema.

So, every local extremum is a critical point that we observed already.

Unfortunately, not all of them are, not all critical points are local extrema.

Otherwise, we could be happy and not bother to go ahead.

So, as an example, here is f of x , y is x cube.

So, we have seen the x cube actually, that is an example of a saddle point in one, the

one variable situation.

And the same thing happens here also, because if you recall how f of x cube, f of x is equal

f cube was, it was a function like this, and there was a problem at 0.

And now you take f of x , y cube, you are taking the same function but stretching it along

the y axis, because y has no role to play as such.

So, the entire y axis will consist of points for which the gradient is 0, but they are

not local extremum.

I will suggest that you check this.

So, a saddle point is a critical point a such that the gradient of f at a exists and equals

0, but a is not a local extremum.

So, all these points that I just said, along the y -axis, those are all going to be saddle

points for the function f of x, y is x^3 .

So, here is an example of a saddle point, which is what gives it its name.

So, this is the function f of x, y is $x^2 - y^2$.

And if you look at this function, what is happening is on one side, it is like this.

And on the other side, it is like this.

So, because you have plenty of directions, you can have this weird situation where it

is a local minimum along some lines and it is a local maximum along some other lines,

and that is exactly what is happening here.

So, it is, gradient is 0.

So, in fact, all the directional derivatives are 0.

And what is happening is that in each direction, it is either a local maximum or a local minimum.

But globally, it is neither, neither a local maximum, nor a local minimum.

And this function or this picture actually gives us why it is called a saddle point,

because it looks like a saddle of a horse.

Let us look at a couple of other examples of saddle points, one such is going to be

at when x is 0 and y is 0 for the function $\sin x, y$.

So, I will zoom in and then maybe you can see what is happening.

If I go in here, now this is very close to 0, 0.

And let us see if I can raise this up.

So, you can see what is happening.

So, over here in one, on one direction, this is a local minimum, and you can see that very

clearly.

But on the other side, the function is actually decreasing.

So, over here, it is a local maximum.

And so this function is a, this function as a saddle point at the point $0, 0$.

So, if you take the line y is equal to x , then the function is like sine of x square.

So, x square is always positive as we know.

So, on that line, the sine is going to increase from 0 onwards.

So, on both sides it will increase.

On the other hand, if you take the line y is equal to minus x , then you are going to

get sine of minus x square and which is minus sine x square, which will always be negative

when between, when x is between 0 and $\pi/2$.

So, it will, on the other side, it will decrease.

And so this is a saddle point as a result.

So, it is a point at which the gradient is 0 , but it is not a local extremum.

So, now that we have identified the critical points as local minimum, local maximum or

saddle points, we can ask how do we identify them and so for that we will, like the function

of one variable where we add a second derivative test, there will be something similar for

the function of many variables and we will develop theory for that in the next video.

So, the final topic that we will do in this video is the notion of absolute or global

extremum.

So, suppose we have a function defined on a domain D in $\mathbb{R}^n$ and suppose we have a point

in D . So, this point is an absolute maximum or global maximum if the value of f assumed

on that point that is $f(a)$ is bigger than $f(x)$ for all x within that domain.

So, notice here that we are not seeing local here.

So, for local maximum we said on some small ball it has a largest value that is not what

we are seeing here.

We are saying across the entire domain it has the largest value.

Similarly, for absolute minimum or global minimum, it has the smallest value amongst

all points in the domain.

So, here are two examples.

These are the prototypes of absolute maximum and absolute minimum.

So, this is a function of x square plus y square and the point $0, 0$ is a global minimum

or the function attains a global minimum at the point $0, 0$.

And this is the function minus x square minus y square and $0, 0$ is a global maximum.

This is clear because we know that squares are always positive.

So, from there it is clear that these are global minimum and maximum respectively for

these functions.

So, these are prototypes of global maxima.

So, then how do we find, how do we know they exist and how do we find them.

So, for the existence similar to the one variable case, where we restricted ourselves to what

we called a closed and bounded interval, here we have a notion called closed and bounded

domain.

So, a domain D in $\mathbb{R}^n$ is called closed if it contains all its boundary points.

So, what do we mean by boundary points.

So, a boundary point is one which is on the boundary of that domain.

So, this is visually very clear what we mean by the boundary of the domain.

There is a technical definition let us not get into that.

So, domain D in $\mathbb{R}^n$ is called bounded if it is contained inside a ball around 0 with finite

radius.

So, what this means is it is bounded means it can vary around but it should be like this.

So, there is some huge ball.

It could be very, very, very large.

And the domain lies entirely inside that.

It could have radius as large as you want maybe 1,000 or 20 million whatever, but there

is some domain, there is some ball of as large radius as you want inside which this domain

lies.

So, it does not go out.

So, for example, the x, y plane is not bounded, because it shoots off.

There is no ball in which, there is no disc in which you can contain that or if you take

just the let us say a as a cone inside that, say the first quadrant in the x, y plane that

is not bounded.

So, you want something which is inside a large circle or a large disc rather.

So, here is the main fact, if the domain D is closed and bounded and f is continuous

on D , continuity is important, then the global maximum and minimum must exist on this domain.

So, closed and bounded guarantees the existence of the global maximum and minimum.

There could be more than one point.

So, there could be several points in which the same thing happens.

For example, we saw the function sine of x, y , several points where it takes the value

1, several points where it is the value minus 1.

So, note that the global maximum and minimum are in particular local maxima or local minima,

unless they are on boundary points.

And this statement is what gives us the key to finding them.

Thus, to find the global maximum and minimum we find the critical points inside the domain

D. So, remember, critical points are inside a domain.

We have to have, the critical points are defined inside where there is a ball around this point

lying inside D.

So, then that leaves us with boundary points.

So, you restrict your function now to the boundary.

It is in one, boundary means it is one dimension less.

So, you found find the critical points on the boundary.

Then the boundary itself may have a boundary.

So, you find the critical points on that, and then you check, so you continue this process.

So, if you are in n dimensions, you have to drop dimension all the way till you reach

a point.

And then you check the value of f on all the critical points that you obtain.

And amongst them you choose whatever is maximum or whatever is minimum those will be of global

minimum or maximum.

We have done this exactly this process when we did the one variable functions, when we

check for critical points within the open interval, and then we checked on the boundary

points.

Let us do an example to set ideas.

So, find the absolute maximum and minimum of the function, f of x , y is x cube plus

y cube minus $3x$ minus $3y$ square plus 1 over the square with vertices $0, 0, 2, 0, 2, 2,$

$0, 2.$

Let me first draw this square.

So, here $0, 0$, here is $2, 0$, here is $0, 2$, here is $2, 2$, and here is 2 comma 0 .

Let us what happens for this.

So, now, here, there is an open part inside.

So, this is the open part of the square.

So, first, we have to check what happens, what are the critical points inside this for

this function, then we have to restrict this function on each of the edges.

So, then we will restrict this function on this edge, this edge, this edge and this edge,

see what happens.

And then finally, we have to check what happens to the functions, to the function on these

four points.

That is the strategy.

That is what we are going to do.

Let us start with what happens inside the square.

So, for that, we will find gradient.

Now, first of all, this is a polynomial function.

So, all the issues with, whether partial derivatives exist, etcetera, they do not play any role.

So, we do not have to bother about points where gradient does not exist or they are

not differentiable when you view them on the sides, etcetera.

So, that is not an issue.

So, let us write down what is the gradient.

So, this is $3x^2 - 3$, $3y^2 - 3$, minus $6y$.

Let us evaluate, let us set this to 0 and see what are the critical points.

So, $3x^2 - 3 = 0$, and $3y^2 - 6y = 0$, let us try to solve this.

So, this is saying $x^2 - 1 = 0$.

So, $x^2 = 1$.

The other equation is $y^2 - 2y = 0$.

So, $y - 2 = 0$.

So, we know the solutions for these.

So, the possible solutions are $x = \pm 1$ and $y = 0$ or 2 .

These other possible solutions.

Now, let us see which of these lie inside our square.

So, when $x = -1$ that does not lie inside our square, so we can discard that.

So, the critical points coming from, coming inside here are the points where the x coordinate

is 1, the y coordinate is 0 or the x coordinate is 1 and the y coordinate is 2.

So, this point 1 comma 2 actually lies on the boundary, but we can consider it anyway

just to check, although in principle this is not a critical point when we restrict to

the open part, but we can keep it anyway.

So, these are the critical points for the square.

So, this is one set of critical points.

Now, let us restrict it to the boundary.

So, first, let us do it on the bottom boundary.

So, that is where y is 0.

So, now we want to look at the function f of x comma 0, where x is between 0 and 2,

actually, strictly between, but again, we can be considerate and take the end points

as well because anyway we are going to take them.

So, f of x comma 0 is x^3 minus $3x$ plus 1.

So, the derivative let us set that to 0.

That means $3x^2$ minus 3 is 0.

Well, we have solved this equation above.

So, that means we get x is plus minus 1.

So, which means we should take the points, 1 comma 0, minus 1 comma 0 does not lie in

the, in this interval.

So, 1 comma 0 is already there.

So, these two points, actually both these points are on the boundary.

So, we will take them anyway.

So, this is already taken care of.

Similarly, we can take the top edge.

So, for that the function is f of x comma 2.

So, the function is x cube minus, x cube plus 8 minus $3x$ minus 3 times 4, which is minus

12 plus 1, that is x cube minus $3x$ and plus some constant, does not really matter, so

minus 3.

So, I set it to 0.

And again, I get the same thing.

So, but these are not on the boundary.

So, I do not have to bother about this.

So, I do not get anything here.

Let us do the two ages, two other ages.

So, we have f of 0 comma y between 0 and 2 .

So, if we do that, we get y^3 minus $3y^2$ plus 1 .

But actually, we have solved this as well.

So, this gives us $3y^2$ is minus $6y$ is 0 , so y times y minus 2 is 0 which means y

is 0 or y is 2 .

So, the points that we get from here are 0 comma 0 and 0 comma 2 , which anyway we are

going to take, so let us add them in 0 comma 0 , 0 comma 2 .

And for the other part, where we have f of 2 comma y , we will get the same function but

with a different constant, so plus c , so we will get the same points.

So, we have added them in already.

So, that means we have taken all the points which are critical points on the inside on

each edge and now we will add the boundary points, the corner points as well.

So, that is 2 comma 2, 2 comma 0, and the others are already taken.

So, these are the six points at which we need to check for maximum or minimum.

And if we do that, we get, so if I evaluate that, let us see what we get.

So, I had 1 comma 0, I get 1 minus 3 plus 1, so that is 2 minus 3, so I get minus 1.

For 1 comma 2 I get 1 plus 8 minus 3, so 9 minus 3 is 6 minus 3 times 2 square, so minus

12 so 6 minus 12 so minus 6 plus 1 so minus 5.

For 0 comma 0 I get 1.

For 0 comma 2 I get 8 minus 12 plus 1, so minus 4 plus 1 so minus 3.

And for 2 comma 2, so I get 8 plus 8 16 minus 6 is 10 minus 12 so minus 2 plus 1, so minus

1.

And for 2 comma 0 I get 8 minus 6 so 2 plus 1 so 3.

So, the absolute maximum is 2 comma 0, so absolute maximum occurs at 2 comma 0 and absolute

minimum occurs at 1 comma 2 and the values are 3 and minus 5, respectively.

So, I hope this example explains what you have to do in order to get the absolute maximum

and minimum over a closed and bounded domain.

Thank you.