



## Solve with us 3.1 - Not Graded



This assignment will not be graded and is only for practice.

Key Points: 1Key Points: 1

Definition of a vector space  $V$ : Definition of a vector space  $V$ :

- A vector space  $V$  over  $R$  is a set along with two functions:

1.  $+: V \times V \rightarrow V$  :  $V \times V \rightarrow V$  ( $V$  is closed under vector addition i.e., if  $x, y \in V \Rightarrow x + y \in V, \forall x, y \in V \Rightarrow x + y \in V, \forall x, y \in V$ )

2.  $.: R \times V \rightarrow V$  :  $R \times V \rightarrow V$  ( $V$  is closed under scalar multiplication i.e., if  $x \in V$  and  $c \in R \Rightarrow c.x \in V, \forall x \in V$ , and  $\forall c \in R$   
 $x \in V$  and  $c \in R \Rightarrow c.x \in V, \forall x \in V$ , and  $\forall c \in R$ )

- Axioms of vector additions of a vector space

1.  $v_1 + v_2 = v_2 + v_1, \forall v_1, v_2 \in V$   $v_1 + v_2 = v_2 + v_1, \forall v_1, v_2 \in V$ .

2.  $v_1 + (v_2 + v_3) = (v_1 + v_2) + v_3, \forall v_1, v_2$  and  $v_3 \in V$   $v_1 + (v_2 + v_3) = (v_1 + v_2) + v_3, \forall v_1, v_2$  and  $v_3 \in V$ .

3. There exists an element  $0$  (called the zero vector of  $V$ ) in  $V$  such that  $0 + v = v, \forall v \in V$   
 $0 + v = v, \forall v \in V$ .

4. There exists a vector  $v'$  in  $V$  such that  $v' + v = v + v' = 0, \forall v \in V$   
 $+ v = v + v' = 0, \forall v \in V$  ( $v'$  is called the inverse ( or negative ) of  $v$  with respect to addition).

- Axioms relating vector addition and scalar multiplication of a vector space (since this is usual multiplication in real numbers, we suppress the "." sign ).

1. For each vector  $v \in V$   $1v = v.1v = v$ .

2. For each vector  $v \in V$  and for each pair  $a, b \in R, (a + b)v = av + bv. a, b \in R,$   
 $(a + b)v = av + bv$ .

3. For each  $a \in R$  and for each pair  $v_1, v_2 \in V, a(v_1 + v_2) = av_1 + av_2. v_1, v_2 \in V, a(v_1 + v_2) = av_1 + av_2$ .

4. For each vector  $v \in V$  and for each pair  $a, b \in R, (ab)v = a(bv). a, b \in R, (ab)v = a(bv)$ .

Consider a set  $V = \{(1, -y) \mid y \in R\} \subseteq R^2$   $V = \{(1, -y) \mid y \in R\} \subseteq R^2$  with the usual addition and scalar multiplication as in  $R^2$ . Answer the following questions (1) and (2).

1 point

Which of the following option(s) is (are) true?

[Hint:Hint: Check the axioms of vector space.]

☐

$(1, 2) \in V$   $(1, 2) \in V$ .

☐

$(2, 3) \in V$   $(2, 3) \in V$ .

☐

$V = R^2$   $V = R^2$ .

☐

$v_1 = (1, -5)$  and  $v_2 = (1, 7) \in V \Rightarrow v_1 + v_2 \in V$   $v_1 = (1, -5)$  and  $v_2 = (1, 7) \in V \Rightarrow v_1 + v_2 \in V$ .

☐

$v = (1, 9) \in V, c \in R \Rightarrow cv \in V, \forall c \in R$   $v = (1, 9) \in V, c \in R \Rightarrow cv \in V, \forall c \in R$ .





For all  $v \in V, cv \in V$  For all  $v \in V, cv \in V$  only for  $c = 1c = 1$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$(1, 2) \in V(1, 2) \in V$ .

For all  $v \in V, cv \in V$  For all  $v \in V, cv \in V$  only for  $c = 1c = 1$ .

1 point

Which of the following option(s) is (are) true?



Zero element does not exist in  $VV$ .



Zero element of  $VV$  is  $(0,0)$ .

☐ Inverse element of  $(1, 5)$  is  $(-1, -5)$ .

☐ Inverse element of  $(1, 5)$  is  $(1, -5)$ .

☐ Inverse element of  $(1, 5)$  does not exist.



Inverse element of any element of  $VV$  does not exist.

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

Zero element does not exist in  $VV$ .

Inverse element of  $(1, 5)$  does not exist.

Inverse element of any element of  $VV$  does not exist.

Consider a set  $V = \{(x, y, z) \mid x + y + z = 0, x, y, z \in R\} \subseteq R^3$   $V = \{(x, y, z) \mid x + y + z = 0, x, y, z \in R\} \subseteq R^3$  with the

usual addition:  $(x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2)$ ,  $(x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2)$ ,

where  $(x_1, y_1, z_1), (x_2, y_2, z_2) \in V$  where  $(x_1, y_1, z_1), (x_2, y_2, z_2) \in V$  and

scalar multiplication:  $c(x, y, z) = (cx, cy, cz)$ ,  $c \in R$  and  $(x, y, z) \in V$ .  $c(x, y, z) =$

$(cx, cy, cz)$ ,  $c \in R$  and  $(x, y, z) \in V$ .

Answer the following questions (3) and (4).

1 point

Which of the following option(s) is (are) true?

[Hint:Hint: For a vector  $v = (x, y, z)v = (x, y, z)$ , check whether the condition  $x + y + z = 0$   $x + y + z = 0$  is satisfied or not.]



$(1, 2, -3) \in V \in V$ .



$(-1, 2, 1) \in V \in V$ .



$V = R^3 V = R^3$ .



$(5, -7, 2) \in V(5, -7, 2) \in V$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$(1, 2, -3) \in V \in V$ .

$(5, -7, 2) \in V(5, -7, 2) \in V$ .

1 point

Inverse element of  $(-2, -4, 6)$  with respect to vector addition is

- ☐  $(2, 4, -6)$ .  
☐  $(2, -4, -6)$ .  
☐  $(2, 4, 6)$ .  
☐  $(-2, 4, -6)$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$(2, 4, -6)$ .

1 point

Consider a set  $V = \{(x, y, z) \mid x + y + z = 0, x, y, z \in R\} \subseteq R^3$   $V = \{(x, y, z) \mid x + y + z = 0, x, y, z \in R\} \subseteq R^3$  with the usual addition as in  $R^3$  and scalar multiplication is defined as

$$c(x, y, z) = \begin{cases} (0, 0, 0) & c = 0 \\ (cx, cy, cz) & c \neq 0 \end{cases} \quad (x, y, z) \in V, c \in R$$

$c = 0 \implies c(x, y, z) = (0, 0, 0)$   $(x, y, z) \in V, c \in R$

Which of the following option(s) is (are) true?

[Hint: By adding any two elements in  $V$ , or by multiplying a scalar with any element in  $V$ , check whether the resultant element is in  $V$  or not.]

☐

$V$  is closed under addition i.e.,  $(v_1, v_2 \in V \implies v_1 + v_2 \in V)$

☐

$V$  is closed under scalar multiplication i.e.,  $(c \in R, v \in V \implies cv \in V)$

$(c \in R, v \in V \implies cv \in V)$ .

☐

$V$  has zero element with respect to addition.

☐

$1.v = v$   $1.v = v$ , where  $1 \in R$  and  $v \in V$

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$V$  is closed under addition i.e.,  $(v_1, v_2 \in V \implies v_1 + v_2 \in V)$

$V$  has zero element with respect to addition.

1 point

Consider usual addition and scalar multiplication on  $R^2$  and the subset

$V = \{(1, -y) \mid y \in R\} \subseteq R^2$

Which of the following option(s) is (are) true?

[Hint: Take two arbitrary elements  $(1, -y_1)$  and  $(1, -y_2)$  from  $V$  and check the conditions given in the options.]

☐

For each element  $v \in V$  and for each pair  $a, b \in R, (a + b)v = av + bv$

$(a + b)v = av + bv$ .

☐

For each  $a \in R$  and for each pair  $v_1, v_2 \in V, a(v_1 + v_2) = av_1 + av_2$

$$) = av_1 + av_2.$$



For each element  $v \in V$  and for each pair  $a, b \in R$ ,  $(ab)v = a(bv)$ ,  $a, b \in R$ ,  $(ab)v = a(bv)$ .



The set  $V$  with the usual addition and scalar multiplication in  $R^2$  is a vector space.

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

For each element  $v \in V$  and for each pair  $a, b \in R$ ,  $(a + b)v = av + bv$ ,  $a, b \in R$ ,

$$(a + b)v = av + bv.$$

For each  $a \in R$  and for each pair  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ ,  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ .

For each element  $v \in V$  and for each pair  $a, b \in R$ ,  $(ab)v = a(bv)$ ,  $a, b \in R$ ,  $(ab)v = a(bv)$ .

1 point

Consider a set  $V = \{(x, y, z) \mid x + y + z = 0, x, y, z \in R\} \subseteq R^3$

with the usual addition as in  $R^3$  and scalar multiplication is defined as

$$c(x, y, z) = \begin{cases} (0, 0, 0) & c = 0 \\ (cx, cy, cz) & c \neq 0 \end{cases} \quad (x, y, z) \in V, c \in R$$

$$c = 0 \implies (0, 0, 0) \in V, c \in R$$

Which of the following option(s) is (are) true?



For each element  $v \in V$  and for each pair  $a, b \in R$ ,  $(a + b)v = av + bv$ ,  $a, b \in R$ ,

$$(a + b)v = av + bv$$



For each  $a \in R$  and for each pair  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ ,  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ .



For each element  $v \in V$  and for each pair  $a, b \in R$ ,  $(ab)v = a(bv)$ ,  $a, b \in R$ ,  $(ab)v = a(bv)$ .



$V$  is a vector space.

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

For each  $a \in R$  and for each pair  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ ,  $v_1, v_2 \in V$ ,  $a(v_1 + v_2) = av_1 + av_2$ .

Key Points: 2

- Definition of a subspace: A subset  $W$  of a vector space  $V$  is a subspace of  $V$  if it is a vector space with respect to the same operations defined on  $V$ .
- Criteria to check a subset  $W$  of a vector space  $V$  is a subspace:
  - Zero vector should be in  $W$  i.e.,  $0 \in W$ .
  - If two vectors  $v_1$  and  $v_2 \in W$ , then  $v_1 + v_2 \in W$ .
  - Let  $c \in R$  and if  $v \in W$ , then  $cv \in W$ .

1 point

Let  $W = \{(x, y) \mid x + 3y = 1, x, y \in \mathbb{R}\} \subseteq \mathbb{R}^2$ . Which of the following options are true?

☐

Zero element exists in  $W$ .

☐

If  $v_1$  and  $v_2 \in W$ , then  $v_1 + v_2 \notin W$ .

☐

Let  $c \in \mathbb{R}$  and if  $v \in W$ , then  $cv \in W$ .

☐

$W$  is not a subspace of  $\mathbb{R}^2$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

If  $v_1$  and  $v_2 \in W$ , then  $v_1 + v_2 \notin W$ .

$W$  is not a subspace of  $\mathbb{R}^2$ .

1 point

Let  $W = \{A \mid A \in M_{2 \times 2}(\mathbb{R}) \text{ and } A \text{ is a diagonal matrix}\}$ .

Which of the following option(s) is (are) true?

☐

Zero vector does not exist in  $W$ .

☐

If  $v_1$  and  $v_2 \in W$ , then  $v_1 + v_2 \in W$ .

☐

Let  $c \in \mathbb{R}$  and if  $v \in W$ , then  $cv \in W$ .

☐

$W$  is a subspace of  $M_{2 \times 2}(\mathbb{R})$ .

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

If  $v_1$  and  $v_2 \in W$ , then  $v_1 + v_2 \in W$ .

Let  $c \in \mathbb{R}$  and if  $v \in W$ , then  $cv \in W$ .

$W$  is a subspace of  $M_{2 \times 2}(\mathbb{R})$ .

Key Points: 3

- Linear combination: Let  $V$  be a vector space and  $v_1, v_2, \dots, v_k \in V$ . The linear combination of  $v_1, v_2, \dots, v_k$  with coefficients  $a_1, a_2, \dots, a_k \in \mathbb{R}$  is the vector  $v = \sum_{i=1}^k a_i v_i$ .
- A vector  $v \in V$  is said to be a linear combination of  $v_1, v_2, \dots, v_k \in V$  if there exist  $a_1, a_2, \dots, a_k \in \mathbb{R}$  so that  $v = \sum_{i=1}^k a_i v_i$ .

1 point

Which of the following vectors are linear combinations of  $(2, 1)$  and  $(6, 3)$  with usual addition and scalar multiplication in  $\mathbb{R}^2$ ?

☐  $(2, 1) + 5(6, 3)$ 
☐  $(2, 1)$ 
☐  $(6, 3)$ 
☐  $(3, 6)$

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

$(2, 1) + 5(6, 3)$

$(2, 1)$

$(6, 3)$

1 point

Let  $V$  be a vector space and  $v_1, v_2, v_3$  and  $v_4 \in V$ . If  $v_1$  is a linear combination of  $v_i, i = 2, 3, 4$  i.e.,  $v_1 = av_2 + bv_3 + cv_4$ , then which of the following options are true?

☐

If  $a \neq 0$ , then  $v_2$  is a linear combination of  $v_i, i = 1, 3, 4$ .

☐

If  $a = 0$  and  $b \neq 0$ , then  $v_3$  is a linear combination of  $v_i, i = 1, 4$ .

☐

$v_2$  is a linear combination of  $v_2$ .

☐ None of the above.

**No, the answer is incorrect.**

**Score: 0**

**Accepted Answers:**

If  $a \neq 0$ , then  $v_2$  is a linear combination of  $v_i, i = 1, 3, 4$ .

If  $a = 0$  and  $b \neq 0$ , then  $v_3$  is a linear combination of  $v_i, i = 1, 4$ .

$v_2$  is a linear combination of  $v_2$ .