Week 4 - Graded Assignment 4

The due date for submitting this assignment has passed.
Due on 2026-03-11, 23:59 IST.
You may submit any number of times before the due date. The final submission will be considered for grading.

You have last submitted on: 2026-03-11, 08:07 IST

Note: This assignment will be evaluated after the deadline passes. You will get your score 48 hrs after the deadline. Until then the score will be shown as Zero.

Multiple Select Questions (MSQ)

1 point
Which of the following statements are correct?

- [ ] If AA and BB are two square matrices, then Rank(AB)(AB)=Rank(BA)(BA).
- [x] If AA and BB are two $2 \times 2$2 × 2 square matrices, then Rank(AB) ≤ (AB) ≤ Rank(B)(B).
- [x] If AA and BB are two $2 \times 2$2 × 2 square matrices, then Rank(AB) ≤ (AB) ≤ Rank(A)(A).
- [x] If AA and BB are $n \times n$n × n matrices of rank nn, then ABAB has rank nn.

Yes, the answer is correct.
Score: 1
Accepted Answers:
If AA and BB are two $2 \times 2$2 × 2 square matrices, then Rank(AB) ≤ (AB) ≤ Rank(B)(B).
If AA and BB are two $2 \times 2$2 × 2 square matrices, then Rank(AB) ≤ (AB) ≤ Rank(A)(A).
If AA and BB are $n \times n$n × n matrices of rank nn, then ABAB has rank nn.

1 point
Match the vector spaces (with the usual scalar multiplication and vector addition as in $M_{3 \times 3}(R)$M3×3 (R)) in column A with their bases in column B and the dimensions of the vector spaces in column C in Table : M2W4GA1.

Choose the correct option.

- [ ] a → → i → → 2.
- [ ] b → → i → → 2.
- [x] a → → ii → → 3.
- [ ] c → → iii → → 1
- [ ] b → → iii → → 2.
- [ ] c → → i → → 1.

Partially Correct.
Score: 0.33

Accepted Answers:
a → → ii → → 3.
b → → iii → → 2.
c → → i → → 1.
1 point

Consider the vector space $V = \{\begin{bmatrix} a & b \\ b & c \end{bmatrix} | a, b, c \in R\}$V = {[abbc] | a, b, c ∈ R}. The subset $S = \{\begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\}$S = {[2113], [1001]} is a linearly independent subset of VV . Which of the following matrices AA are vectors in VV such that $S \cup \{A\}$S ∪ {A} is a basis for VV ?

- [ ] $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$[1101]
- [x] $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$[0110]
- [x] $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$[2003]
- [ ] $\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$[0111]
- [x] $\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$[1−1−11]
- [ ] $\begin{bmatrix} 1 & -1 \\ -1 & 0 \end{bmatrix}$[1−1−10]

Yes, the answer is correct.
Score: 1
Accepted Answers:
$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$[0110]

$\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$[2003]

$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$[1−1−11]

1 point
Let $S = \{v_1, v_2, v_3\}$S = {v1, v2, v3} be a set of three vectors in R^4R4. In order to determine the linear dependence/independence of SS, we construct a matrix AA whose i^{th} ith column is given by v_i vi, for $i \in \{1, 2, 3\}$i ∈ {1, 2, 3}. Let RR denote the reduced row echelon form (RREF) of AA, and let r_{ij} rij denote the $(i, j)^{th}$ (i, j)th entry of RR. Choose the correct statements from the following.

- [ ] RR does not have any row comprising entirely of zeros if SS is linearly independent.
- [x] If kk is the number of rows in RR comprising entirely of zeros, then the dimension of Span(S)(S) is given by $4 - k$4 − k.
- [ ] If r_{11} r11 and r_{23} r23 are the pivots, then $\{v_1, v_2\}$ {v1, v2} is a basis for Span(S)(S).
- [x] If r_{11} r11 and r_{23} r23 are the pivots, then $\{v_1, v_3\}$ {v1, v3} is a basis for Span(S)(S).
- [x] rank(A)(A) = rank(R)(R), i.e., row operations do not alter the rank of a matrix.

Yes, the answer is correct.
Score: 1
Accepted Answers:
If kk is the number of rows in RR comprising entirely of zeros, then the dimension of Span(S)(S) is given by 4 − k4 − k.
If r_{11} r11 and r_{23} r23 are the pivots, then {v_1, v_3}{v1, v3} is a basis for Span(S)(S).
rank(A)(A) = rank(R)(R), i.e., row operations do not alter the rank of a matrix.

Numerical Answer Type (NAT):

Find the rank of the matrix AA, where A = [a_{ij}]A = [aij] is of order 2021 × 20212021 × 2021 and $a_{i,j}$ = min{i, j}ai,j = min{i, j}, i, j = 1, 2, … 2021i, j = 1, 2, … 2021
[Hint:Hint: First do it for smaller matrices, such as 3 × 33 × 3 or 4 × 44 × 4.]

2021

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 2021
1 point

Consider a subspace W = {(x_1, x_2, x_3, x_4, x_5) | 8x_1 + 10x_5 = 0 and 9x_2 + 6x_4 = 0}W = {(x1, x2, x3, x4, x5) | 8x1 + 10x5 = 0 and 9x2 + 6x4 = 0} of R^5 R5. The dimension of WW is

3

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 3
1 point

Consider a matrix A = [2−41 1 1−31−4 3−72 3]A = $\begin{bmatrix} & & & \\ 2 & 1 & -4 & -3 \\ -7 & 1 & 1 & -4 \\ 3 & & & \end{bmatrix}$. The rank of the matrix is

3

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 3
1 point

Consider two matrices A = [3 9 8 −7 −10 −9 2 6 −7]A = [3 −7 2 9 −10 6 8 −9 −7] and B = [3 −7 2 9 −10 6 8 −9 −7]B = [3 −7 2 9 −10 6 8 −9 −7]. Then Rank(A)(A) - Rank (B)(B) is

0

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 0
1 point

Consider the subspace V = {(x, y, z) | z = 7x + 5y, where $x, y, z \in R$}V = {(x, y, z) | z = 7x + 5y, where x, y, z ∈ R} of R^3 R3 and span({$(p, 0, 1), (0, q, 1)$}{(p, 0, 1), (0, q, 1)}) = VV . Find the value of $\frac{1}{pq}$pq1.

35

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 35
1 point

Suppose VV is a vector space defined as V = {A | $A \in M_{4\times4}(R)$, AV = {A | A ∈ M4×4(R), A is an upper triangular matrix, and the sum of the diagonal entries is zero }}. What is the cardinality of a basis of VV ?

9

Yes, the answer is correct.
Score: 1
Accepted Answers:
(Type: Numeric) 9.0
1 point

UU is the vector space of all upper triangular matrices of order 3. VV is the vector space of all lower triangular matrices of order 3. The dimensions of the vector spaces $U, V, U \cap V$U, V, U ∩ V are a, b, ca, b, c respectively. Find $a + b + c$a + b + c. The usual rules of addition and scalar multiplication apply for all three spaces.

15