

**This assignment will not be graded and is only for practice.**

1) Let  $L_1$  and  $L_2$  be affine subspaces of  $\mathbb{R}^3$ , where  $L_1 = U$  and  $L_2 = (2, 0, 1) + W$ , for some **1 point** vector subspaces  $U$  and  $W$  of  $\mathbb{R}^3$ . Let a basis for  $U$  be given by  $\{(2, 0, 1), (1, 1, 0), (0, 1, 0)\}$  and a basis for  $W$  be given by  $\{(1, 0, 1), (0, 1, 1)\}$ . Suppose there is a linear transformation  $T : U \rightarrow W$  such that  $(0, 1, 0) \in \ker(T)$ ,  $T(2, 0, 1) = (0, 1, 1)$  and  $T(1, 1, 0) = (1, 0, 1)$ . An affine mapping  $f : L_1 \rightarrow L_2$  is obtained by defining  $f(u) = (2, 0, 1) + T(u)$ , for all  $u \in U$ . Which of the following options are true?

- ☐  $L_1 = \mathbb{R}^3$
- ☐  $L_2 = \{(x, y, z) \mid x + y - z = 1\}$
- ☐  $L_2 = \mathbb{R}^3$
- ☐  $f(x, y, z) = (x - 2z + 2, z, x - z + 1)$
- ☐  $f(x, y, z) = (x - 2z, z, x - z)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$L_1 = \mathbb{R}^3$$

$$L_2 = \{(x, y, z) \mid x + y - z = 1\}$$

$$f(x, y, z) = (x - 2z + 2, z, x - z + 1)$$

**Solution:**

**Recall the definition of Affine map:**

Let  $L_1$  and  $L_2$  be affine subspaces of  $V_1$  and  $V_2$ , respectively, and let  $U$  and  $W$  be the unique subspaces that correspond to  $L_1$  and  $L_2$ , respectively.

- A map  $f : L_1 \rightarrow L_2$  is an affine map if there exists a linear transformation  $T : U \rightarrow W$  such that

$$f(v_1 + u) = v_2 + T(u),$$

where  $v_1$  and  $v_2$  are elements of  $V_1$  and  $V_2$ , respectively such that  $L_1 = v_1 + U$  and  $L_2 = v_2 + W$ .

**Before we proceed to the solution you may want to recall the following weekly contents:**

- Watch : Week 7: Lecture 2.
- Solve : Activity 7.2: Questions 4, 5 and 7.

**Discussion on Option 1:** It is given that  $L_1 = U$  and  $\{(2, 0, 1), (1, 1, 0), (0, 1, 0)\}$  is a basis of the vector subspace  $U$ .

2) The dimension of  $U$  is

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 3

**1 point**

$L_1 = U$  is a subspace of  $\mathbb{R}^3$ . To know whether  $L_1 = U$  is a proper subspace of  $\mathbb{R}^3$  or not, we just have to compare their dimensions.

3) Now, can you see why  $L_1 = U = \mathbb{R}^3$ ?

**1 point**

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Feedback:

**Feedback:** Dimension of a proper subspace of a vector space is always less than the dimension of the vector space. Here,  $\dim(L_1) = \dim(\mathbb{R}^3)$ .

Accepted Answers:

Yes

**Discussion on Options 2 and 3:** It is given that  $L_2 = (2, 0, 1) + W$  and  $\{(1, 0, 1), (0, 1, 1)\}$  is a basis for the vector subspace  $W$ . Now to find the explicit description of  $L_2$ , we have to find the explicit description of  $W$  first.

$$\begin{aligned} W &= \{a(1, 0, 1) + b(0, 1, 1) \mid a, b \in \mathbb{R}\} \\ &= \{(a, b, a+b) \mid a, b \in \mathbb{R}\} \end{aligned}$$

As  $W$  is a subspace of  $\mathbb{R}^3$ , we want to express it using the usual  $x, y, z$  co-ordinate and the relation between them. From the above description we have,  $x = a, y = b, z = a + b$ , which implies that  $z = x + y$ . So

$$W = \{(x, y, z) \mid x + y - z = 0, \text{ where } x, y, z \in \mathbb{R}\}$$

4) Now, are you able to find the equation of  $L_2$ ?

**1 point**

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Feedback:

$$L_2 = (2, 0, 1) + W = (2, 0, 1) + \{(x, y, z) \mid x + y - z = 0\} = \{(2+x, y, z+1) \mid x + y - z = 0\}$$

Accepted Answers:

Yes

5) Do you agree that the sets  $\{(2+x, y, z+1) \mid x+y-z=0\}$  and  $\{(a, b, c) \mid a+b-c=1\}$  **1 point** are the same?

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Feedback:

**Feedback:**

$$a = 2 + x, \quad b = y, \quad c = z + 1 \quad \text{such that } x + y - z = 0.$$

$$a + b - c = 2 + x + y - z - 1 = x + y - z + 1 = 0 + 1.$$

Accepted Answers:

Yes

### Discussion on Option 4 and 5:

**Step 1:** Here, we have to find an algebraic expression for the affine map  $f : L_1 \rightarrow L_2$ . To get an algebraic expression for  $f$ , we have to find the algebraic expression which represents the linear transformation  $T : U \rightarrow W$ .

Given:

$$T(0, 1, 0) = (0, 0, 0), \quad T(2, 0, 1) = (0, 1, 1). \quad T(1, 1, 0) = (1, 0, 1)$$

6) Is  $T$  uniquely determined by the above information?

**1 point**

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Feedback:

**Feedback:** As  $\{(0, 1, 0), (2, 0, 1), (1, 1, 0)\}$  forms a basis of  $U$  and  $T$  is defined for each element of this basis, then the above description completely and uniquely describes  $T$ .

Accepted Answers:

Yes

**Step 2:** Since  $\{(0, 1, 0), (2, 0, 1), (1, 1, 0)\}$  is a basis of  $U = \mathbb{R}^3$ , we can write any element  $(x, y, z) \in U$  as a linear combination of  $\{(0, 1, 0), (2, 0, 1), (1, 1, 0)\}$ .

$$(1) \quad (x, y, z) = a(0, 1, 0) + b(2, 0, 1) + c(1, 1, 0)$$

Apply  $T$  on both sides:

$$T(x, y, z) = aT(0, 1, 0) + bT(2, 0, 1) + cT(1, 1, 0)$$

$$T(x, y, z) = a(0, 0, 0) + b(0, 1, 1) + c(1, 0, 1)$$

To get an algebraic expression for  $T(x, y, z)$ , we just have to know the values of  $a$ ,  $b$  and  $c$  in terms of  $x, y$  and  $z$ .

7) Now, can you find the values of  $a$ ,  $b$  and  $c$  using equation (1)?

**1 point**

☐ Yes

☐ No

No, the answer is incorrect.

Score: 0

Feedback:

**Feedback:** Form equation (1), we get

$$2b + c = x$$

$$a + c = y$$

$$b = z$$

Accepted Answers:

Yes

Solve the above system for  $a$ ,  $b$  and  $c$ .

8) What is the value of  $a$  ?

**1 point**

- ☐  $z$
- ☐  $x - 2z$
- ☐  $y + 2z - x$
- ☐  $x - y$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$y + 2z - x$

9) What is the value of  $b$  ?

**1 point**

- ☐  $z$
- ☐  $x - 2z$
- ☐  $y + 2z - x$
- ☐  $x - y$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$z$

10) What is the value of  $c$  ?

**1 point**

- ☐  $z$
- ☐  $x - 2z$
- ☐  $y + 2z - x$
- ☐  $x - y$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$x - 2z$

**Step 3:**

$$T(x, y, z) = (0, z, z) + (x - 2z, 0, x - 2z) = (x - 2z, z, x - z)$$

By definition:

$$f(x, y, z) = (2, 0, 1) + T(x, y, z) \quad \text{where } (x, y, z) \in U.$$

By substituting the value  $T(x, y, z)$  in the above expression, we get

$$f(x, y, z) = (x - 2z + 2, z, x - z + 1) \quad \text{for all } (x, y, z) \in U.$$

**Your score is: 0/10**