AQ5.2: Activity Questions 2 - Not Graded

This assignment will not be graded and is only for practice.
Level 1

1 point
Rank of a matrix AA is equal to
- [ ] the row rank of the matrix AA.
- [ ] the column rank of the matrix AA.
- [ ] the nullity of the matrix AA.
- [ ] the number of dependent variables in the system of equations $Ax = 0$Ax = 0.
- [ ] the number of independent variables in the system of equations $Ax = 0$Ax = 0.

No, the answer is incorrect.
Score: 0
Accepted Answers:
the row rank of the matrix AA.
the column rank of the matrix AA.
the number of dependent variables in the system of equations $Ax = 0$Ax = 0.
Let nullity of the matrix $A_{3\times 5}$A3×5 be 2. Find the rank of the matrix $A_{3\times 5}$A3×5.

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 3
1 point

Find the rank of the matrix $\begin{bmatrix}1&2&3\\4&5&6\\2&4&6\end{bmatrix}$.

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 2
1 point
1 point
If AA is an $m \times n$m × n matrix, then which of the following statements is true?
- ○ $rank(A) + nullity(A) = m$rank(A) + nullity(A) = m
- ○ $rank(A) + nullity(A) = n$rank(A) + nullity(A) = n
- ○ $rank(A) + nullity(A) = mn$rank(A) + nullity(A) = mn
- ○

$rank(A) + nullity(A) = max\{m, n\}$rank(A) + nullity(A) = max{m, n}

$rank(A) + nullity(A) = min\{m, n\}$rank(A) + nullity(A) = min{m, n}

No, the answer is incorrect.

Score: 0

Accepted Answers:

$rank(A) + nullity(A) = n$rank(A) + nullity(A) = n

1 point

Which of the following is true for a homogeneous system of linear equations $Ax = 0$Ax = 0?

- [ ] It may have no solution.
- [ ] It can have infinitely many solutions.
- [ ] It can have a unique solution.
- [ ] Whenever it has a non-trivial(non zero) solution, it must have infinitely many solutions

No, the answer is incorrect.

Score: 0

Accepted Answers:

It can have infinitely many solutions.
It can have a unique solution.
Whenever it has a non-trivial(non zero) solution, it must have infinitely many solutions

1 point

Which of the following options are correct for a square matrix AA of order $n \times n$n × n, where nn is any natural number?

- [ ] If the determinant is non-zero, then the nullity of AA must be 0.
- [ ] If the determinant is non-zero, then the nullity of AA may be non-zero.
- [ ] If the nullity of AA is non-zero, then the determinant of AA must be 0.
- [ ] If the nullity of AA is non-zero, then the determinant of AA may be non-zero.

No, the answer is incorrect.

Score: 0

Accepted Answers:

If the determinant is non-zero, then the nullity of AA must be 0.
If the nullity of AA is non-zero, then the determinant of AA must be 0.

1 point

Choose the set of correct statements.

- [ ] If nullity of a $3 \times 3$3 × 3 matrix is cc for some natural number cc, $0 \leq c \leq 3$0 ≤ c ≤ 3, then the nullity of $-A$−A will also be cc.
- [ ] $nullity(A + B) = nullity(A) + nullity(B)$nullity(A + B) = nullity(A) + nullity(B)
- [ ] Nullity of the zero matrix of order $n \times n$n × n, is nn.
- [ ] Nullity of the zero matrix of order $n \times n$n × n, is 00.
- [ ] There exist square matrices AA and BB of order $n \times n$n × n, such that nullity of both AA and BB is 00, but the nullity of $A + B$A + B is nn.

No, the answer is incorrect.

Score: 0

Accepted Answers:

If nullity of a $3 \times 33 \times 3$ matrix is cc for some natural number cc, $0 \leq c \leq 30 \leq$ c ≤ 3, then the nullity of $-A-$A will also be cc.
Nullity of the zero matrix of order $n \times n$n × n, is nn.
There exist square matrices AA and BB of order $n \times n$n × n, such that nullity of both AA and BB is 00, but the nullity of $A + B$A + B is nn.
The rank of an $n \times n$n × n matrix all of whose entries are equal to 1 is

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 1
1 point
1 point
If AA is a $3 \times 43 \times 4$ matrix, then which of the following options are true?
☐ rank(A) must be less than or equal to 3.
☐ nullity(A) must be greater than or equal to 1.
☐
If AA has 22 columns which are non-zero and not multiples of each other, while the remaining columns are linear combinations of these 22 columns, then $nullity(A)$ = 2nullity(A) = 2.
☐
If AA has 22 columns which are non-zero and not multiples of each other, while the remaining columns are linear combinations of these 22 columns, then $nullity(A)$ = 1nullity(A) = 1.

No, the answer is incorrect.
Score: 0
Accepted Answers:
rank(A) must be less than or equal to 3.
nullity(A) must be greater than or equal to 1.
If AA has 22 columns which are non-zero and not multiples of each other, while the remaining columns are linear combinations of these 22 columns, then $nullity(A)$ = 2nullity(A) = 2.

Level 2

1 point
Let Ax = 0Ax = 0 be a homogeneous system of linear equations which has infinitely many solutions, where AA is an $m \times n$m × n matrix (where, $m > 1, n >$ 1m > 1, n > 1). Which of the following statements are possible?
☐
$rank(A)$ = $m$rank(A) = m and $m < n$m < n.
☐
$rank(A)$ = $m$rank(A) = m and $m > n$m > n.
☐
$rank(A)$ = $m$rank(A) = m and $m = n$m = n.
☐
$nullity(A)$ = $n$nullity(A) = n.
☐
$nullity(A) \neq 0$nullity(A) = 0.

No, the answer is incorrect.
Score: 0
Accepted Answers:
$rank(A)$ = $m$rank(A) = m and $m < n$m < n.

$nullity(A) = n$nullity(A) = n.
$nullity(A) \neq 0$nullity(A) = 0.

Consider the coefficient matrix AA of the following system of linear equations to answer questions 11 and 12:

$3x_1 + 2x_2 + x_3 = 0$ 3x1 + 2x2 + x3 = 0

$x_1 + x_3 = 0$ x1 + x3 = 0

1 point

Which one of the following vector spaces represents the null space of AA appropriately?

○ $\{(-t, t, t) \mid t \in R\}$ {(−t, t, t) | t ∈ R}

○ $\{(t_1, t_2, \frac{t_2 - t_1}{2}) \mid t_1, t_2 \in R\}$ {(t1, t2, 2t2−t1) | t1, t2 ∈ R}

○ $\{(t, -t, t) \mid t \in R\}$ {(t, −t, t) | t ∈ R}

○ $\{(t_1, t_2, \frac{t_1 + t_2}{2}) \mid t_1, t_2 \in R\}$ {(t1, t2, 2t1+t2) | t1, t2 ∈ R}

No, the answer is incorrect.
Score: 0
Accepted Answers:
$\{(-t, t, t) \mid t \in R\}$ {(−t, t, t) | t ∈ R}

1 point

What will be the rank of AA and nullity of AA?

○ $rank(A) = 3, nullity(A) = 2$rank(A) = 3, nullity(A) = 2

○ $rank(A) = 2, nullity(A) = 1$rank(A) = 2, nullity(A) = 1

○ $rank(A) = 1, nullity(A) = 2$rank(A) = 1, nullity(A) = 2

○ $rank(A) = 2, nullity(A) = 0$rank(A) = 2, nullity(A) = 0

No, the answer is incorrect.
Score: 0
Accepted Answers:
$rank(A) = 2, nullity(A) = 1$rank(A) = 2, nullity(A) = 1

Let $Ax = 0$Ax = 0 be a homogeneous system of linear equations which has a unique solution, where $A \in R^{n \times n}$ A ∈ Rn×n. What is the nullity of AA?

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 0

1 point

Suppose $x_1 \neq 0$x1 = 0 solves $Ax = 0$Ax = 0, where $A \in R^{2 \times 4}$ A ∈ R2×4. What is the minimum number of elements in a linearly independent subset of null space of AA that also spans the set of solutions of $Ax = 0$Ax = 0?

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 2
1 point
1 point
Consider the following sets:
S_1 = {(1, 2, 3), (4, 5, 6), (8, 9, 10)}S1 = {(1, 2, 3), (4, 5, 6), (8, 9, 10)}
S_2 = {(1, 2, 1), (2, 4, 2), (1, 2, 5)}S2 = {(1, 2, 1), (2, 4, 2), (1, 2, 5)}
S_3 = {(1, 2, 3), (4, 5, 6), (7, 8, 10)}S3 = {(1, 2, 3), (4, 5, 6), (7, 8, 10)}
S_4 = {(1, 2), (4, 5)}S4 = {(1, 2), (4, 5)}
S_5 = {(1, 2), (4, 5), (3, 8)}S5 = {(1, 2), (4, 5), (3, 8)}
Choose the set of correct options

- [ ] S_1 S1 is a basis of R^3 R3.
- [ ] S_2 S2 is a basis of R^3 R3.
- [ ] S_3 S3 is a basis of R^3 R3.
- [ ] S_4 S4 is a basis of R^2 R2.
- [ ] S_5 S5 is a basis of R^2 R2.

No, the answer is incorrect.
Score: 0
Accepted Answers:
S_3 S3 is a basis of R^3 R3.
S_4 S4 is a basis of R^2 R2.

Check Answers