



Solve with us 10 - Not Graded



This assignment will not be graded and is only for practice.

Key Points: 1

- Gradient of a function: Gradient of a function: If f is a scalar valued function defined on a domain D in R^n , then the gradient ∇f is defined as the vector valued function $(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n})$ ($\partial x_1 \partial f, \partial x_2 \partial f, \dots, \partial x_n \partial f$).
- If ∇f exists and is continuous in an open ball around the point $(a_1, a_2, \dots, a_n)$, then the directional derivative of the function f at $(a_1, a_2, \dots, a_n)$ in the direction of a vector $\mathbf{v}$ is given by the dot product of the gradient of the function at $(a_1, a_2, \dots, a_n)$ with the unit vector in the direction of $\mathbf{v}$ i.e.,

$$D_{\mathbf{v}} f(a_1, a_2, \dots, a_n) = \nabla f(a_1, a_2, \dots, a_n) \cdot \frac{\mathbf{v}}{\|\mathbf{v}\|} \\ = \frac{\nabla f(a_1, a_2, \dots, a_n) \cdot \mathbf{v}}{\|\mathbf{v}\|}$$

- If ∇f exists and is continuous in an open ball around the point $(a_1, a_2, \dots, a_n)$, then the gradient is the direction in which the directional derivative at P is maximized amongst all directions, i.e., the directional derivative at P is maximized amongst all the directions in the direction of the vector $\frac{\nabla f(a_1, a_2, \dots, a_n)}{\|\nabla f(a_1, a_2, \dots, a_n)\|}$ $|\nabla f(a_1, a_2, \dots, a_n)|$.
- If ∇f exists and is continuous in an open ball around the point $(a_1, a_2, \dots, a_n)$, then the direction opposite to the gradient is the direction in which the directional derivative at P is minimized amongst all directions, i.e., the directional derivative at P is minimized amongst all the direction in the direction of the vector $\frac{-\nabla f(a_1, a_2, \dots, a_n)}{\|\nabla f(a_1, a_2, \dots, a_n)\|}$ $|\nabla f(a_1, a_2, \dots, a_n)|$.
- If ∇f exists and is continuous in an open ball around the point $(a_1, a_2, \dots, a_n)$, then the maximum value amongst all directional derivatives at P is

$$\nabla f(a_1, a_2, \dots, a_n) \cdot \frac{\nabla f(a_1, a_2, \dots, a_n)}{\|\nabla f(a_1, a_2, \dots, a_n)\|} = \frac{\|\nabla f(a_1, a_2, \dots, a_n)\|^2}{\|\nabla f(a_1, a_2, \dots, a_n)\|} = \|\nabla f(a_1, a_2, \dots, a_n)\| \\ = \|\nabla f(a_1, a_2, \dots, a_n)\|$$

Similarly, If ∇f exists and is continuous in an open ball around the point $(a_1, a_2, \dots, a_n)$, then the minimum value amongst all directional derivatives at P is

$$\nabla f(a_1, a_2, \dots, a_n) \cdot \frac{-\nabla f(a_1, a_2, \dots, a_n)}{\|\nabla f(a_1, a_2, \dots, a_n)\|} = -\frac{\|\nabla f(a_1, a_2, \dots, a_n)\|^2}{\|\nabla f(a_1, a_2, \dots, a_n)\|} = -\|\nabla f(a_1, a_2, \dots, a_n)\| \\ = -\|\nabla f(a_1, a_2, \dots, a_n)\|$$

1 point

Find the gradient of the scalar valued function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$.
 $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$.

☐

$(6x^2, 2y, 12z)$

☐



$$(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)$$

☐

$$(6x^2 + y^2, 2xy + 2y, 12z)(6x^2 + y^2, 2xy + 2y, 12z)$$

☐

$$(6x^2 + yz, 2y + xz, 12z + xy)(6x^2 + yz, 2y + xz, 12z + xy)$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)(6x^2 + y^2 + yz, 2xy + 2y + xz, 12z + xy)$$

1 point

Find the directional derivative of $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$

$f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ at $(2, 1, 2)(2, 1, 2)$ in the direction of a vector $(0, 3, 0)(0, 3, 0)$.

☐

3030

☐

00

☐

33

☐

1010

No, the answer is incorrect.

Score: 0

Feedback:

Accepted Answers:

1010

1 point

Find the direction in which the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$

$f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ increases the fastest from the point $(1, 1, 1)(1, 1, 1)$?

☐

$(1, 1, 1)(1, 1, 1)$

☐

$(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$

☐

$\sqrt{1, 3}$

☐

$\sqrt{1, 3}$

☐

1)

☐

$(8, 5, 13)(8, 5, 13)$

☐

$(\frac{8}{\sqrt{258}}, \frac{5}{\sqrt{258}}, \frac{13}{\sqrt{258}})(\frac{8}{\sqrt{258}}, \frac{5}{\sqrt{258}}, \frac{13}{\sqrt{258}})$

☐

$\sqrt{8, 258}$

☐

$\sqrt{5, 258}$

☐

$\sqrt{13}$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$\left(\frac{8}{\sqrt{258}}, \frac{5}{\sqrt{258}}, \frac{13}{\sqrt{258}}\right) \quad 258$$

√

$$8, \quad 258$$

√

$$5, \quad 258$$

√

$$13)$$

1 point

Find the maximum directional derivative of the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ at the point $P = (1, 2, 3)$.

○

$$\sqrt{1821} \quad 1821$$

√

○

$$4343$$

○

$$\sqrt{14} \quad 14$$

√

○

$$155155$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$\sqrt{1821} \quad 1821$$

√

1 point

Find the minimum directional derivative of the function $f(x, y, z) = 2x^3 + xy^2 + 6z^2 + y^2 + xyz$ at the point $P = (1, 0, 0)$.

○

$$-1-1$$

○

$$66$$

○

$$-6-6$$

○

$$11$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$-6-6$$

1 point

Let $f(x, y) = x^2 y^2 + y^3$. Find the directional derivative of f at the point $(1, 1)$ in the direction which forms an angle (counterclockwise) of $\frac{\pi}{3}$ with the positive x -axis.

○

$$\sqrt{29} \quad 29$$

✓

○

$$\sqrt{29} + \sqrt{2} \quad 29$$

✓

$$+ \quad 2$$

✓

○

$$\sqrt{2} \quad 2$$

✓

○

$$1 + \frac{5\sqrt{3}}{2} + 25 \quad 3$$

✓

No, the answer is incorrect.

Score: 0

Feedback:

Accepted Answers:

$$1 + \frac{5\sqrt{3}}{2} + 25 \quad 3$$

✓

Key Points: 2Key Points: 2

- Let $f(x, y)$ be a function defined on a domain D in R^2 containing some open ball around the point (a, b) . Equations of the tangent lines to the function f at the point (a, b) in the direction of unit vector (u_1, u_2) :

i) Parametric equations: $x(t) = a + tu_1$, $y(t) = b + tu_2$, $z(t) = f(a, b) + tf_u(a, b)$

ii) Symmetric equations: $\frac{x-a}{u_1} = \frac{y-b}{u_2} = \frac{z-f(a,b)}{f_u(a,b)}$

iii) Vector form: $(x(t), y(t), z(t)) = (a, b, f(a, b)) + t(u_1, u_2, f_u(a, b))$

- The equation of the tangent plane of the graph of a function of two variables $z = f(x, y)$ at the point (x_0, y_0, z_0) (where, $z_0 = f(x_0, y_0)$) is given by

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) - 1(z - z_0) = 0$$

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) + f(x_0, y_0) = z$$

- Linear approximation of the function $z = f(x, y)$ at the point (x_0, y_0, z_0) is given by

$$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) \quad L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

- Let $z = f(\tilde{x}) = f(x_1, x_2, \dots, x_n)$ $z = f(\tilde{x}) = f(x_1, x_2, \dots, x_n)$ be a function defined on a domain DD in R^2 R^2 containing some open ball around the point $\tilde{a} = (a_1, a_2, \dots, a_n)$ $\tilde{a} = (a_1, a_2, \dots, a_n)$. Equations of the tangent lines to the function f at the point $\tilde{a}$ in the direction of unit vector $\tilde{u} = (u_1, u_2, \dots, u_n)$ $\tilde{u} = (u_1, u_2, \dots, u_n)$:

i) Parametric equations: Parametric equations: $x_i(t) = a_i + t u_i$ $x_i(t) = a_i + t u_i$,
 $z(t) = f(\tilde{a}) + t f_{\tilde{u}}(\tilde{a})$ $z(t) = f(\tilde{a}) + t f_{\tilde{u}}(\tilde{a})$.

ii) Vector form: Vector form: $(\tilde{x}(t), z(t)) = (\tilde{a}, f(\tilde{a})) + t(\tilde{u}, f_{\tilde{u}}(\tilde{a}))$. $(\tilde{x}(t), z(t)) = (\tilde{a}, f(\tilde{a})) + t(\tilde{u}, f_{\tilde{u}}(\tilde{a}))$.

- Let $f(\tilde{x})$ $f(\tilde{x})$ be a function defined on domain DD in R^n R^n containing some open ball around the point $\tilde{a}$. Suppose ∇f exists and is continuous on some open ball around the point $\tilde{a}$. Equation of tangent plane to f at $\tilde{a}$:

$$z = f(\tilde{a}) + \nabla f(\tilde{a}) \cdot (\tilde{x} - \tilde{a}) \quad z = f(\tilde{a}) + \nabla f(\tilde{a}) \cdot (\tilde{x} - \tilde{a})$$

$$z = f(\tilde{a}) + \sum \frac{\partial f}{\partial x_i}(\tilde{a})(x_i - a_i) \quad z = f(\tilde{a}) + \sum \frac{\partial f}{\partial x_i}(\tilde{a})(x_i - a_i)$$

1 point

Find the equation of the tangent plane to the graph of the function given by

$$z = f(x, y) = \sqrt{r^2 - x^2 - y^2} \quad z = f(x, y) = \sqrt{r^2 - x^2 - y^2}$$

at the point $(0, 0, r)$.

☐

$$x + y + z = r \quad x + y + z = r$$

☐

$$2x + 2y + 2z = r \quad 2x + 2y + 2z = r$$

☐

$$z = r \quad z = r$$

☐

$$x + y = r \quad x + y = r$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$z = r \quad z = r$$

1 point

Find the equation of the tangent plane to the graph of the function given by

$$z = f(x, y) = \frac{1}{\sqrt{c}} \sqrt{d - ax^2 - by^2} \quad z = f(x, y) = \frac{1}{\sqrt{c}} \sqrt{d - ax^2 - by^2}$$

☐

$$1 \quad d - ax^2 - by^2$$

$\sqrt{\quad}$
at the point $(1, 2, f(1, 2))(1, 2, f(1, 2))$.

☐

$$ax + 2by + \sqrt{c(d - a - 4b)} z = 0ax + 2by + \quad c(d - a - 4b)$$

$\sqrt{\quad}$

$$z = 0$$

☐

$$ax + 2by + \sqrt{c(d - a - 4b)} z = d ax + 2by + \quad c(d - a - 4b)$$

$\sqrt{\quad}$

$$z = d$$

☐

$$2ax + 4by + \sqrt{c(d - a - 4b)} z = d 2ax + 4by + \quad c(d - a - 4b)$$

$\sqrt{\quad}$

$$z = d$$

☐

$$ax + 2by + \sqrt{c(d + a + 4b)} z = d ax + 2by + \quad c(d + a + 4b)$$

$\sqrt{\quad}$

$$z = d$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$ax + 2by + \sqrt{c(d - a - 4b)} z = d ax + 2by + \quad c(d - a - 4b)$$

$\sqrt{\quad}$

$$z = d$$

1 point

Find the equation of the tangent plane to the graph of the function $z = f(x, y) = x^2 yz = f(x, y) = x^2 y$ at the point $(1, 1, 1)(1, 1, 1)$.

☐

$$2x + y - z = 22x + y - z = 2$$

☐

$$2x + y + z = 22x + y + z = 2$$

☐

$$2x + y - z = 02x + y - z = 0$$

☐

$$2x + y = 22x + y = 2$$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$$2x + y - z = 22x + y - z = 2$$

Key Points: 3Key Points: 3

- Definition:Definition: Consider a scalar valued multivariable function $f: R^n \rightarrow Rf: R^n \rightarrow R$. A point $(a_1, a_2, \dots, a_n) \in R^n(a_1, a_2, \dots, a_n) \in R^n$ is said to be a critical point of the function f if $\nabla f(a_1, a_2, \dots, a_n) = 0\nabla f(a_1, a_2, \dots, a_n) = 0$ or at least one of $f_{x_i}(a_1, a_2, \dots, a_n)f_{x_i}(a_1, a_2, \dots, a_n)$ for $i = 1, 2, \dots, ni = 1, 2, \dots, n$ does not exist.
- Steps to find critical points of a function,

i) Find the partial derivatives f_{x_i} for $i = 1, 2, \dots, n$. If one of $f_{x_i}(a_1, a_2, \dots, a_n) = 0$ for $i = 1, 2, \dots, n$ does not exist then $(a_1, a_2, \dots, a_n)$ is a critical point.

ii) Otherwise, find the all possible point $(a_1, a_2, \dots, a_n) \in \mathbb{R}^n$ such that $f_{x_i}(a_1, a_2, \dots, a_n) = 0$ for all $i = 1, 2, \dots, n$, simultaneously.

- A function $f(x_1, x_2, \dots, x_n)$ has a local minimum at the point $(a_1, a_2, \dots, a_n)$ if $f(x_1, x_2, \dots, x_n) \geq f(a_1, a_2, \dots, a_n)$ for all points $(x_1, x_2, \dots, x_n)$ in some open ball around $(a_1, a_2, \dots, a_n)$.
- A function $f(x_1, x_2, \dots, x_n)$ has a local maximum at the point $(a_1, a_2, \dots, a_n)$ if $f(x_1, x_2, \dots, x_n) \leq f(a_1, a_2, \dots, a_n)$ for all points $(x_1, x_2, \dots, x_n)$ in some open ball around $(a_1, a_2, \dots, a_n)$.
- Local extrema are critical points.
- Not all critical points are local extrema. Those critical points at which the gradient exists and is 0 but are not local extrema, are called saddle points.
- A global (or absolute) maximum/minimum is a point for which the function value is maximum/minimum amongst all points in the domain. It always exists when the domain is closed and bounded and can be found by comparing the function values at all critical points in the domain as well as on successive boundaries.

1 point

Let $f(x, y) = \frac{x^3}{3} + y^2 + 2xy - 6x - 3y + 4$. Which of the following options are true about f ?

☐

$f_y = 0$ is a straight line passing through the origin.

☐

$(-1, \frac{5}{2})$ and $(3, -\frac{3}{2})$ are critical points.

☐

$f_x(-1, \frac{5}{2}) = 0$

☐

$f_y(3, -\frac{3}{2}) = 2$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$(-1, \frac{5}{2})$ and $(3, -\frac{3}{2})$ are critical points.

$f_x(-1, \frac{5}{2}) = 0$

1 point

Choose the correct statement(s) about the maxima/minima of $f(x, y) = x^4 + y^4$.

☐

There are no critical points.

☐

There is exactly one critical point.

☐

$(0, 0)$ is a point of local minima.

No, the answer is incorrect.

Score: 0

Feedback:

Accepted Answers:

There is exactly one critical point.

$(0, 0)$ is a point of local minima.

1 point

Suppose $f(x, y) = x^2 + 2y^2 + 3xy - 5y$ be a scalar valued function. Which of the following is (are) critical point(s) of $f(x, y)$?

☐

$(-10, 15)$

☐

$(0, 0)$

☐

$(15, -10)$

☐

$(15, 10)$

No, the answer is incorrect.

Score: 0

Feedback:

$f_x(x, y) = 2x + 3y$ and $f_y(x, y) = 3x + 4y - 5$

Accepted Answers:

$(15, -10)$

1 point

Which of the following statements are correct?

☐

The minimum value of the function $f(x, y) = x^2 + y^2 + 8$ is 00.

☐

The minimum value of the function $f(x, y) = x^2 + y^2 + 8 - 4x - 4y$ is 00.

☐

The maximum value of the function $f(x, y) = -x^2 - y^2$ is 00.

☐

The maximum value of the function $f(x, y) = x^2 + y^2 - 2x + 1$ is 00.

No, the answer is incorrect.

Score: 0

Accepted Answers:

The minimum value of the function $f(x, y) = x^2 + y^2 + 8 - 4x - 4y$ is 00.

The maximum value of the function $f(x, y) = -x^2 - y^2$ is 00.

1 point

Find the point on the plane $2x - y + 2z = 16$ closest to the origin.

☐

$(\frac{32}{9}, -\frac{16}{9}, \frac{32}{9})$

☐

$(1, -1, \frac{13}{2})$

☐

$(\frac{32}{9}, 1, \frac{89}{18})$

☐

$(1, 0, -1)$

No, the answer is incorrect.

Score: 0

Feedback:

Accepted Answers:

22 10 22 2022 2 10 2022