



L6.2: Image and kernel of linear transformations



Transcript Language

English



Hello, and welcome to the Maths 2 component of the online B.Sc. program on data science.

In this video we are going to study the notion of the image and kernel

of linear transformations. So, remember previously that we have studied the notion of the null space,

so this is some, for a matrix. So, this is some analogous thing for a linear transformation.

So, for a matrix we know what is the row space or the column space and that is, the analogous notion



for a linear transformation is the image. So, let us start by, from the definitions.

So, let $f: V \rightarrow W$ be a linear transformation and define the kernel of f , which is denoted by $\ker f$

of f as kernel of f is all those vectors v , such that $f(v) = 0$. So, V and W are vector spaces,

f is a linear transformation and we are looking at all those vectors v so that $f(v) = 0$. Now,

it is easy to check that kernel of f , a kernel of f is actually a subspace of V , which means it is

a vector space in its own right. What that means is if you take a linear combination of elements of

kernel of f , they still belong to kernel of f . Define the image of f . This is denoted by

$\text{Im } f$. Sometimes we will use a little i and sometimes we will use capital I .

So, that may, that notation will be a bit vague. So, we will define it as all those vectors w .

So, that there exists v in V . So, that w is equal to $f(v)$. So, this is nothing but the usual

definition of the range of a function. If you have a function between two spaces or two sets, we

call the range of f as the set of values which the function takes. So, that this is exactly the range

of the function. So, image of f is another name for the range of the function f . And I think you

have studied range of a function in Maths 1. In case you have not, I just recall for you that

range of a function is the set of values which the function f takes.

So, anyway, if you did not follow what I just said, we will anyway look at some examples.

Before that, let me also comment on the previous slide that the image of f is also a subspace of W .

So, if two vectors are in the image, the sum is also in the image, because you can take, so if w_1

is f of v_1 and w_2 is f of v_2 , then you can look at f of v_1 plus v_2 that is exactly w_1 plus w_2

because remember f is a linear transformation and that means w_1 plus w_2 is also in the image.

Similarly, if you have α times w , where w is in the image,

then w is f of v . So, you look at α times v , then f of αv is α times f of v is

α times w . So, αw is also in the image. Both of these remember use, I mean, in the proofs

that I gave, the statements that I just made, the fact that it is a linear transformation was

crucially used. So, this is not true for functions and so on for other things.

So, let us look at some examples to get our idea straight. So, look at the function

from $\mathbb{R}^2$ to $\mathbb{R}^2$, f of x, y is $2x$ comma y . First of all, why is this a linear transformation,

because each coordinate is a linear combination. Remember that linear combinations, if each

coordinate is a linear combination that is exactly what a linear transformation is when it comes to

functions from $\mathbb{R}^n$ to $\mathbb{R}^m$. So, $2x$ is a linear transformation of x and y ,

and y is a linear transformation of x and y . So, this is indeed a

linear transformation. So, each of these are linear combinations.

So, this is a linear transformation. So, what is the kernel of f ? The kernel of f is

all those values so that f of that number x comma, that vector x comma y is 0 . 0 where, 0 in the

codomain $\mathbb{R}^2$. So, 0 in the vector space $\mathbb{R}^2$. So, 0 in $\mathbb{R}^2$ is 0 comma 0 . So, in the previous slide

also note that we said f of v is equal to 0 . By that 0 we mean the 0 in W . So, here we have,

we want to look for those x, y such that f of x, y is equal to 0 comma 0 . So, it is clear that if

$2x$ comma y is 0 comma 0 , then $2x$ is 0 and y is 0 , which will tell us that x is 0 and y is 0 .

So, the kernel is exactly 0 comma 0 . So, the kernel is the 0 element or the 0 vector in R^2 .

What is the image of f ? The image of f is the entire R^2 . Why is that? So, if you want to write

a vector W in R^2 as the image of some element, then what would you do? You would look at,

suppose w is equal u comma v , then you look at u by 2 comma v .

And if you apply f on u by 2 comma v , you get 2 times u by 2 , which is u and v . So, that tells us

that u comma v is in the image. So, every vector u comma v is in the image that means the image is

the entire space R^2 , the entire vector space. Consider the example, R^2 to R^2 , f of x, y is $2x$

comma 0 . Again, this is a linear transformation because each coordinate is a linear combination.

So, what is the image and what is the kernel? So, kernel of f is 0 comma y . Why is this?

Suppose f of x, y is equal to 0, we want to, from there we want to solve for x comma y .

So, that means $2x$ comma 0 is equal to 0 comma 0. That means $2x$ is 0 and 0 is equal to 0.

So, 0, 0 does not yield us anything on x comma y , any restriction on x comma y .

$2x$ equals 0 yields us the restriction x is equal to 0, that means y can be anything. So,

the kernel of f is 0 comma y . So, this is the entire y -axis. 0 comma y is the y -axis.

And what is the image of f ? Image of f is, well, you have f of x comma y is $2x$ comma 0. So,

now suppose you look for an element u comma v and you ask when can u comma v be equal to $2x$ comma 0.

So, the first thing you need is that v is 0. That means this element must, this vector must lie on

the x -axis. And then you have to ask whether on the x -axis it can take all possible values.

So, you get the equation u is $2x$, which if you solve, you get x is u by 2,

which means if you put x comma y equal to u by 2 comma 0,

then the image of that element f of x comma y is u comma 0. So, you get

that every element, every vector on the x-axis is indeed in the image. And you already saw that

the image is contained in the x-axis that means the image is equal to the x-axis.

So, let us go on to studying the

relation between the kernel and injectivity, which, so we remember that a function is

injective or one-one, if $f(v_1)$ is equal to $f(v_2)$ implies v_1 is equal to v_2 and for linear

transformations this is the same as demanding that $f(v) = 0$ implies $v = 0$. We prove this actually.

So, what does this have to do with the kernel? Well, the first one may, you may not be,

it may not be clear, but the second statement makes it clear what it has to do with the kernel.

$f(v) = 0$ implies $v = 0$. So,

if a linear transformation is one-one or injective then the kernel must be $\{0\}$. That is what we are

saying, because if $f(v) = 0$, then v must be 0 . So, the only possible choices of vectors which

yield that $f(v) = 0$ is the 0 vector itself. So, if f is one-one or injective, then the kernel of f

is 0. On the other hand, if the kernel is 0, that means whenever $f(v)$ is 0, v is 0, so that means

the condition for being one-one is satisfied. So, kernel of f is 0 also means that

the linear transformation is one-one. So, being the linear transformation being one-one

is equivalent to kernel of f is 0. That is what we have proved here. So, linear transformation f is

one-one if and only if kernel of f is 0. So, what do I mean by kernel of f is 0. What I mean here

is that it is a 0 subspace. So, this should be better written as kernel of f is the 0 subspace,

which means it consists of the set 0. And remember, 0 itself is a

vector space. It satisfies all the properties of a vector space. And it is a subset of V .

So, it is a subspace of V . So, kernel of f is the 0 subspace. That is what we mean by

these two statements here. Kernel of f is 0 and kernel of f is 0. So, be careful when we

write this. The 0 in this case here on the right is a 0 subspace and not the 0 vector.

It means it is a set consisting of the 0 vector.

So, let us again see the relation like kernel is related to being injective let us study what

is the relation between the image and the surjectivity of a linear transformation.

So, recall that a function $f: V \rightarrow W$ is onto or surjective if for each w in W there is some

v in V such that $f(v) = w$. In other words, the range of this function is the entire W . So,

this is actually, this part is the same for any function. This is not, this has nothing to do with

linear transformations per se, what the statement in this slide.

So, it follows from the definition that a function f being onto is equal to range of f being W that

is exactly what we just commented. So, for linear transformations, in particular, this means that

because remember range is the same as image, only thing is when we say range we think of it as a set

and when we say image we think of it as a subspace. This is the main difference.

Otherwise, there is no real difference between the range and image as such.

So, a linear transformation is onto if and only if the image of f is W . And this image of f is W is

being concluded as a subspace. Image of f is actually a subspace of W always and we say that

it is onto. We have proved that it is onto, is equivalent to the subspace image of f actually

been the entire space. But this is not very different from what happened for functions.

Let us now proceed and make the connection between kernels and null spaces. So, so far

there is no matrix in the picture. So, now, we are going to introduce some matrix in the picture

and we will associate the kernel of this linear transformation with that matrix.

So, let $f: V \rightarrow W$ be a linear transformation,

let β be $v_1, v_2, \dots, v_n$ and γ be $w_1, w_2, \dots, w_m$ which are both, both of which are

ordered basis. β is an ordered basis of V and γ is an ordered basis for W .

So, now, indeed we can talk about a matrix, because remember in the previous video,

we have talked about how we, if you have a linear transformations along with

an ordered basis for each of V and W , there is a corresponding matrix. And how do we write down

that matrix? That write, we write down that matrix by looking at f of v_i and expressing that in terms

of w_j s, as a linear combination of the w_j s and whatever coefficients we get go into the matrix.

So, the coefficient of f , coefficients of f of v_i go into the i th column of that matrix.

So, let A be the corresponding matrix. So, just recall that for v in,

so suppose v is $\sum c_j v_j$ then f of v is equal to $\sum c_j$,

f of v_j that is by the linear transformation property, axioms of being a linear transformation.

And then, f of, so $\sum f$ of v_j can be further written as $\sum A_{ij} w_i$. So, f of v_j

is $\sum A_{ij} w_i$. So, that is corresponding the j th column of the matrix A , that is what these,

where these coefficients are coming from. That is exactly what we just commented.

So, when is f of v equal to 0? So, f of v is 0 if and only if each

coefficient in this linear combination of w_i s is 0. So, what is the coefficient of w_i ? So, you have

to take that $\sum i$ is equal 1 to m outside and take the $\sum j$ is equal to 1 to n inside

and then you will get summation $A_{ij}c_j$ over j that is the coefficient of w_i . And then we know that

if you have a linear combination of w_i which is 0 then each coefficient must be 0. Why is that,

because w_i are a basis, remember, and basis satisfies that it is linearly independent.

So, if summation of some linear combination, this particular linear combination of

w_i is 0 then each coefficient is 0. So, $f(v) = 0$ is exactly the same as saying that

summation $A_{ij}c_j$ is equal to 0. And now this equation should ring a bell in your mind.

So, the left hand side is saying, so $f(v) = 0$ means this is exactly the definition of

v being in the kernel of f . And the right hand side is saying that if you look at the vector c

consisting of c_1, c_2, c_n , then that vector is in the null space of A .

So, the relation we have between the kernel and the null space is the following.

That if you have a vector v , which when you express as a linear combination of the v_j s,

there, remember, it is an ordered basis, so you can write it as summation $c_j v_j$ in a unique way.

So, this vector is in the kernel if and only if this happens precisely when if you write these

coefficients of the v_j s as a column, then this column must be in the null space of

the matrix A . So, that is the relation between kernels and null spaces.

So, let us similarly describe what is the relation between images and column spaces? So, let V be a,

$f: V \rightarrow W$ be a linear transformation, let β be $v_1, v_2, \dots, v_n$, γ be $w_1, w_2, \dots, w_m$ both of these

are ordered basis, β for V and γ for W and let A be the matrix corresponding to f with

respect to β and γ . So, we know how to write this. The j th column of A corresponds to

the coefficients arising as expressing f of v_j in terms of w_i s.

So, now, again recall that for v is $\sum v_j$, $c_j v_j$ in V , f of v is $\sum c_j \sum A_{ij} w_i$,

which if you take the summations, interchange the summations we can do that because

it is a finite sum then you get that this is $\sum_{i=1}^m \left(\sum_{j=1}^n c_j A_{ij} \right) w_i$,

$\sum_{j=1}^n c_j \sum_{i=1}^m A_{ij} w_i$. So, the coefficient of w_i is $\sum_{j=1}^n c_j A_{ij}$ over j .

So, suppose now w is $\sum d_i w_i$ and

it is, so this is a vector in W . So, these d_i s are unique, because w_i s form a basis. So,

you can write w as $\sum d_i w_i$. So, when does this lie in the image.

So, this lies in the image precisely when there exists a scalar c_j so that $\sum A_{ij} c_j$ is d_i .

So, the scalars from the coefficients of something in the image, something of the form $f(v)$

is $\sum A_{ij} c_j$. So, that means these d_i s should be $\sum A_{ij} c_j$ in order for

w to be $f(v)$, where, what is v ? v is equal to $\sum c_j v_j$.

So, what that means is, we can now express this in terms of

columns. What it means is that $\sum d_i w_i$ is in the image of f if there exists a column vector c

such that the column vector d , which is $d_1, d_2, \dots, d_m$ is equal to A times c .

So, but remember A times c is exactly a linear combination of the columns of A .

So, what that means is that

w is in the image of f where w is given by $\sum d_i w_i$ if and only if you

look at the vector d is equal to $d_1, d_2, \dots, d_m$ and this is in the column space of A .

So, that is a relation between the image of f and the column space of this matrix A .

Now, let us see if we can use this to find the basis

of the kernel and the image of a linear transformation. So, again let $f: V \rightarrow W$ be a

linear transformation, choose ordered basis β and γ for V and W , respectively.

Let A be the corresponding matrix. So, the point, I mean, the first point here is that

we can look at the kernel and the null space and what we have really shown is that

the kernel is isomorphic to the null space. That is what really we showed

in the slide on kernels and null spaces. Why is that? That is because for every element of

the kernel, we have a unique element of the null space. So, you can define a function and

it is easy to verify that this is indeed a linear transformation from the kernel to the null space.

And it is also equally clear that this function is both one-one and onto that something you can

check. So, that tells us that the kernel is actually isomorphic to the null space.

So, in particular, if you look at the vectors $c_{11}, c_{12}, c_{1n}, c_{21}, c_{22}, c_{2n}$ and c_{k1}, c_{k2}, c_{kn}

these form a basis for the null space of A precisely when v_1 , the corresponding v_1 primes,

v_2 primes, v_k primes lie in kernel of f , form a basis for kernel of f . And what is the relation

between v_i s and, v_i primes and these columns? The coefficients of v_i prime when expressed in

terms of v_j are exactly corresponding to the j th, the i th vector in this collection.

So, in other words, what is the main point? The main point is I can read off a basis

for the kernel if I am, if I know how to get a basis for the null space.

And indeed, we do know how to get a basis for the null space.

Similar to the previous situation, we can find the relation between images and column spaces

specifically about basis between them using the isomorphism that we have discussed earlier.

So, if we consider the column vectors $d_{11}, d_{12}, d_{1m}, d_{21}, d_{22}, d_{2m}, d_{r1}, d_{r2}, d_{rm}$

then these form a basis for the column space of A precisely when if you look at the corresponding

linear combinations of the w_i s, w_i prime is $\sum_{j=1}^m d_{ij} w_j$ these form

a basis for image of f . So, this follows because of the previous isomorphism that was discussed.

So, once we have this isomorphism note also that if you look at the columns of A

which form a spanning set of the column space, then these correspond to the images of f of v_i .

So, the images f of v_i which form a spanning set for image of f .

So, what is the point of all this? So, as we have noted before, we can convert the problem of

finding a basis for the image space to the problem of finding a basis for the column space.

And, well, that is something we know how to do, because to find a basis of the column space,

what do we do, we look at the columns, then we do row reduction and once it is in row, reduced row

echelon form, you look at the columns containing the pivot elements, and then the original columns,

those are the ones which form a basis for the column space. So, now, what this is telling us is,

once we know those columns which form a basis for the column space, you look at the corresponding

linear combinations of the w_j s and that will give you a basis for the image of f .

So, this video, we have seen the notion of image and kernel and we have discussed these

relationships between the kernel and the null space of the associated matrix once we fix basis,

ordered basis, and similarly, the relationship between the image of a linear transformation

and the column space of a matrix which we obtain by fixing ordered basis.

So, we have seen that these are isomorphic. You know the interplay between these two. And using

these we can get basis by considering these corresponding matrices that we obtain and then

obtaining basis for those using row reduction, which we know how to do from the last week. So,

in the next video, we will implement all of this in concrete examples. Thank you.