



## AQ9.3: Activity Questions 3 - Not Graded



This assignment will not be graded and is only for practice.

### Directional Derivatives:

#### Level 1:

1 point

Which of the following statements is(are) true?

☐

The directional derivative of  $f(x, y)$  at a point  $(a, b)$  in the direction of a unit vector  $u$  is a non negative real number.

☐

The directional derivative of any function is always a scalar.

☐

The rate of change of  $f(x_1, x_2, \dots, x_n)$  in the direction of a unit vector

$u = (a_1, a_2, \dots, a_n)$  is called the directional derivative of  $f$  in the direction of  $u$ .

☐

The directional derivative of any function is a vector because it is dependent on the direction of a unit vector.

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:

The directional derivative of any function is always a scalar.

The rate of change of  $f(x_1, x_2, \dots, x_n)$  in the direction of a unit vector

$u = (a_1, a_2, \dots, a_n)$  is called the directional derivative of  $f$  in the direction of  $u$ .

If the directional derivative of  $f(x, y) = x + xy - 2y^2$  at the point  $(2, 3)$  in the direction of the vector  $(2\sqrt{2}, 2\sqrt{2})$  is  $a\sqrt{2}$  where  $a \in \mathbb{R}$ , then find the value of  $a$ .

, 2

) is  $a\sqrt{2}$  where  $a \in \mathbb{R}$ , then find the value of  $a$ .

where  $a \in \mathbb{R}$ , then find the value of  $a$ .

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:

(Type: Numeric) -3

1 point

If the directional derivative of  $f(x, y, z) = xe^y + ye^z + ze^x$  at point

$(1, 0, 1)$  in the direction of a vector  $(2, -2, 1)$  is  $ke$  where  $k \in \mathbb{R}$ , then find the value of  $6k$ .

**No, the answer is incorrect.**

**Score: 0**

#### Accepted Answers:

(Type: Numeric) 2

1 point

1 point

Suppose  $f$  is a multivariable function defined on domain  $\mathbb{R}^2$ . Which of the following is equal to the rate of change of  $f$  at point  $(x_1, y_1)$  in the direction of a vector  $u = (u_1, u_2)$

☐


$$\lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(u_1, u_2)) + f(x_1, y_1)}{h} \rightarrow 0 \lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(u_1, u_2)) + f(x_1, y_1)}{h}$$

○

$$\lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(u_1, u_2)) - f(x_1, y_1)}{h} \rightarrow 0 \lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(u_1, u_2)) - f(x_1, y_1)}{h}$$

○

$$\lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) - f(x_1, y_1)}{h} \rightarrow 0 \lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) - f(x_1, y_1)}{h}$$

√

$$u_1, \frac{u_2}{\sqrt{u_1^2 + u_2^2}}$$

√

$$u_2)) - f(x_1, y_1)$$

○

$$\lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) + f(x_1, y_1)}{h} \rightarrow 0 \lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) + f(x_1, y_1)}{h}$$

√

$$u_1, \frac{u_2}{\sqrt{u_1^2 + u_2^2}}$$

√

$$u_2)) + f(x_1, y_1)$$

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

$$\lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) - f(x_1, y_1)}{h} \rightarrow 0 \lim_{h \rightarrow 0} \frac{f((x_1, y_1) + h(\frac{u_1}{\sqrt{u_1^2 + u_2^2}}, \frac{u_2}{\sqrt{u_1^2 + u_2^2}})) - f(x_1, y_1)}{h}$$

√

$$u_1, \frac{u_2}{\sqrt{u_1^2 + u_2^2}}$$

√

$$u_2)) - f(x_1, y_1)$$

If the directional derivative of  $f(x, y) = x \sin(y)$  at the point  $(1, 0)$  in the direction of a unit vector  $(u_1, u_2)$  ( $u_1, u_2 \in \mathbb{R}$ ) is 0.8, then find the absolute value of  $u_1$ .

(Enter your answer upto one decimal place only)

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

(Type: Range) 0.5, 0.7

1 point

**Level 2:**

Consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined as:  $f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & x, y \neq 0 \\ 0 & x = y = 0 \end{cases}$  Find the directional derivative of  $f$  at  $(0, 0)$  in the direction of the

$$\text{vector } u = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$$

√

, 21). (Enter your answer upto one decimal place only)

**No, the answer is incorrect.****Score: 0**

**Accepted Answers:**

(Type: Range) 1.4,1.6

1 point

Suppose  $d$  is the directional derivative of a scalar valued multivariable function  $f(x, y) = x^2 + y^2$  at a point  $P(1, 2)$  in the direction of the line  $PQ$ , where point  $Q$  is at  $(4, 6)$ .

The value of  $d$  is (Enter your answer upto one decimal place only)

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

(Type: Range) 4.3,4.5

1 point

Suppose  $f(x, y, z) = ax^2y + yz^2 - z^2x$ , where  $a \in \mathbb{R}$ , is a scalar valued multivariable function defined on domain  $D \subset \mathbb{R}^3$ . For what value of  $a$ , does the directional derivative of the given function at the point  $(1, 1, 1)$  in the direction of the vector  $u = (1, 2, -2)$  equal 33?

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

(Type: Numeric) 2

1 point

The temperature at a point  $(x, y, z)$  is given by

$$T(x, y, z) = 10e^{-x^2+y^2-4z^2}$$

where  $T$  is measured in  $^{\circ}C$  and  $x, y, z$  are measured in meters. If the rate of change of temperature at  $(2, -1, 0)$  in the direction of a vector  $u = (1, -1, 1)$  is  $k \cdot \frac{e^{-3}}{\sqrt{3}}$  where  $k \in \mathbb{R}$ , then the value of  $k$  is (approximate to two decimal places)

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

(Type: Numeric) -20

1 point

Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be given by  $f(x, y) = x^2 + y^2 - 2xy$ . Suppose  $D_u f(x_0, y_0)$  is the directional derivative at  $(x_0, y_0)$  in the direction of the unit vector  $u = \left(\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}\right)$ .

$\sqrt{\frac{1}{2}}$

$-1$ ) and  $D_v f(x_0, y_0)$  is the directional derivative at  $(x_0, y_0)$  in the direction of the unit vector  $v = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ .

$\sqrt{\frac{1}{2}}$

). Consider a matrix  $A = \begin{bmatrix} D_u f(1, 2) & D_u f(2, 1) \\ D_v f(1, 2) & D_v f(2, 1) \end{bmatrix}$ . What is the value of  $\det(A)$ ?

**No, the answer is incorrect.****Score: 0****Accepted Answers:**

(Type: Numeric) 0

1 point

[Check Answers](#)

Your score is: 0/10