Practice Assignment 4 - Not Graded

This assignment will not be graded and is only for practice.

Multiple Select Questions (MSQ)

1 point
Match the vector spaces (with the usual scalar multiplication and vector addition as in $M_{3\times 3}(R)$M3×3 (R)) in column A with their bases in column B in Table : M2W4P1.

Choose the correct option.

- [ ] a → → i
- [ ] a → → ii
- [ ] b → → iii
- [ ] c → → iii
- [ ] c → → i

No, the answer is incorrect.
Score: 0
Accepted Answers:
a → → ii
b → → iii
c → → i

1 point
Match the sets of vectors in column A with their properties of linear dependence or independence in column B and the dimension of the vector spaces in column C spanned by the sets.

- [ ] a → → ii → → 4, b → → i → → 3
- [ ] c → → iv → → 1, d → → iii → → 2
- [ ] c → → iv → → 2, d → → iii → → 1

No, the answer is incorrect.
Score: 0
Accepted Answers:
a → → ii → → 4, b → → i → → 3
c → → iv → → 1, d → → iii → → 2

1 point
Which of the following options are true?

- [ ] The number of linearly independent vectors in a vector space can be more than the dimension of the vector space.
- [ ] Dimension of the vector space spanned by the vectors $\begin{bmatrix}1 & -1\\ 1 & 1\end{bmatrix}$[11−11], $\begin{bmatrix}-1 & 1\\ -1 & 1\end{bmatrix}$[−11−111], and $\begin{bmatrix}-1 & 1\\ 1 & 1\end{bmatrix}$[−11111] is 2.

☐
Dimension of the vector space spanned by the vectors $\begin{bmatrix}1 & -1\\ 1 & 1\end{bmatrix}\begin{bmatrix}1 & 1\\ -1 & 1\end{bmatrix}$, $\begin{bmatrix}-1 & 1\\ -1 & -1\end{bmatrix}\begin{bmatrix}-1 & -1\\ 1 & -1\end{bmatrix}$, and $\begin{bmatrix}-2 & 2\\ -2 & -2\end{bmatrix}\begin{bmatrix}-2 & -2\\ 2 & -2\end{bmatrix}$ is 1.
☐ The number of spanning vectors can be more than the dimension of the vector space.

No, the answer is incorrect.
Score: 0
Accepted Answers:
Dimension of the vector space spanned by the vectors $\begin{bmatrix}1 & -1\\ 1 & 1\end{bmatrix}\begin{bmatrix}1 & 1\\ -1 & 1\end{bmatrix}$, $\begin{bmatrix}-1 & 1\\ -1 & -1\end{bmatrix}\begin{bmatrix}-1 & -1\\ 1 & -1\end{bmatrix}$, and $\begin{bmatrix}-2 & 2\\ -2 & -2\end{bmatrix}\begin{bmatrix}-2 & -2\\ 2 & -2\end{bmatrix}$ is 1.
The number of spanning vectors can be more than the dimension of the vector space.

1 point
Let AA be a nonzero $2 \times 2$2 × 2 matrix. Which of the following options are true?
☐
The rank of AA must be strictly less than 22.
☐
If the rank of the matrix is 2, then the dimension of the vector space spanned by the vectors corresponding to each column of AA, must be 2.
☐
The rank of AA must be at least 1.
☐
The rank of AA may be 0.
☐
If the rank of the matrix is $2 - 12 - 1$, then there exists one vector corresponding to a column of AA, which can be expressed as a linear combination of the vectors corresponding to each of the remaining columns of AA.

No, the answer is incorrect.
Score: 0
Accepted Answers:
If the rank of the matrix is 2, then the dimension of the vector space spanned by the vectors corresponding to each column of AA, must be 2.
The rank of AA must be at least 1.
If the rank of the matrix is $2 - 12 - 1$, then there exists one vector corresponding to a column of AA, which can be expressed as a linear combination of the vectors corresponding to each of the remaining columns of AA.

Numerical Answer Type (NAT)

Suppose $A = \begin{pmatrix}8-m & 2 & 4\\ 3 & 0 & 0\\ m & m-2 & \end{pmatrix}$A = $\begin{pmatrix}8-\text{m} & 3\text{m} & 20 & \text{m} & 4 & 0 & -2\end{pmatrix}$. For what value of mm is the rank of AA at most 2?

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) -1.0
1 point

If rank of the matrix $\begin{bmatrix} 0 & -1 & a \\ 2 & 0 & -4 \\ 3 & -9 & -6 \end{bmatrix}\begin{bmatrix} 0 & 2 & 3 \\ -1 & 0 & -9 \\ a & -4 & -6 \end{bmatrix}$ is 2 then find the value of aa.

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 0
1 point

Find out the value of aa for which the matrix $\begin{bmatrix} a & -5 \\ 0 & -1 \end{bmatrix}$[a0−5−1] will be in the spanning set of the matrices $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$[100−1] and $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$[0010] in $M_{2\times 2}(R)$M2×2(R) with usual matrix addition and scalar multiplication.

No, the answer is incorrect.
Score: 0
Accepted Answers:
(Type: Numeric) 1
1 point

Comprehension Type Question:

Suppose in a village there are four farmers A, B, C and D, each owning 1 acre of land. They cultivate paddy, pulses and/or sugarcane in their lands as follows: Farmer A uses 50% of his land for paddy, 30% for pulses and the remaining for sugarcane. Farmer B uses 40% of her land for paddy and she divides her remaining land equally for pulses and sugarcane. Farmer C uses the whole land for paddy only, and Farmer D uses 30% for paddy, 30% for pulses and the remaining for sugarcane. Using the above information, answer the following questions.

1 point
Suppose the area used by a farmer for different crops is denoted by a row vector. Let S be the span of the resulting four row vectors . Choose the correct set of options.
☐ The row vectors corresponding to the area used for different crops by Farmer D can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmers A, B, and C.
☐ The row vectors corresponding to the area used for different crops by Farmer D can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmer A and Farmer B.
☐ The row vectors corresponding to the area used for different crops by Farmer C can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmer A and Farmer B.
☐ The row vectors corresponding to the area used for different crops by Farmer D can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmer A and Farmer C.
No, the answer is incorrect.
Score: 0
Accepted Answers:

The row vectors corresponding to the area used for different crops by Farmer D can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmers A, B, and C.

The row vectors corresponding to the area used for different crops by Farmer D can be obtained as a linear combination of the row vectors corresponding to the area used for different crops by Farmer A and Farmer B.

1 point

Let SS be the vectors space defined in the previous question, with the usual addition and scalar multiplication on R^3 R3. Which of the following sets will not be a basis of SS?

- [ ] {(5, 3, 2)}{(5, 3, 2)}
- [ ] {(5, 3, 2), (4, 3, 3)}{(5, 3, 2), (4, 3, 3)}
- [ ] {(5, 3, 2), (4, 3, 3), (10, 0, 0)}{(5, 3, 2), (4, 3, 3), (10, 0, 0)}
- [ ] {(5, 3, 2), (4, 3, 3), (3, 3, 4)}{(5, 3, 2), (4, 3, 3), (3, 3, 4)}
- [ ] {(5, 3, 2), (4, 3, 3), (10, 0, 0), (3, 3, 4)}{(5, 3, 2), (4, 3, 3), (10, 0, 0), (3, 3, 4)}

No, the answer is incorrect.

Score: 0

Accepted Answers:

{(5, 3, 2)}{(5, 3, 2)}

{(5, 3, 2), (4, 3, 3)}{(5, 3, 2), (4, 3, 3)}

{(5, 3, 2), (4, 3, 3), (3, 3, 4)}{(5, 3, 2), (4, 3, 3), (3, 3, 4)}

{(5, 3, 2), (4, 3, 3), (10, 0, 0), (3, 3, 4)}{(5, 3, 2), (4, 3, 3), (10, 0, 0), (3, 3, 4)}

Check Answers