

Cosmological Implications of the GWTC-3 Modified-Propagation Anomaly: Early-Anchor Recalibration, CMB-Lensing Refit, and Early-Universe H_0 Inference Bias

Aiden B. Smith
Independent Researcher
 (Dated: February 10, 2026)

We project the GWTC-3 O3 modified-gravity propagation posterior into late- and early-universe inference tests using an updated Planck-facing calibration chain. The upstream O3 anomaly replication repository is archived on Zenodo (DOI: 10.5281/zenodo.18585598). A 60-restart Planck+MG global refit defines an updated early anchor, $H_0^{\text{Planck, MG}} = 68.005$ (p50), $\Omega_m^{\text{Planck, MG}} = 0.30643$ (p50), and $A_{\text{lens}} = 1.0428$ (p50).

After rebasing late-time transfer sweeps to this anchor, the constrained anchor-relief posterior is $\mathcal{R}_{\text{anchor}}^{\text{GR}} = 0.1545$ (mean; p16/p50/p84 = 0.108/0.147/0.189). A 10-case robustness grid gives posterior-shift relief mean 0.5296 (p50 0.5125), while a joint SN+BAO+CC transfer model yields $\mathcal{R}_{\text{joint}} = 0.8329$ (mean) but no Bayes preference for explicit transfer terms ($\log \text{BF} = -0.533$).

For CMB lensing, direct CAMB propagation of MG draws still predicts baseline suppression at $L \sim 100$ and $L \sim 300$ (medians -15.29% and -9.49%). However, an MG-aware lensing refit recovers good fits (median $\chi^2 = 8.06$ vs Planck-reference 9.04; 100% of refit draws better than reference) with median $M_\star^2(z=0)/M_\star^2(z \gg 1) \simeq 0.901$. Compressed- θ_\star inversions under GR assumptions bias inferred H_0 upward (mean $\Delta H_0 = +1.88 \text{ km s}^{-1} \text{ Mpc}^{-1}$ for fixed Ω_m , or $+4.55$ for lensing-proxy Ω_m), indicating that early-universe inference assumptions can materially shift recovered expansion rates in this MG-conditioned scenario.

I. SCOPE AND FRAMING

This work treats the O3 modified-propagation signal phenomenologically: given the inferred posterior, what cosmological consequences follow? The O3 anomaly analysis and data products are archived on Zenodo [1]. We do not re-argue detection significance in this manuscript.

Modified GW propagation has been explored in theory-forward frameworks [14, 15]. Here, we use a data-driven posterior and update the pipeline to answer three questions in one chain:

1. How much late-time Hubble tension relief remains after recalibrating the early anchor?
2. Does Planck 2018 lensing necessarily reject this posterior, or can an MG-aware refit absorb the suppression?
3. How much can GR-based early-universe compression bias inferred H_0 if MG truth is assumed?

II. PIPELINE SUMMARY

Posterior draws are taken from `outputs/finalization/highpower_multistart_v2/M0_start101` and propagated through four linked stages:

1. **Global Planck+MG recalibration:** 60-restart multistart fit (cpuset 0-59) to establish updated early-anchor reference values.
2. **Late-time rebasing:** constrained/pilot transfer sweeps are rebased to the new Planck-like anchor and recompressed into a final relief posterior.

3. **CMB lensing forecasts:** baseline draw-level CAMB projection to Planck 2018 lensing bandpowers, followed by an MG-aware two-parameter lensing refit.

4. **Compressed early-universe inversion:** GR inversion of $\theta_\star = r_d/D_M(z_\star)$ under fixed- Ω_m and lensing-proxy- Ω_m assumptions.

These are targeted forecasts and refits, not a full MG TT/TE/EE perturbation-sector likelihood analysis.

III. RESULTS

A. Updated early anchor from the global Planck+MG fit

The 60-restart Planck+MG run completed all restarts with 5 converged minima and 55 max-evaluation exits. Using converged minima only, we obtain:

$$H_0^{\text{Planck, MG}} = 68.005302 \text{ (p50)}, \quad \Omega_m^{\text{Planck, MG}} = 0.30643039 \text{ (p50)}, \quad (1)$$

With local reference $H_0^{\text{local}} = 73.0$, the baseline gap used in rebased relief calculations is

$$\Delta H_0^{\text{base}} = \left| H_0^{\text{local}} - H_0^{\text{Planck, MG}} \right| = 4.994698. \quad (2)$$

B. Late-time relief after rebasing

After rebasing constrained transfer sweeps to the updated early anchor and applying Monte Carlo calibration:

$$\mathcal{R}_{\text{anchor}}^{\text{GR}} = 0.1545 \quad (\text{mean}; \text{p16/p50/p84} = 0.1085/0.1475/0.1891) \quad (3)$$

Independent robustness and joint-fit diagnostics are:

- 10-case robustness grid: posterior-shift relief mean 0.5296 (p50 0.5125, p84 0.5453), with zero failed cases.
- Joint SN+BAO+CC transfer fit: relief posterior mean 0.8329 (p50 0.8386), but

$$\log \text{BF}_{\text{transfer/no-transfer}} = -0.533, \quad (4)$$

so explicit transfer terms are not favored in this setup.




FIG. 1. Rebased relief sensitivity to high- z transfer bias from constrained and pilot sweeps. The final calibrated posterior is centered at $\mathcal{R}_{\text{anchor}}^{\text{GR}} \approx 0.15$.

C. CMB lensing: baseline suppression and MG-aware refit

Baseline draw-level CAMB projection against Planck 2018 lensing bandpowers (`consent8`, 64 draws) gives:

$$\left. \frac{C_L^{\phi\phi}(\text{MG})}{C_L^{\phi\phi}(\text{Planck ref})} \right|_{L \approx 106} = 0.847^{+0.091}_{-0.127}, \quad (5)$$

$$\left. \frac{C_L^{\phi\phi}(\text{MG})}{C_L^{\phi\phi}(\text{Planck ref})} \right|_{L \approx 286} = 0.905^{+0.068}_{-0.080}, \quad (6)$$

with median suppressions of -15.29% and -9.49% . The baseline fit quality is poor relative to the Planck-reference model:

$$\chi_{\text{MG,baseline}}^2 (\text{median}) = 51.77, \quad \chi_{\text{Planck ref}}^2 = 9.04, \quad (7)$$

and only 3.1% of draws outperform the reference. A 32-draw cross-check from an independent posterior sample is more discrepant (-18.66% at $L \approx 106$, -11.29% at $L \approx 286$; $p_{\text{better}} = 0$).

We then perform an MG-aware lensing refit (32 draws) with a phenomenological effective- M_\star^2 amplitude plus ℓ -tilt response. This removes the baseline mismatch:

$$\chi_{\text{MG refit}}^2 (\text{median}) = 8.06, \quad (8)$$

better than the Planck-reference $\chi^2 = 9.04$ in 100% of refit draws. The fitted median response corresponds to

$$\frac{M_\star^2(z=0)}{M_\star^2(z \gg 1)} \simeq 0.901 \quad (9)$$

(about a 9.9% drop), with small residual suppression at $L \approx 286$.




FIG. 2. Planck 2018 lensing bandpowers with baseline MG projection and MG-aware refit overlay. The refit absorbs the baseline suppression and restores near-reference fit quality.

D. Early-universe GR inversion biases inferred H_0 upward

Using the rebased early-anchor assumptions in compressed- θ_\star inversion:

- fixed $\Omega_m = \Omega_m^{\text{Planck, MG}}$: $H_{0, \text{inferred}}$ mean 72.394 (p50 73.170), with mean $\Delta H_0 = +1.876 \text{ km s}^{-1} \text{ Mpc}^{-1}$ relative to draw-level truth;
- lensing-proxy Ω_m : $H_{0, \text{inferred}}$ mean 75.065 (p50 75.226), with mean $\Delta H_0 = +4.547 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

Relative to the recalibrated early anchor $H_0^{\text{Planck, MG}} = 68.005$, the posterior medians shift by approximately:

$$\Delta H_0^{\text{truth}} \approx +2.39, \quad \Delta H_0^{\text{fixed inversion}} \approx +5.16, \quad \Delta H_0^{\text{lensing inversion}} \approx +2.39 \quad (10)$$




FIG. 3. Draw-level H_0 truth versus GR-inferred H_0 under compressed- θ_* inversion with fixed- Ω_m (left) and lensing-proxy- Ω_m (right). Both assumptions bias inferred H_0 upward, with larger shift in the lensing-proxy case.

IV. INTERPRETATION

The updated pipeline supports three conclusions.

First, recalibrating the Planck-like early anchor materially lowers the constrained anchor-relief estimate (now centered near ~ 0.15), while robustness-grid and joint-transfer diagnostics still indicate that meaningful upward H_0 shifts are available in broader transfer-conditioned settings.

Second, baseline CAMB propagation of the O3 MG posterior predicts lensing suppression, but this does not by itself imply exclusion once MG-aware response freedom is included: the phenomenological refit reaches χ^2 comparable to or better than the Planck-reference model.

Third, if MG truth holds while early-universe compression is interpreted under GR assumptions, inferred H_0 can be biased high by order $+2$ to $+5$ $\text{km s}^{-1} \text{Mpc}^{-1}$ relative to posterior truth, and by up to $\sim +7$ $\text{km s}^{-1} \text{Mpc}^{-1}$ relative to the recalibrated Planck-like anchor median.

Within this study scope, the largest systematic lever is not only late-time transfer, but also model-assumption bias in early-universe inference compression.

REPRODUCIBILITY

Core scripts used in this follow-up are:

- `scripts/run_planck_global_mg_refit_multistart.py`
- `scripts/rebase_bias_transfer_sweep_to_planck_ref.py`
- `scripts/run_hubble_tension_final_relief_posterior.py`
- `scripts/run_hubble_tension_mg_forecast_robustness_grid.py`
- `scripts/run_joint_transfer_bias_fit.py`
- `scripts/run_hubble_tension_cmb_forecast.py`
- `scripts/run_hubble_tension_mg_lensing_refit.py`
- `scripts/run_hubble_tension_early_universe_bias.py`

ACKNOWLEDGMENTS

This work relied extensively on A.I.-assisted tools for code development, pipeline execution support, figure generation, and manuscript drafting/editing.

DATA AVAILABILITY AND DOIS

The follow-up uses posterior products from the O3 anomaly pipeline and public cosmology datasets. Data provenance and DOIs are:

- O3 modified-gravity tension anomaly repository (Zenodo): DOI 10.5281/zenodo.18585598.

- O3 search-sensitivity injection data used in up-stream calibration (Zenodo): DOI 10.5281/zenodo.7890437.
- GWTC-3 catalog paper: DOI 10.1103/PhysRevX.13.041039.
- Pantheon+ cosmology constraints: DOI 10.3847/1538-4357/ac8e04.
- SH0ES local- H_0 reference: DOI 10.3847/2041-8213/ac5c5b.
- SDSS DR12 BOSS consensus BAO (source of `sdss_DR12Consensus_bao.dat`): DOI 10.1093/mnras/stx721.
- eBOSS DR16 cosmological compilation (source class for `sdss_DR16_LRG_BAO_DMDH.dat`): DOI 10.1103/PhysRevD.103.083533.
- DESI 2024 BAO cosmological constraints (source class for `desi_2024_gaussian_bao_ALL_GCcomb_mean.txt`): DOI 10.1088/1475-7516/2025/02/021.
- Cosmic-chronometer compilation components used in `Hz_BC03_all.dat`: DOIs 10.1088/1475-7516/2012/08/006, 10.1103/PhysRevD.71.123001, and 10.1088/1475-7516/2010/02/008.
- Planck 2018 cosmological-parameter and lensing references: DOIs 10.1051/0004-6361/201833910 and 10.1051/0004-6361/201833886.

-
- [1] A. B. Smith, “O3 Modified Gravity Tension Replication,” Zenodo (2026), DOI: 10.5281/zenodo.18585598.
 - [2] LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration, “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run — O3 search sensitivity estimates,” Zenodo (2023), DOI: 10.5281/zenodo.7890437.
 - [3] R. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration), “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run,” *Phys. Rev. X* **13**, 041039 (2023), DOI: 10.1103/PhysRevX.13.041039.
 - [4] D. Brout *et al.*, “The Pantheon+ Analysis: Cosmological Constraints,” *Astrophys. J.* **938**, 110 (2022), DOI: 10.3847/1538-4357/ac8e04.
 - [5] A. G. Riess *et al.*, “A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km s⁻¹ Mpc⁻¹ Uncertainty from the Hubble Space Telescope and the SH0ES Team,” *Astrophys. J. Lett.* **934**, L7 (2022), DOI: 10.3847/2041-8213/ac5c5b.
 - [6] S. Alam *et al.*, “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample,” *Mon. Not. R. Astron. Soc.* **470**, 2617 (2017), DOI: 10.1093/mnras/stx721.
 - [7] S. Alam *et al.*, “Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory,” *Phys. Rev. D* **103**, 083533 (2021), DOI: 10.1103/PhysRevD.103.083533.
 - [8] DESI Collaboration, “DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations,” *J. Cosmol. Astropart. Phys.* **02** (2025) 021, DOI: 10.1088/1475-7516/2025/02/021.
 - [9] M. Moresco *et al.*, “Improved constraints on the expansion rate of the Universe up to $z \sim 1.1$ from the spectroscopic evolution of cosmic chronometers,” *J. Cosmol. Astropart. Phys.* **08** (2012) 006, DOI: 10.1088/1475-7516/2012/08/006.
 - [10] J. Simon, L. Verde, and R. Jimenez, “Constraints on the redshift dependence of the dark energy potential,” *Phys. Rev. D* **71**, 123001 (2005), DOI: 10.1103/PhysRevD.71.123001.
 - [11] D. Stern *et al.*, “Cosmic chronometers: constraining

- the equation of state of dark energy. I: $H(z)$ measurements,” *J. Cosmol. Astropart. Phys.* **02** (2010) 008, DOI: 10.1088/1475-7516/2010/02/008.
- [12] N. Aghanim *et al.* (Planck Collaboration), “Planck 2018 results. VI. Cosmological parameters,” *Astron. Astrophys.* **641**, A6 (2020), DOI: 10.1051/0004-6361/201833910.
- [13] N. Aghanim *et al.* (Planck Collaboration), “Planck 2018 results. VIII. Gravitational lensing,” *Astron. Astrophys.* **641**, A8 (2020), DOI: 10.1051/0004-6361/201833886.
- [14] E. Belgacem, Y. Dirian, S. Foffa, and M. Maggiore, “Modified gravitational-wave propagation and standard sirens,” *Phys. Rev. D* **98**, 023510 (2018), DOI: 10.1103/PhysRevD.98.023510.
- [15] A. Nishizawa, “Generalized framework for testing gravity with gravitational-wave propagation,” *Phys. Rev. D* **97**, 104037 (2018), DOI: 10.1103/PhysRevD.97.104037.