

# Discrete Mathematics

Sina Aleyaasin

September 2, 2017

### **Abstract**

A compilation of notes based on *Discrete Mathematics and its Applications* (7<sup>th</sup> Edition) – K. Rosen.

# Contents

# Chapter 1

## Logic and Proofs

### 1.1 Propositional Logic

#### 1.1.1 Propositions

A proposition is a declarative sentence that has a truth value of true (T) or false (F). It may be denoted by a propositional variable (such as  $p, q, r, s$ ). Compound propositions can be made from one or more propositions by means of *logical connectives*. The definition of a logical connective can be expressed by its corresponding truth table.

#### 1.1.2 Logical Connectives

A logical connective operates on one or more propositions to yield a new proposition.

##### Negation $\neg$

Negation is a unary operation that yields the logical complement of a proposition.

$p$	$\neg p$
T	F
F	T

Table 1.1: Truth table for negation

##### Conjunction $\wedge$

The conjunction of two propositions is true if both propositions are true, and is false otherwise.

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Table 1.2: Truth table for conjunction

##### Disjunction $\vee$

The disjunction of two propositions is true if either or both are true, and is false otherwise.

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Table 1.3: Truth table for disjunction

##### Exclusive Disjunction $\oplus$

The exclusive disjunction of two propositions is true if one is the logical complement of the other, and is false otherwise.

#### 1.1.3 Conditional Statements

A conditional statement is a proposition composed from a *premise*  $p$  and a *conclusion*  $q$ . It represents

$p$	$q$	$p \oplus q$
T	T	F
T	F	T
F	T	T
F	F	F

Table 1.4: Truth table for exclusive disjunction

the implication of  $q$  by  $p$ , and is denoted

$$p \rightarrow q$$

The only case in which this statement is false is when  $p$  is true but  $q$  is false.

$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

Table 1.5: Truth table for a conditional statement

### Converse

The converse of a conditional statement  $p \rightarrow q$  is given by

$$q \rightarrow p$$

### Contrapositive

The contrapositive of a conditional statement is given by

$$\neg q \rightarrow \neg p$$

It is *logically equivalent* to the conditional statement itself; that is to say, the truth tables for a conditional statement and its contrapositive are identical.

### Inverse

The inverse of a conditional statement is given by

$$\neg p \rightarrow \neg q$$

Since a conditional statement is logically equivalent to its contrapositive, and the inverse is the contrapositive of the converse, then the inverse is thus logically equivalent to the converse.

## Biconditional Statements

A biconditional statement is the conjunction of a conditional statement and its converse. It is denoted by

$$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$$

and is only true when both  $p$  and  $q$  have the same truth value.

$p$	$q$	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

Table 1.6: Truth table for a biconditional statement.

### 1.1.4 Logical & Bit operations

A *bit* (portmanteau of binary and digit) is the smallest unit of information representable by a computer. The value of a bit can be either 1 or 0, analogous to the truth value of a proposition, T or F, respectively. One may consider bit operators analogous to logical operators.

Logical Connective	Bit Operator
$\vee$	OR
$\wedge$	AND
$\oplus$	XOR

Table 1.7: Correspondence between logical connectives and bit operators

## 1.2 Applications of Propositional Logic

### 1.2.1 Logic Gates

## 1.3 Propositional Equivalences

### 1.3.1 Compound Propositions

A compound proposition may be categorised into one of three categories:

### Tautology

A *tautology* is a compound proposition that is *true* in all possible cases. An arbitrary tautology is denoted by

$$\top$$

### Contradiction

A *contradiction* is a compound proposition that *false* in all possible cases. An arbitrary contradiction is denoted by

$$\perp$$

### Contingent

A *contingent* proposition is one that is neither a tautology nor a contradiction. Its truth value is contingent on the particular configuration of truth values of its constituent propositions.

## 1.3.2 Logical Equivalences

The compound propositions  $p$  and  $q$  are *logically equivalent* if

$$p \leftrightarrow q \equiv \top \quad (1.1)$$

This statement may be equally expressed by

$$p \equiv q$$

The following are important logical equivalences:

#### Domination Laws

Disjunction with a tautology is a tautology, and conjunction with a contradiction is a contradiction

$$p \vee \top \equiv \top \quad (1.2a)$$

$$p \wedge \perp \equiv \perp \quad (1.2b)$$

#### Identity Laws

Conjunction with a tautology, and disjunction with a contradiction, are *identity* operations on a proposition

$$p \wedge \top \equiv p \quad (1.3a)$$

$$p \vee \perp \equiv p \quad (1.3b)$$

### Negation laws

Conjunction of a proposition with its negation is a contradiction, and disjunction of a proposition with its negation is a tautology

$$p \wedge \neg p \equiv \perp \quad (1.4a)$$

$$p \vee \neg p \equiv \top \quad (1.4b)$$

### Double Negation Law

A proposition is the negation of its negation

$$\neg(\neg p) \equiv p \quad (1.5a)$$

### Idempotent Laws

Both the conjunction and disjunction of a proposition with itself yields itself

$$p \wedge p \equiv p \quad (1.6a)$$

$$p \vee p \equiv p \quad (1.6b)$$

### Absorption Laws

The former two equivalences may be *absorbed* into each other as in the following

$$p \wedge (p \vee p) \equiv p \quad (1.7a)$$

$$p \vee (p \wedge p) \equiv p \quad (1.7b)$$

### Commutative Laws

The commutativity of a binary logical connective can be determined by examining its operation on two propositions with complementary truth values. From Table ??, one observes that

$$\top \wedge \perp \equiv \perp \wedge \top \equiv \perp$$

Likewise, from Table ??

$$\top \vee \perp \equiv \perp \vee \top \equiv \top$$

This leads to the *commutative laws*

$$p \wedge q \equiv q \wedge p \quad (1.8a)$$

$$p \vee q \equiv q \vee p \quad (1.8b)$$

A similar argument may be used to demonstrate the commutativity of  $\oplus$  and  $\leftrightarrow$ .

### Associative Laws

Both conjunction and disjunction are *associative*

$$p \vee (q \vee r) \equiv (p \vee q) \vee r \quad (1.9a)$$

$$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r \quad (1.9b)$$

### Distributive Laws

Conjunction and disjunction *distribute* as in the following

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r) \quad (1.10a)$$

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r) \quad (1.10b)$$

### De Morgan's Laws

De Morgan's laws demonstrate how conjunctions and disjunctions are negated

$$\neg(p \wedge q) \equiv \neg p \vee \neg q \quad (1.11a)$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q \quad (1.11b)$$

### Logical Equivalences with $\rightarrow$

DO TABLE

### Logical Equivalences with $\leftrightarrow$

DO TABLE

### 1.3.3 Propositional Satisfiability

Consider the set of  $N$  propositions

$$\{s_1, s_2, \dots, s_N\} = S_N$$

There exist  $2^N$  possible configurations of truth values for the propositions contained in  $S_N$ . Let  $p$  be a compound proposition that is some amalgamation of the propositions in  $S_N$  through logical connectives. We say that  $p$  is *satisfiable* if there exists a configuration of truth values in  $S_N$  such that  $p$  is true. We call this configuration a *solution* of  $p$ . If there exist no solution to  $p$ , we say the proposition is *unsatisfiable*. That is, a proposition is unsatisfiable if and only if its negation is a tautology.

## 1.4 Predicates and Quantifiers

### 1.4.1 Predicates

A *predicate* is a statement regarding a *subject* that becomes a proposition when the subject is specified. A predicate may be considered as a *propositional function*; that is, a mapping from the subject to a bivalent truth value. We denote a unary predicate as

$$P(x)$$

where  $P$  is the predicate condition, and  $x$  is a variable that denotes the subject. A predicate may have multiple subjects; the predicate

$$P(x_1, x_2, \dots, x_n)$$

is referred to as an *n-ary* predicate.

### 1.4.2 Quantifiers

The variable  $x$  in the statement  $P(x)$  is a *free* variable. In the previous section we discussed how  $P(x)$  becomes a proposition when  $x$  is set to a value. Here, we discuss another way in which  $P(x)$  becomes a proposition: *binding*  $x$  with a *universal quantifier*.

### Universal Quantifier

Consider the set

$$D = \{x_1, x_2, \dots, x_n\}$$

which we shall call the *domain of discourse*. One can construct a compound proposition that is the conjunction of the predicate  $P(x)$  for all  $x$  in  $D$ . That is

$$P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n) \equiv \bigwedge_{i=1}^n P(x_i)$$

Such a prescription is sufficient when the domain of discourse is *finite*. What is the equivalent proposition if  $D$  is *not* finite?

$$\forall x P(x)$$

where we use the **universal quantifier**  $\forall$ , read as “for all”.

### Existential Quantifier

Analagous to the universal quantifier, the proposition

$$P(x_1) \vee P(x_2) \vee \cdots \vee P(x_n) \equiv \bigvee_{i=1}^n P(x_i)$$

may be denoted for a non-finite universe of discourse by

$$\exists x P(x)$$

where we use the **existential quantifier**  $\exists$ , read as “there exists”.

$(s - n)$ -ary predicate. Thus, the act of *nesting* quantifiers is a way in which a predicate can be made into either a proposition or a predicate of lower *arity*. The total number of nested quantifiers may not exceed the arity of the predicate it quantifies.

### 1.4.3 Negating Quantifiers

Using the Associative Laws (??), one may generalise De Morgan’s Laws (??) for  $n$  propositions

$$\neg \left( \bigwedge_{i=1}^n p_i \right) \equiv \bigvee_{i=1}^n \neg p_i \quad (1.12a)$$

$$\neg \left( \bigvee_{i=1}^n p_i \right) \equiv \bigwedge_{i=1}^n \neg p_i \quad (1.12b)$$

Suppose

$$p_i \equiv P(x_i)$$

for some predicate  $P$  where the  $x_i$  are members of a finite domain of discourse  $D$ . What then if  $D$  is *not* finite?

$$\neg \forall x P(x) \equiv \exists x \neg P(x) \quad (1.13a)$$

$$\neg \exists x P(x) \equiv \forall x \neg P(x) \quad (1.13b)$$

Equations ?? are known as *De Morgan’s Laws for Quantifiers*.

### 1.4.4 Nested Quantifiers

Consider the  $n$ -ary predicate

$$P(x_1, x_2, \dots, x_n)$$

As discussed previously, this statement becomes a proposition if and only if all the free variables are either *bound* or *set* to a value. Suppose we bind (or set)  $s < n$  of the free variables. The result is an