ANATOMY 3 ENT MOCK 2

1- The external ear is situated:

A- Below the temporal region
B- In front of the parotid region
C- In front of the mastoid
D- Behind the temporomandibular joint

2- Regarding the muscles of the auricle:

A- Auricularis posterior arises from the mastoid process
B- Auricularis anterior arises from the mastoid process
C- They are divided into extrinsic and intrinsic groups
D- They are rudimentary in humans

3- The external ear is supplied by:

A- The posterior auricular artery
B- The superficial temporal artery
C- Branches of the internal carotid artery
D- Branches of the external carotid artery

4- The lateral wall of the tympanic cavity:

A- Separates it from the external acoustic canal
B- Is entirely bony
C- Is formed in 4/5 by the tympanic membrane
D- Carries the promontory

5- Regarding the roof and floor of the tympanic cavity:

A- The roof is the tegmen tympani, a thin plate of bone
B- The roof separates it from the middle cranial fossa
C- The floor separates it from the jugular bulb
D- The floor separates it from the internal carotid artery

6- Regarding the stapes:

A- It weighs about 2 mg
B- It has a footplate
C- It is the smallest ossicle
D- It articulates with the malleus

7- The tympanic plexus is formed by:

A- The chorda tympani
B- A branch from the geniculate ganglion
C- The tympanic branch of the glossopharyngeal nerve (Jacobson's nerve)
D- Sympathetic fibres from around the internal carotid artery

8- The membranous labyrinth includes:

A- The vestibule
B- The utricle and saccule
C- The semicircular ducts
D- The cochlear duct

9- Near the basal turn of the cochlea:

A- The basal turn bulges into the tympanic cavity as the promontory
B- The scala vestibuli communicates with the vestibule
C- The oval window opens on the scala tympani
D- The scala tympani presents the round window

10- The oval window (fenestra vestibuli):

A- Is closed by the secondary tympanic membrane
B- Is closed by the footplate of the stapes and the annular ligament
C- Is on the lateral wall of the vestibule
D- Connects the vestibule with the middle ear

11- The membranous labyrinth:

A- Is a tube of epithelial origin
B- Has the saccule in its anterior (hearing) part
C- Has an anterior part for hearing, the cochlear duct
D- Has a posterior part for balance: utricle, saccule, semicircular ducts

12- Regarding the hair cells of the cochlea:

A- Fibres of inner hair cells make up 90 to 95% of the cochlear nerve
B- Outer hair cells are contractile and sharpen frequency selectivity
C- Inner hair cells perform mechano-electrical transduction
D- Outer hair cells are the main true sensory receptors

13- The cochlear nerve:

A- Carries static balance from the maculae
B- Carries hearing
C- Runs through the internal auditory canal to the cochlear fossa
D- Its fibres reach the spiral ganglion of Corti

14- Regarding the position of the larynx:

A- It is lower in children than in adults
B- It is higher in children (C2 to C3) than in adults
C- It lies in the middle part of the visceral sheath of the neck
D- In newborns it touches the soft palate, so breathing is nasal until about 4 months

15- The cricoid cartilage:

A- Lies at C6 and completely encircles the lumen
B- Has a narrow anterior arch and a broad posterior lamina
C- Is an incomplete ring open behind
D- Is signet-ring shaped

16- The cricothyroid joint:

A- Joins the arytenoid to the cricoid
B- Allows tilting of the cricoid that tenses the vocal cords
C- Is a flat synovial joint
D- Joins the lower cornu of the thyroid cartilage to the side of the cricoid

17- The cricothyroid muscle:

A- Is supplied by the recurrent laryngeal nerve
B- Is the only muscle on the outer aspect of the larynx
C- Is called the tuning fork of the larynx
D- Is supplied by the external branch of the superior laryngeal nerve

18- The glottis (rima glottidis):

A- Is triangular
B- Is wider in children, so oedema is rarely serious
C- Is narrower than the vestibular cleft
D- Is bounded by the two vocal cords

19- The lymphatics of the larynx:

A- The supraglottic region drains to submental nodes
B- The infraglottic region drains to pretracheal and paratracheal nodes
C- Are very rich in the supraglottic zone
D- Are absent in the vocal cords

20- Check the correct statement(s) about the thyroid gland:

A- It secretes parathyroid hormone
B- It secretes thyroid hormones and calcitonin
C- It is a unique, median endocrine gland of the neck
D- Its relations with the recurrent nerves and parathyroids are the major risks of thyroidectomy

21- The infrahyoid muscles:

A- Are supplied by the anterior rami of the upper three cervical nerves
B- Are enclosed in a fascial sheath
C- Fix or depress the hyoid during swallowing and speech
D- Elevate the hyoid

22- The faces of the lateral lobe:

A- Posterior face: in contact with the sternum
B- Medial face: applied to the trachea and lower larynx
C- Anterolateral face: convex and superficial
D- Posterior face: moulded on the vasculo-nervous bundle of the neck

23- The posterior surface of the isthmus:

A- Is attached to the trachea by the median ligament of Berry
B- Covers the 2nd, 3rd and 4th tracheal rings
C- Lies behind the trachea
D- Covers only the cricoid

24- The parathyroid glands:

A- Are small orange-yellow glands in the fat of the thyroid space
B- Lie on the front of the isthmus
C- The inferior one lies below the end of the inferior thyroid artery
D- The inferior one lies on the posteromedial border of the lateral lobe

25- The thyroid veins:

A- The inferior veins descend to the left brachiocephalic vein
B- The superior vein joins the internal jugular, often via Farabeuf's thyro-linguo-facial trunk
C- The middle veins join the internal jugular directly
D- The inferior veins join the internal jugular

ANSWER KEY

1- A, C, D
2- A, C, D
3- A, B, D
4- A, C
5- A, B, C
6- A, B, C
7- B, C, D
8- B, C, D
9- A, B, D
10- B, C, D
11- A, C, D
12- A, B, C
13- B, C, D
14- B, C, D
15- A, B, D
16- B, C, D
17- B, C, D
18- A, C, D
19- B, C, D
20- B, C, D
21- A, B, C
22- B, C, D
23- A, B
24- A, C, D
25- A, B, C