
Tranche-2 sealed test: 113 group-mean forecast series
next observation
landmark
model 1.1
model 1.0.0 (same training)
analog kNN
pooled-curve ratio
persistence
6.69
8.95
6.92
9.40
10.23
18.25
8.50
15.25
7.98
15.01
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
pooled RMSE, group means (pp of preoperative IOP)


Next observation, group means of public sealed studies
model 1.0.0 (221 targets)
model 1.1 (221 targets)
±5 pp
forecast (% of preoperative IOP)
observed next follow-up (% of preoperative IOP)
20
40
60
80
100
120
140


XEN gel stent: early course followed by the 1.1 transient
(shaded: 90% interval of the 1.1 next-follow-up forecast)
observed (xen-phaco)
model 1.1 next follow-up (xen-phaco)
model 1.0.0 next follow-up (xen-phaco)
observed (xen-alone)
model 1.1 next follow-up (xen-alone)
model 1.0.0 next follow-up (xen-alone)
IOP (% of preoperative)
30
40
50
60
70
10^{-1}
10^{0}
10^{1}
months after surgery (log scale)


Goniotomy (KDB): low day 1, then a rise far above the prior
(shaded: 90% interval of the 1.1 next-follow-up forecast)
observed (kdb-goniotomy)
model 1.1 next follow-up (kdb-goniotomy)
model 1.0.0 next follow-up (kdb-goniotomy)
IOP (% of preoperative)
40
50
60
70
80
90
10^{-1}
10^{0}
months after surgery (log scale)