
Tranche-4 sealed test: 133 group-mean forecast series; H1–H4 pass
H5 fails: initial state 10.94 vs 10.69 pp for 1.2.0
next observation
landmark
model 1.3
model 1.2.0 (same training)
analog kNN
pooled-curve ratio
persistence
6.64
7.92
6.67
8.00
10.67
17.08
8.52
16.03
8.27
15.02
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
pooled RMSE, human group means (pp of preoperative IOP)
Micropulse cyclophotocoagulation (two studies):
the sub-procedure level improves the initial state
IOP (% of preoperative)
45
50
55
60
65
70
10^{-1}
10^{0}
10^{1}
months after surgery (log scale)
observed (8625108:mp-tscpc)
1.3 initial state (8625108:mp-tscpc)
1.2.0 initial state (8625108:mp-tscpc)
observed (10900656:mp-cpc)
1.3 initial state (10900656:mp-cpc)
1.2.0 initial state (10900656:mp-cpc)


Initial state by procedure: where the sub-procedure level helps and hurts
1.2.0
1.3
angle mims
(57 groups, 290 targets)
9.7
10.1
trabeculectomy
(21 groups, 127 targets)
9.6
9.6
tube
(14 groups, 76 targets)
10.2
11.9
cyclodestruction
(16 groups, 55 targets)
14.7
14.1
nonpenetrating
(4 groups, 24 targets)
8.0
8.1
subconjunctival mims
(5 groups, 21 targets)
9.1
8.5
cataract
(5 groups, 19 targets)
20.3
19.9
0
5
10
15
20
initial-state RMSE, human control group means (pp)
Valved tube (one study, two baseline subgroups):
the valved-tube level makes the initial state worse
IOP (% of preoperative)
25
30
35
40
45
50
55
60
10^{0}
10^{1}
months after surgery (log scale)
observed (11722142:su-pap-grade-0)
1.3 initial state (11722142:su-pap-grade-0)
1.2.0 initial state (11722142:su-pap-grade-0)
observed (11722142:su-pap-grade-3)
1.3 initial state (11722142:su-pap-grade-3)
1.2.0 initial state (11722142:su-pap-grade-3)