
Sealed test, group means: 25 studies, 43 forecast series
next observation
landmark
model (robust prior)
analog kNN
pooled-curve ratio
persistence
preoperative level (f = 1)
7.74
9.92
9.77
14.82
8.98
14.24
8.64
12.89
43.53
43.53
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
20.0
pooled RMSE (pp of preoperative IOP; bars clipped at 19)


Next observation, group means of public sealed studies
persistence (103 targets)
model (103 targets)
±5 pp
forecast (% of preoperative IOP)
observed next follow-up (% of preoperative IOP)
20
40
60
80
100
120
140


GATT: early pressure rise after day 1 (not modelled)
(shaded: 90% interval of the next-follow-up forecast)
observed (oag-solo-gatt)
model before any follow-up, U1 (oag-solo-gatt)
model next follow-up, U2 (oag-solo-gatt)
observed (acg-phaco-gatt)
model before any follow-up, U1 (acg-phaco-gatt)
model next follow-up, U2 (acg-phaco-gatt)
IOP (% of preoperative)
40
60
80
100
120
140
10^{-1}
10^{0}
10^{1}
months after surgery (log scale; first point = day 1)


Canaloplasty: very low day 1, then rebound
(shaded: 90% interval of the next-follow-up forecast)
observed (canaloplasty)
model before any follow-up, U1 (canaloplasty)
model next follow-up, U2 (canaloplasty)
observed (phacocanaloplasty)
model before any follow-up, U1 (phacocanaloplasty)
model next follow-up, U2 (phacocanaloplasty)
IOP (% of preoperative)
40
50
60
70
80
90
10^{-1}
10^{0}
10^{1}
months after surgery (log scale; first point = day 1)