
Sealed test: 40 group-mean forecast series
next observation
landmark
model 1.0.0 (r3)
5.75
7.50
r0, no deviations (same training)
6.14
7.37
analog kNN
6.56
9.95
pooled-curve ratio
8.03
10.10
persistence
6.34
9.50
0
2
4
6
8
10
12
pooled RMSE, group means (pp of baseline thickness)


Next observation, group means of public sealed studies
r0 (74 targets)
model 1.0.0 (74 targets)
±5 pp
forecast (% of baseline thickness)
observed next follow-up (% of baseline thickness)
30
40
50
60
70
80
90
100
110
120


Ranibizumab, multicentre cohorts: the model tracks repeated-injection plateaus
(shaded: 90% interval of the next-follow-up forecast)
observed (dme-innovator)
initial-state prediction (dme-innovator)
next follow-up (dme-innovator)
observed (namd-innovator)
initial-state prediction (namd-innovator)
next follow-up (namd-innovator)
observed (brvo-innovator)
initial-state prediction (brvo-innovator)
next follow-up (brvo-innovator)
central thickness (% of baseline)
72.5
70.0
67.5
65.0
62.5
60.0
57.5
55.0
5
10
15
20
25
months after the first treatment


Single dexamethasone implant: the rebound after wear-off is missed
(shaded: 90% interval of the next-follow-up forecast)
observed (dex)
initial-state prediction (dex)
next follow-up (dex)
central thickness (% of baseline)
110
100
90
80
70
60
2.0
2.5
3.0
3.5
4.0
4.5
5.0
5.5
6.0
months after the first treatment