

УДК 51, 53
ББК 22.19

Сборник научных трудов Международной конференции «Разностные схемы и их приложения», посвященной 90-летию профессора В.С.Рябенного
Институт прикладной математики им. М.В. Келдыша РАН,
Москва, 27-31 мая 2013 г.

В сборнике представлены доклады по вычислительной и прикладной математике. В них отражено современное состояние теории и приложений численных методов для решения уравнений математической физики, обсуждается проблематика теории разностных потенциалов и искусственных граничных условий, рассматриваются вопросы описания и расчетов турбулентных течений.

Proceedings of the International Conference «Difference schemes and applications» in Honor of the 90-th Birthday of Prof. V.S. Ryaben'kii
Keldysh Institute of Applied Mathematics, May 27-31, 2013

The collection contains papers on computational and applied mathematics. They reflect the present state of the theory and application of numerical methods for solving the equations of mathematical physics, the problems of the theory of difference potentials and artificial boundary conditions are discussed, progress in calculation of turbulent flows are considered.

Программный комитет искренне благодарит за поддержку Российскую Академию Наук, Международную Ассоциацию по Математике и Компьютерам в Моделировании, Государственный Университет Штата Северная Каролина, США, Бюро Научных Исследований Армии США и Европейское Бюро Аэрокосмических Исследований и Разработок (EOARD), Лабораторию FlowModellium "Математическое моделирование нелинейных процессов в газовых средах" МФТИ.



Редакционная коллегия: К.В. Брушлинский, М.С. Гавреева, В.Т. Жуков,
А.В. Северин, Н.А. Чмыхова

Федеральное государственное бюджетное учреждение науки
Институт прикладной математики
имени М.В. Келдыша Российской академии наук

Москва, 2013

ISBN: 978-5-98354-010-1

© ИПМ им. М.В. Келдыша РАН

ОГЛАВЛЕНИЕ

	Стр.
В.С. Рябенский КЛЮЧЕВАЯ ИНФОРМАЦИЯ ДЛЯ УПРАВЛЕНИЯ РЕШЕНИЯМИ ЛИНЕЙНЫХ РАЗНОСТНЫХ СХЕМ В СОСТАВНЫХ ОБЛАСТЯХ ...	11
S. Abarbanel, A. Ditkowski WAVE PROPAGATION IN ADVECTED ACOUSTICS WITHIN A NON- UNIFORM MEDIUM UNDER THE EFFECT OF GRAVITY	14
G.I. Barenblatt, A.G. Chorin, V.M. Prostokishin THE SIMPLEST TURBULENT FLOW: SHEAR FLOW AT VERY LARGE REYNOLDS NUMBERS. A SURVEY OF THE TOTAL OF STUDIES	15
M.A. Botchev EFFICIENT TIME-STEPPING-FREE TIME INTEGRATION OF THE MAXWELL EQUATIONS	16
A. Chertock A SIMPLE EULERIAN FINITE-VOLUME METHOD FOR COMPRESSIBLE FLUIDS IN DOMAINS WITH MOVING BOUNDARIES	17
T. Hagstrom BOUNDARY CONDITIONS FOR HIGH-RESOLUTION SIMULATIONS OF WAVES	18
D.S. Kamenetskiy, J.E. Bussioletti, C.L. Hilmes F.T. Johnson, V. Venkatakrishnan, L.B. Wigton NUMERICAL EVIDENCE OF MULTIPLE SOLUTIONS FOR THE REYNOLDS--AVERAGED NAVIER--STOKES EQUATIONS FOR HIGH--LIFT CONFIGURATIONS	19
A.V. Karpeev, I.V. Glazyrin, O.G Kotova IONIZATION-ASSISTED RELATIVISTIC ELECTRON GENERATION WITH MONOENERGETIC FEATURES FROM LASER THIN FOIL INTERACTION	22
A. Kurganov CENTRAL SCHEMES: A POWERFUL BLACK-BOX SOLVER FOR NONLINEAR HYPERBOLIC PDES	23

The results were compared with all published experimental data and an instructive confirmation was found. An analogous investigation with the same conclusions was performed for boundary layer flows and wall-jets.

The authors came to the conclusion that the von Karman-Prandtl law is not correct. It should be excluded from the teaching process and replaced by proposed and experimentally approved power law.

EFFICIENT TIME-STEPPING-FREE TIME INTEGRATION OF THE MAXWELL EQUATIONS

M.A. Botchev¹

*¹Department of Applied Mathematics, University of Twente,
Enschede, the Netherlands,
m.a.botchev@utwente.nl*

Solution of the time dependent Maxwell equations is an important problem arising in many applications ranging from nanophotonics to geoscience and astronomy. The problem is far from trivial, and solutions typically exhibit complicated wave properties as well as damping behavior. Usually, special staggered time stepping schemes are used [1]. Although their time step may be severely restricted by the CFL condition, performance of these schemes is hard to beat by modern implicit or exponential time integration schemes [2]. We show that in some cases so-called time-stepping-free schemes provide a very efficient alternative to the standard schemes. These schemes employ the matrix exponential function and can be implemented by special block Krylov subspace techniques [3,4].

Numerical examples demonstrating the efficiency of the proposed approach are presented, coming from the fields of nanophotonics and geoscience.

References:

1. M.A. Botchev, J.G.Verwer, Numerical integration of damped Maxwell equations // SIAM J. Sci. Comput. 2009, vol. 31(2), pp. 1322-1346 .
2. J.G.Verwer,, M.A. Botchev, Unconditionally stable integration of Maxwell's equations // Linear Algebra and its Applications, 2009, vol. 431, Issues 3-4, pp. 300-317.
3. M.A. Botchev, V. Grimm, M. Hochbruck, Residual, restarting and Richardson iteration for the matrix exponential. // To appear in SIAM J. Sci. Comput., accepted in February 2013.
4. M.A. Botchev, A block Krylov subspace time-exact solution method for linear ODE systems. To appear in Numer. Lin. Algebra Appl., accepted in December 2012.