

Text: 4414 words

Preserved neural specialization for non-social information in autism

James C. McPartland, Ph.D.¹, Jia Wu, Ph.D.¹, Christopher A. Bailey¹,

Robert T. Schultz, Ph.D.^{2,3} & Ami Klin, Ph.D.¹

¹Yale Child Study Center, New Haven, CT

²Department of Pediatrics, University of Pennsylvania, Philadelphia, PA

³Center for Autism Research, Children's Hospital of Philadelphia, PA

This research was supported by funding from NIH (KL2RR024138) and NIMH (R03MH079908) to James McPartland and NICHD (PO1HD003008) to Fred Volkmar.

Correspondence and reprint requests to:

James McPartland

Yale Child Study Center

230 South Frontage Road

New Haven, CT 06520

james.mcpartland@yale.edu

Phone: (203) 785-7179

Fax: (203) 764-4373

Abstract

Context: Aberrant neural response to human faces in autism has been attributed to atypical social development and consequently reduced developmental exposure. The specificity of this deficit in neural specialization remains unclear.

Objective: To contrast neural specialization for social information (human faces) and non-social information (letters) in individuals with autism and typical development.

Design: Event-related potentials elicited by faces, inverted faces, houses, letters, and pseudoletters were recorded, and an electrophysiological marker of neural specialization (N170) was extracted.

Setting: University medical setting.

Participants: 36 individuals with autism spectrum disorder and 18 typically developing individuals matched for age, race, sex, handedness, and cognitive ability.

Main Outcome Measures: N170 amplitude and latency, behavioral performance on standardized measures of face recognition and word reading/decoding.

Results: Individuals with autism displayed slowed face processing and decreased sensitivity to face inversion, despite comparable brain response to letters. Brain responses were associated with behavioral performance in both groups.

Conclusions: Individuals with autism display atypical neural specialization for social information but intact specialization for non-social information. Results concord with specific dysfunction in social brain systems rather than non-specific problems with information processing.

Preserved neural specialization for non-social information in autism

The ability to efficiently perceive the human face is a crucial and early-emerging social ability. Specialized processing for faces emerges in the first days of life¹⁻⁴ and is honed by developmental experience⁵. Faces come to be encoded using configural processing mechanisms⁶, reflected in disproportionate impairments in recognizing both upside down faces (the inversion effect⁷) and facial features out of context⁸. Functional neuroimaging studies show that faces elicit selective, right-lateralized hemodynamic activity in a portion of occipitotemporal cortex, the fusiform gyrus⁹⁻¹¹, and intra-cranial electrophysiological recordings reveal face-related negative electrical activity originating from this portion of cortex¹². Likewise, event-related potentials (ERPs) recorded from corresponding scalp regions show a negative-going electrical deflection approximately 170 milliseconds after viewing a face (N170)¹³. The N170 reflects structural encoding, an early stage of face processing preceding higher-order processes like recognition¹⁴, and is sensitive to perturbations in face configuration, including inversion¹⁵. Neural generators of the N170 have been localized to occipitotemporal sites including the fusiform gyrus¹⁶⁻¹⁸, as well as the superior temporal sulcus¹⁹, lingual gyrus¹⁸, and posterior inferotemporal gyrus^{18,20}.

These processing strategies and brain regions are also observed in the processing of visual stimuli with which viewers have extensive experience, or *perceptual expertise*^{21,22}. Experts at perceiving and discriminating among exemplars within a visually homogenous class (e.g., Greebles, birds, or cars^{23,24}) develop face-like patterns of brain activity, both in terms of hemodynamic²⁵ and electrophysiological response, as indexed by the N170²⁶. According to this model, these brain regions subserve a processing style rather than specific content, and face-related brain activity reflects human beings' extensive experience processing human faces during development²⁷.

Analogous specialization through developmental experience occurs in brain mechanisms subserving letter and word processing. Perception of printed letters²⁸ and words²⁹ selectively activates left fusiform gyrus and elicits a left-lateralized N170 in literate children as young as eight years of age^{30,31}. A maturational course

independent of higher-order phonological or semantic processes^{32,33} and an early time course suggest that the “letter N170” marks pre-linguistic processes related to visual perception of form³¹ and, like the N170 elicited by faces, automatic perceptual categorization within a domain of expertise³⁴. Perceptual expertise effects for letters are revealed by enhanced N170 amplitude to familiar alphabets but not foreign alphabets or nonsensical letter approximations (pseudoletters)³⁵. Converging evidence from neuroimaging studies and source localization analyses suggest left-lateralized sources in the fusiform gyrus and the inferior occipitotemporal cortex^{16,34,36}. Though letter-related brain activity is typically contralateral to face processing areas¹⁶, there is some degree of functional overlap; under special circumstances, such as precocious reading ability, right fusiform gyrus is recruited for letter and word recognition³⁷.

Because face perception is a well-understood and socially important behavior, it has been employed as an avenue to understand social development in autism spectrum disorder (ASD). In ASD, decreased attention to human faces is evident by 6 to 12 months^{38,39}, and abnormalities in face perception and recognition have been observed throughout the lifespan⁴⁰⁻⁴⁴. Individuals with ASD often exhibit abnormal viewing patterns to faces^{45,46}, and neuroimaging studies reveal hypoactivation of the fusiform gyrus during face viewing^{42,47}. Studies of electrophysiological correlates of face perception suggest delayed N170 to human faces and decreased sensitivity to face inversion in individuals with ASD, as well as first degree relatives⁴⁸⁻⁵⁴.

One theoretical explanation for these observed differences in face perception in ASD focuses on the role of developmental exposure to faces. The social motivation hypothesis⁵⁵ posits that, due to abnormalities in social drive very early in childhood, children with ASD do not attend to faces during sensitive developmental periods. Consequently, people with ASD fail to develop face processing expertise and associated patterns of behavioral and brain specialization⁵⁶. Because the social motivation hypothesis implicates social drive as the dysfunction from which face perception difficulties originate (and not atypical functioning of brain regions subserving perceptual expertise), it presumes that individuals with ASD, given appropriate exposure to and interest in a stimulus class, should develop perceptual expertise in terms of both behavioral and brain specialization⁵⁷. This notion is supported by a single-case study revealing behavioral and neural indices of

perceptual expertise in a child with ASD during perception of cartoon characters associated with a circumscribed interest⁵⁸. Though others have attempted to investigate brain response associated with expertise in this population⁵⁹, research has been stymied by difficulty finding shared areas of expertise in ASD; whereas groups of study participants experienced in perceiving faces are common, groups of individuals with ASD who share a common non-face area of expertise are rare.

The current work circumvented this difficulty by examining brain activity reflecting perceptual expertise for letters of the alphabet. As described above, development of perceptual expertise for letters and words is well studied and elicits brain activity similar to faces in terms of temporal characteristics and scalp topography (despite lateralization differences). This is a novel and uniquely appropriate comparison because, despite developmental disinterest towards faces and characteristic weakness in language, facility with reading has been a noted strength in ASD since Kanner's original account⁶⁰. High-functioning individuals on the autism spectrum display age-appropriate skills in single word-reading and word-decoding ability⁶¹⁻⁶³, and a subgroup possesses precocious interest and proficiency in reading, or *hyperlexia*⁶⁴⁻⁶⁶. In this study, electrophysiological and behavioral methods were applied to compare perceptual expertise for faces and letters in individuals with ASD. Experiments contrasted neural response to faces versus houses, faces versus inverted faces, and letters versus pseudoletters and compared these parameters to behavioral measures assessing proficiency in face recognition and letter and word perception. Consistent with previous work, it was hypothesized that individuals with ASD would exhibit impaired face recognition and delayed brain response to faces, as well as decreased sensitivity to face inversion. In keeping with the notion that these atypicalities reflect developmental sequelae of social deficits, it was predicted that similar anomalies would not be observed for non-social stimuli; individuals with ASD would show typical skills in terms of letter and word perception and comparably enhanced response to letter stimuli with respect to unfamiliar pseudoletters. As prior work has revealed relationships among neural correlates of face perception and behavioral measures of face recognition⁴⁹, exploratory analyses examined relationships among neural and behavioral measures of face and letter perception.

Methods

Participants

Two groups participated in the study: individuals with ASD and medically and neuropsychiatrically healthy individuals with typical development. Exclusionary criteria for participants with ASD included seizures, neurological disease, history of serious head injury, sensory or motor impairment that would impede completion of the study protocol, active psychiatric disorder (other than ASD; screened with the Child Symptom Inventory: Fourth Edition ⁶⁷), or medication known to affect brain electrophysiology. Additional exclusionary criteria for typical participants included the above plus learning/language disability or family history of ASD. From an existing pool of subjects involved in ongoing research at the Yale Child Study Center, participants were selected based on having a Full Scale IQ (Differential Ability Scales: Second Edition ⁶⁸; Wechsler Intelligence Scale for Children – Fourth Edition ⁶⁹; Wechsler Adult Intelligence Scale: Third Edition ⁷⁰) in the average range or higher (Standard Score of 80 or above). All individuals with ASD had a pre-existing diagnosis that was confirmed with gold standard diagnostic assessments for research: combination of parent interview (Autism Diagnostic Interview-Revised ⁷¹; ADI-R), semi-structured social and communication assessment (Autism Diagnostic Observation Schedule ⁷²), and clinical diagnosis based on DSM-IV-TR ⁷³ criteria by an expert clinician. The ADI was not administered to one subject because a parent was unavailable for interviewing. Two individuals were included in the sample who failed to meet ADI-R onset criteria; for both of these high-functioning, verbal individuals, problems were not detected until enrolled in school with peers. In addition to the aforementioned exclusionary criteria, typical participants were recruited to match the ASD sample in terms of sex, ethnicity (determined by self or parent report), handedness (Edinburgh Handedness Inventory ⁷⁴), chronological age, and Full Scale IQ (Wechsler Abbreviated Scale of Intelligence ⁷⁵), and groups did not significantly differ on any of these variables. Behavioral assessments could not be administered to one typical participant due to time limitations. All procedures were approved by the Human Investigation Committee at Yale School of Medicine and were carried out in accordance with the Declaration of Helsinki (1975/1983). Of an initial sample of 57 individuals with ASD and 25 typically developing participants, adequate artifact-free

data was obtained from 36 and 18 participants, respectively, in Block 2, and 32 and 17, respectively, in Block 1. Table 1 displays demographic data for the larger sample; variation in sample between blocks did not introduce significant differences on matching factors.

[PLEASE INSERT TABLE 1 ABOUT HERE]

EEG procedures

Stimuli. Stimuli were administered in pseudorandom sequence in two counterbalanced blocks. The first block consisted of gray-scale digitized images of neutral faces, houses, inverted faces, and inverted houses (not included in current analyses), all displayed from a direct frontal perspective. The second block included letters and a confabulated alphabet of pseudoletters³⁵. Example stimuli are displayed in the legends of Figures 1 and 2. Subjects were presented with 23 stimuli from each category four times, for a total of 92 stimuli per category. Stimuli were standardized in terms of size (subtended approximately five degrees of visual angle), background color (gray), and average luminance. To maximally engage participants with stimuli, participants completed a one-back task during administration, pressing a button when a stimulus repeated (9 times for each stimulus category). Because this behavioral task was confounded with face recognition, attention to task was monitored in real time through closed-circuit video, enabling pausing of data collection and redirection of attention to stimulus presentation if needed.

Data collection. Stimuli were presented on a Pentium-IV computer controlling a 51 cm color monitor (75-Hz, 1024x768 resolution) running E-Prime 2.0 software⁷⁶. Displays were viewed at a distance of 90 cm in a sound attenuated room with low ambient illumination. EEG was recorded using NetStation 4.3. A 256 lead Geodesic sensor net (Electrical Geodesics Incorporated;⁷⁷ was dampened with potassium-chloride electrolyte solution, placed on the participant's head, and fitted according to the manufacturer's specifications. Impedances were kept below 40 kilo-ohms. ERP was recorded continuously throughout each stimulus presentation trial, consisting of a fixation cross (randomly varying from 250-750 ms), stimulus (500 ms), and blank screen (500

ms). The EEG signal was amplified (x1000) and filtered (0.1 Hz high-pass filter and 100 Hz elliptical low-pass filter) via a preamplifier system (Electrical Geodesics Incorporated). The conditioned signal was multiplexed and digitized at 250 Hz using an analog-to-digital converter (National Instruments PCI-1200) and a dedicated Macintosh computer. The vertex electrode was used as a reference, and data were re-referenced to an average reference after data collection.

Data editing and reduction. Data were averaged for each subject by stimulus type across trials. Averaged data were digitally filtered with a 30 Hz low-pass filter and transformed to correct for baseline shifts. The window for segmentation of the ERP was set from 100 ms before and 500 ms after stimulus onset. NetStation artifact detection settings were set to 200 μ v for bad channels, 150 μ v for eye blinks, and 150 μ v for eye movements. Channels with artifacts on more than 50 percent of trials were marked as bad channels and replaced through spline interpolation. Segments that contained eye blinks, eye movement, and those with more than 20 bad channels were also excluded. Participants with less than 46 good trials for any stimulus category were excluded from analysis. Electrodes of interest were selected based on maximal observed amplitude of the N170 to faces and letters in grand averaged data and to conform to those used in previous research. Data were averaged across eight electrodes over the left (95, 96, 97, 106, 107, 108, 116, 117) and right lateral posterior scalp (151, 152, 153, 160, 161, 162, 170, 171). The time window for N170 analysis, extending from 108 ms to 327 ms post-stimulus onset, was chosen by visual inspection of grand averaged data and then customized for each subject to confirm that the component of interest was captured at each electrode. Peak amplitude and latency to peak were averaged across each electrode group within the specified time window and were extracted for each participant for each stimulus category.

Data analysis. N170 amplitudes and latencies to peak were separately analyzed using univariate repeated measures analyses of variance (ANOVA) with two within-subject factors, each with two levels: Condition (face/house; face/inverted face, letter/pseudoletter) and hemisphere (left/right). The between subjects factor was Group (ASD/Typical).

Behavioral procedures

Face perception. Face recognition was measured with the Benton Facial Recognition Test ⁷⁸.

Participants viewed a grayscale image of a face and specified one or three matches from an array of six faces, varying in shadowing and orientation.

Letter perception. The Letter-Word Identification and Word Attack subtests of the Woodcock-Johnson Tests of Achievement – Third Edition ⁷⁹ required the participant to read words aloud, with the former using genuine English words and the latter using novel words. Both subtests yielded a standard score (Mean = 100, SD = 15) derived from an age-based standardization sample.

Data analysis. Between-group differences in behavioral measures were analyzed with independent samples t-tests. Interrelationships among behavioral measures and ERP parameters (N170 latency, amplitude) were computed using Pearson Product-Moment Correlations.

Results

Electrophysiological measures

Faces versus houses. Tables 2 and 3 display N170 latency and amplitude, respectively, in all conditions for both groups and in both hemispheres. Figure 1 displays waveforms depicting ERPs to faces and houses. Faces elicited N170s with shorter latencies (main effect of Condition; $F(1,47) = 30.10, p \leq .01$) and larger amplitudes (main effect of Condition; $F(1,47) = 49.77, p \leq .01$) across hemispheres for both groups. Right-lateralization was evident only in typically developing individuals (Hemisphere by Group interaction; $F(1,47) = 7.22, p \leq .01$). N170 amplitude to houses was reduced in the left hemisphere across groups (Hemisphere by Condition interaction; $F(1,47) = 6.80, p \leq .01$), and, relative to typical individuals, bilaterally in the ASD group (Condition by Group interaction; $F(1,47) = 25.02, p \leq .05$). Predicted differences in latency between groups were reflected in a three-way interaction (Group by Hemisphere by Condition interaction; $F(1,47) = 3.09, p \leq .10$); in the right hemisphere, N170 latency to faces was significantly faster (a differences of approximately 20.4 milliseconds) in typically developing individuals than those with ASD ($F(1, 47) = 5.57; p \leq .05$). Figure 2 displays N170 amplitudes for faces and houses, highlighting this difference.

[PLEASE INSERT TABLE 2 ABOUT HERE]

[PLEASE INSERT TABLE 3 ABOUT HERE]

[PLEASE INSERT FIGURE 1 ABOUT HERE]

[PLEASE INSERT FIGURE 2 ABOUT HERE]

Face versus inverted faces. Waveforms depicting ERPs to faces and inverted faces are displayed in Figure 1. Inverted faces elicited N170s with longer latencies than upright faces (main effect of Condition; $F(1,47) = 4.66, p \leq .05$) across hemispheres for both groups. Across faces and inverted faces, typically developing individuals displayed enhanced amplitude in the right hemisphere, while those with ASD exhibited equivalent amplitude in both hemispheres (Hemisphere by Group interaction; $F(1,47) = 7.70, p \leq .01$). Across

hemisphere, typically developing individuals displayed an inversion effect in the expected direction, with larger amplitude to inverted relative to upright faces, whereas individuals with ASD displayed attenuated N170 amplitudes to inverted faces relative to upright faces (Condition by Group interaction; $F(1,47) = 5.84, p \leq .05$).

Letters versus pseudoletters. Figure 3 displays waveforms depicting ERPs to letters and pseudoletters. For both groups, letters elicited N170s with larger amplitudes than pseudoletters across hemispheres (main effect of Condition; $F(1, 52) = 14.67, p \leq .01$). As displayed in Figure 4, paired samples t-tests revealed that this effect was carried by significantly enhanced amplitude to letters versus pseudoletters in the typical group in the left hemisphere ($t(1,17) = 2.12, p \leq .05$) and in the ASD group in both left ($t(1,35) = 2.90, p \leq .01$) and right hemispheres ($t(1,35) = 3.34, p \leq .01$). No other significant between group differences were detected.

[PLEASE INSERT FIGURE 3 ABOUT HERE]

[PLEASE INSERT FIGURE 4 ABOUT HERE]

Behavioral measures

Face perception. Table 4 displays mean score and standard deviation on behavioral measures for both groups. Individuals with ASD obtained significantly lower face recognition scores than typically developing individuals ($t(1,51)=3.29, p \leq .01$). For both groups, N170 latency to faces in the right hemisphere was correlated with face recognition skill; individuals with faster N170s displayed better face recognition performance (ASD: $r = -.39, p \leq .05$; Typical: $r = -.53, p \leq .05$). Among individuals with ASD, N170 amplitude to inverted faces was correlated with face recognition performance; those with better face recognition abilities were more likely to display an enhanced N170 associated with inversion ($r = -.47, p \leq .01$).

Letter perception. Groups performed comparably and in the average range on word reading and decoding tasks. Among typically developing individuals, longer N170 latency to letters in the right hemisphere was correlated with word reading score; those with longer latencies tended to perform better on the measure of single word reading ($r = .64, p \leq .01$).

270
271
272
273
274
275

[PLEASE INSERT TABLE 4 ABOUT HERE]

Comment

The current study contrasted neural specialization for social and non-social information in individuals with ASD and a cohort of typically developing individuals of comparable age, ethnicity, sex, handedness, and cognitive ability. A critical social stimulus with which most adults possess great experience, the human face, was contrasted with a comparably complex visual stimulus with minimal social relevance, houses. Consistent with predictions and with prior research⁴⁹, individuals with ASD displayed a selective processing delay for human faces relative to typical counterparts. This was reflected in a substantial (~20 milliseconds, approximately 10% of total response time) difference in peak latency of the N170 ERP component between groups that was specific to faces and evident in the right hemisphere, the hemisphere in which face processing typically takes place. Individuals with ASD showed reduced hemispheric specialization compared to typical counterparts, who showed a marked right lateralization effect for faces. A second analysis contrasted brain response to upright and inverted human faces. This manipulation disrupts configural processing strategies typically applied to human faces and other expert stimuli and, in typically developing samples, perturbs neural response, resulting in an enhanced N170. In the current experiment, this inversion effect was evident in typically developing individuals; however, those with ASD showed attenuated, rather than enhanced, brain response to inverted faces. Finally, on a behavioral measure of face recognition, individuals with ASD, despite comparable intellectual ability, performed significantly worse than typically developing counterparts. Face recognition performance was associated with processing efficiency for faces; in both groups, individuals with better face recognition abilities displayed faster N170 response. Among individuals with ASD, increased inversion effects, as reflected by a stronger response to inverted faces, was associated with better face recognition performance. This pattern of anomalies, i.e., decreased efficiency of processing, insensitivity to inversion, and impaired face recognition, is hypothesized to reflect underdeveloped expertise for faces, a downstream effect of decreased attention to faces during childhood secondary to reduced social drive from infancy⁵⁵. Indeed, the observed correlation between neural response to face inversion and recognition performance suggests that, in this case, development of expertise and processing proficiency are related.

These findings concord with prior work describing deviant social development in ASD. However, scant evidence to-date informs the specificity of the observed neural processing anomalies to social information. By measuring responses to non-social expert stimuli, the current study sought to demonstrate such selectivity in perceptual deficits in ASD. Following up on work showing N170-related expertise effects for letters, ERP response to letters of Roman alphabet were compared to a confabulated alphabet of pseudoletters³⁵. In contrast to the discrepancies observed during perception of social stimuli, individuals with ASD displayed neural responses comparable to typical counterparts; both groups showed enhanced N170 to familiar letters. Similar results were obtained on behavioral measures, with individuals with ASD obtaining word reading and decoding scores that were comparable to the typical participants in this study and within the average range. These results suggest intact specialization, both behavioral and neural, associated with perceptual expertise for letters in individuals with ASD.

The current findings have significant implications for understanding the neuropathology of autism spectrum disorders. Two prevailing classes of theories attribute autistic impairments to (a) dysfunctional brain structures supporting social information processing⁵⁵ or (b) altered connectivity among distributed brain regions⁸⁰; the former bespeaks the import of content and the latter bespeaks the import of process. Social information processing theories posit that social information is qualitatively unique and that specific brain systems have evolved to support this type of information processing. Connectivity theories, in contrast, have traditionally argued that social information is relevant only insofar as it relies on complex or cortically distributed processing mechanisms. The current work demonstrates, for the first time in a substantial sample of children with ASD, preserved specialization for a cognitive process subserved by distributed cortical regions. Development of letter expertise is an intricate process accruing over time and requiring elaborate communication of anterior and posterior cortical regions⁸¹. The demonstration of capacity for development of perceptual expertise in ASD reveals an example of intact functioning of such brain systems that is not consistent with non-specific, brain-wide dysfunction. Taking into consideration considerable evidence for atypical patterns of connectivity in ASD⁸⁰, current findings emphasize the potential value of studying connectivity *within*

specific brain systems in a developmental context. In this way, scientists may also extricate atypical connectivity as potential cause or consequence of autistic dysfunction; it is likely that origins of dysfunction in functionally specific brain systems would, through developmental maturation, lead to broader connectivity problems. Such research may also serve to clarify to what degree problems with connectivity uniquely differentiate autism from the diversity of developmental and psychiatric disorders also manifesting atypical connectivity, e.g. obsessive-compulsive disorder,⁸² schizophrenia⁸³, ADHD⁸⁴, and intellectual impairment⁸⁵.

This work yields clinically relevant implications for the detection and treatment of ASD. Results are supportive of the broad class of interventions designed to direct the attention of children with ASD to relevant social information. When children are appropriately engaged and attuned to information, in this case, letters, typical patterns of neural specialization develop; given the right input, the brain of a person with autism can function like that of a typical peer, without ostensible reliance on compensatory mechanisms or alternative processing strategies. Findings add to a body of evidence that electrophysiological brain activity to faces represents a viable bio-behavioral risk marker for ASD, as temporal anomalies in neural correlates of face perception have been observed in children with ASD⁵³ and infants at-risk for ASD⁵⁴.

Though the current work replicates initial findings of temporal anomalies to faces⁴⁹, these findings have not fully replicated in all samples⁸⁶⁻⁸⁸. These varied results most likely reflect the phenotypic heterogeneity evident in ASD. Despite the unifying characteristic of social impairment, ASD presents in a remarkable diversity of manifestations, likely representing multiple etiologic pathways and variability in developmental experience⁸⁹. Considering the manner in which face processing (especially in older children and adults) has been actively shaped by experience, it is intuitive that anomalies might emerge in a variety of ways or might not emerge universally⁹⁰. In this regard, like any of the symptoms characterizing autism, anomalous face perception is neither necessary nor specific. It is one potential manifestation of atypical social development that, by virtue of a deep understanding of behavioral and brain bases in typical social development, is a viable avenue for investigating social disability. Variability in electrophysiological studies of face perception may also relate to differences in visual attention⁸⁶, a trend observed in hemodynamic studies^{91,92}. Our employment of a

pre-stimulus fixation crosshair reduces the likelihood that between-group differences are attributable to differences in visual attention; resolution of this matter will ultimately require co-registration of eye-tracking and EEG. Finally, as is the case in all clinical research, variability in results is likely to mirror variability in clinical characterization. Application of gold standard diagnostic criteria has been inconsistent in ERP research; moving forward, adoption of clinically rigorous methods of characterization will be vital to derive directly comparable results.

There are several aspects of the current work that are being revisited and improved upon in ongoing research. Limiting the sample to high-functioning individuals was a necessary first step towards addressing the research questions posed in this study, but it limits generalizability to the broader range of individuals with ASD. Given that even many nonverbal children with ASD are capable of reading, these types of experiments offer a window into domains of strength and preserved neural functions of children on the autism spectrum, important goals for tailoring interventions and proscribing specific treatments. The sample in the current study focused on pre-adolescence, a time of rapid maturation of brain systems subserving face perception. Additional research in younger and older children and adults will elucidate the protracted maturational course of specialization for face perception in ASD and of letter expertise in both typical and atypical development. Of note, many participants in the current study displayed the bifid waveform morphology characteristic of pre-adult face responses⁹³; however, this was not evident for letter N170s. Exploiting the dense spatial sampling afforded by the 256 lead net, analyses in progress are using individual-specific three-dimensional head models (computed with sensor registration images acquired with the Geodesic Photogrammetry System) to localize potentially distinct neural sources for these facets of the developing N170.

Understanding developmental factors is particularly important in the current context in that perceptual expertise for letters is clearly a distinct phenomenon from face expertise, occurring over a relatively compressed period of time rather than from birth. It will thus be essential to examine perceptual expertise for a greater variety of stimuli. Though it has been proposed that, like faces, letters are encoded using a holistic processing strategy⁹⁴, unlike faces, letters are processed at a basic rather than subordinate level of identification²⁸. The

N170 has been posited to denote perceptual expertise at this basic level of identification, while later components, such as the N250, index expertise at the subordinate level of identification⁹⁵. Similar mechanisms underlying perceptual expertise for both faces and letters exist at early processing stages as indexed by the N170, but study of a broader range of electrophysiological components and expert stimuli will paint a clearer picture of perceptual expertise development in ASD.

Acknowledgements

This research was supported by funding from NIH (KL2RR024138) and NIMH (R03MH079908) to James McPartland and NICHD (PO1HD003008) to Fred Volkmar. Sponsors were not involved in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript. The authors report no potential conflicts of interest. James McPartland had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. The authors gratefully acknowledge the contributions of Linda Mayes, Michael Crowley, Sarah Shultz, Joshua Diehl, Stephanie Huckins, Rebecca Pearlson, and Benjamin Aaronson. Portions of this work were presented at the International Meeting for Autism Research in London, England in May 2008 and in Chicago, Illinois in May 2009 and at the American Psychological Association Annual Convention in Toronto, Canada in August 2009.

References

1. Johnson MH, Dziurawiec S, Ellis H, Morton J. Newborns' preferential tracking of face-like stimuli and its subsequent decline. *Cognition*. Aug 1991;40(1-2):1-19.
2. Goren CC, Sarty M, Wu PY. Visual following and pattern discrimination of face-like stimuli by newborn infants. *Pediatrics*. Oct 1975;56(4):544-549.
3. Bushnell I, Sai F, Mullin J. Neonatal recognition of the mother's face. *British Journal of Developmental Psychology*. 1989;7:3-15.
4. Meltzoff AN, Moore MK. Imitation of facial and manual gestures by human neonates. *Science*. Oct 7 1977;198(4312):74-78.
5. Nelson C. The development and neural bases of face recognition. *Infant and Child Development*. 2001;10:3-18.
6. Farah MJ, Tanaka JW, Drain HM. What causes the face inversion effect? *J Exp Psychol Hum Percept Perform*. Jun 1995;21(3):628-634.
7. Yin R. Face recognition by brain-injured patients: A dissociable ability. *Neuropsychologia*. 1970;8:395-402.
8. Tanaka JW, Farah MJ. Parts and wholes in face recognition. *Q J Exp Psychol A*. May 1993;46(2):225-245.
9. Puce A, Allison T, Gore J, McCarthy G. Face-sensitive regions in human extrastriate cortex studied by functional MRI. *Journal of Neurophysiology* 1995.
10. Kanwisher N, McDermott J, Chun MM. The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J Neurosci*. Jun 1 1997;17(11):4302-4311.
11. Haxby JV, Grady CL, Horwitz B, Ungerleider JM, Maisog M, Pietrini P. The functional organization of human extrastriate cortex: A pet-rCBF study of selective attention to faces and locations. *The Journal of Neuroscience*. 1994;14:6336-6353.

12. Allison T, McCarthy G, Novbre A, Puce A, Belger A. Human extrastriate visual cortex and the perception of faces, words, numbers, and colors. *Cerebral Cortex*. 1994;5:544-554.
13. Bentin S, Allison T, Puce A, Perez E, et al. Electrophysiological studies of face perception in humans. *Journal of Cognitive Neuroscience*. Nov 1996;8(6):551-565.
14. Bentin S, Deouell LY, Soroker N. Selective visual streaming in face recognition: evidence from developmental prosopagnosia. *Neuroreport*. Mar 17 1999;10(4):823-827.
15. Rossion B, Gauthier I, Tarr MJ, et al. The N170 occipito-temporal component is delayed and enhanced to inverted faces but not to inverted objects: an electrophysiological account of face-specific processes in the human brain. *Neuroreport*. Jan 17 2000;11(1):69-74.
16. Rossion B, Joyce CA, Cottrell GW, Tarr MJ. Early lateralization and orientation tuning for face, word, and object processing in the visual cortex. *Neuroimage*. Nov 2003;20(3):1609-1624.
17. Itier RJ, Taylor MJ. Inversion and contrast polarity reversal affect both encoding and recognition processes of unfamiliar faces: a repetition study using ERPs. *Neuroimage*. Feb 2002;15(2):353-372.
18. Shibata T, Nishijo H, Tamura R, et al. Generators of visual evoked potentials for faces and eyes in the human brain as determined by dipole localization. *Brain Topogr*. Fall 2002;15(1):51-63.
19. Itier RJ, Taylor MJ. Source analysis of the N170 to faces and objects. *Neuroreport*. Jun 7 2004;15(8):1261-1265.
20. Schweinberger SR, Pickering EC, Jentsch I, Burton AM, Kaufmann JM. Event-related brain potential evidence for a response of inferior temporal cortex to familiar face repetitions. *Cognitive Brain Research*. Nov 2002;14(3):398-409.
21. Diamond R, Carey S. Why faces are and are not special: An effect of expertise. *Journal of Experimental Psychology*. 1986;115(2):107-117.
22. Gauthier II. What constrains the organization of the ventral temporal cortex? *Trends Cogn Sci*. Jan 2000;4(1):1-2.

23. Gauthier I, Skudkarski P, Gore J, Anderson A. Expertise for cars and birds recruits brain areas involved in face recognition. *Nature Neuroscience*. 2000;3(2):191-197.
24. Gauthier I, Williams P, Tarr MJ, Tanaka J. Training 'greeble' experts: a framework for studying expert object recognition processes. *Vision Res*. Aug 1998;38(15-16):2401-2428.
25. Tarr MJ, Gauthier I. FFA: a flexible fusiform area for subordinate-level visual processing automatized by expertise. *Nat Neurosci*. Aug 2000;3(8):764-769.
26. Rossion B, Gauthier I, Goffaux V, Tarr MJ, Crommelinck M. Expertise training with novel objects leads to left-lateralized facelike electrophysiological responses. *Psychol Sci*. May 2002;13(3):250-257.
27. Gauthier I, Nelson C. The development of face expertise. *Current opinion in neurobiology*. Vol 11. 2001:219-224.
28. James KH, James TW, Jobard G, Wong AC, Gauthier I. Letter processing in the visual system: different activation patterns for single letters and strings. *Cogn Affect Behav Neurosci*. Dec 2005;5(4):452-466.
29. McCandliss BD, Cohen L, Dehaene S. The visual word form area: expertise for reading in the fusiform gyrus. *Trends Cogn Sci*. Jul 2003;7(7):293-299.
30. Maurer U, Brem S, Kranz F, et al. Coarse neural tuning for print peaks when children learn to read. *Neuroimage*. Nov 1 2006;33(2):749-758.
31. Bentin S, Mouchetant-Rostaing Y, Giard MH, Echallier JF, Pernier J. ERP manifestations of processing printed words at different psycholinguistic levels: time course and scalp distribution. *J Cogn Neurosci*. May 1999;11(3):235-260.
32. Grossi G, Coch D, Coffey-Corina S, Holcomb PJ, Neville HJ. Phonological processing in visual rhyming: a developmental erp study. *J Cogn Neurosci*. Jul 1 2001;13(5):610-625.
33. Holcomb P, Coffey S, Neville H. Visual and auditory sentence processing: a developmental analysis using event-related brain potentials. *Developmental Neuropsychology*. 1992;8(2-3):203-241.
34. Maurer U, Brem S, Bucher K, Brandeis D. Emerging neurophysiological specialization for letter strings. *J Cogn Neurosci*. Oct 2005;17(10):1532-1552.

- 473 **35.** Wong AC, Gauthier I, Woroch B, DeBuse C, Curran T. An early electrophysiological response
474 associated with expertise in letter perception. *Cogn Affect Behav Neurosci.* Sep 2005;5(3):306-318.
- 475 **36.** Cohen L, Dehaene S, Naccache L, et al. The visual word form area: spatial and temporal
476 characterization of an initial stage of reading in normal subjects and posterior split-brain patients. *Brain.*
477 Feb 2000;123 (Pt 2):291-307.
- 478 **37.** Turkeltaub PE, Flowers DL, Verbalis A, Miranda M, Gareau L, Eden GF. The neural basis of hyperlexic
479 reading: an fMRI case study. *Neuron.* Jan 8 2004;41(1):11-25.
- 480 **38.** Osterling JA, Dawson G. Early recognition of children with autism: A study of first birthday home
481 videotapes. *Journal of Autism & Developmental Disorders.* Jun 1994;24(3):247-257.
- 482 **39.** Maestro S, Muratori F, Cavallaro MC, et al. Attentional skills during the first 6 months of age in autism
483 spectrum disorder. *J Am Acad Child Adolesc Psychiatry.* Oct 2002;41(10):1239-1245.
- 484 **40.** Klin A, Sparrow S, De Bildt A, Cicchetti D, Cohen D, Volkmar F. A normed study of face recognition
485 in autism and related disorders. *Journal of Autism & Developmental Disorders.* 1999;29(6):499-508.
- 486 **41.** Hobson R. The autistic child's appraisal of expressions of emotion. *Journal of Child Psychology and*
487 *Psychiatry.* 1986;27(3):321-342.
- 488 **42.** Schultz RT. Developmental deficits in social perception in autism: the role of the amygdala and fusiform
489 face area. *Int J Dev Neurosci.* Apr-May 2005;23(2-3):125-141.
- 490 **43.** Langdell T. Recognition of faces: an approach to the study of autism. *J Child Psychol Psychiatry.*
491 1978;19(3):255-268.
- 492 **44.** Hobson R, Ouston J, Lee A. What's in a face? The case of autism. *British Journal of Psychology.*
493 1988;79:441-453.
- 494 **45.** Jones W, Carr K, Klin A. Absence of preferential looking to the eyes of approaching adults predicts
495 level of social disability in 2-year-old toddlers with autism spectrum disorder. *Arch Gen Psychiatry.* Aug
496 2008;65(8):946-954.

46. Klin A, Jones W, Schultz R, Volkmar F, Cohen D. Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Arch Gen Psychiatry*. Sep 2002;59(9):809-816.
47. Schultz RT, Gauthier I, Klin A, et al. Abnormal ventral temporal cortical activity during face discrimination among individuals with autism and Asperger syndrome. *Archives of General Psychiatry*. Apr 2000;57(4):331-340.
48. O'Connor K, Hamm JP, Kirk IJ. The neurophysiological correlates of face processing in adults and children with Asperger's syndrome. *Brain Cogn*. Oct 2005;59(1):82-95.
49. McPartland J, Dawson G, Webb SJ, Panagiotides H, Carver LJ. Event-related brain potentials reveal anomalies in temporal processing of faces in autism spectrum disorder. *J Child Psychol Psychiatry*. Oct 2004;45(7):1235-1245.
50. Dawson G, Carver L, Meltzoff AN, Panagiotides H, McPartland J, Webb SJ. Neural correlates of face and object recognition in young children with autism spectrum disorder, developmental delay, and typical development. *Child Dev*. May-Jun 2002;73(3):700-717.
51. Webb SJ, Dawson G, Bernier R, Panagiotides H. ERP evidence of atypical face processing in young children with autism. *J Autism Dev Disord*. Oct 2006;36(7):881-890.
52. O'Connor K, Hamm JP, Kirk IJ. Neurophysiological responses to face, facial regions and objects in adults with Asperger's syndrome: An ERP investigation. *Int J Psychophysiol*. Jan 29 2007.
53. Dawson G, Webb SJ, Wijsman E, et al. Neurocognitive and electrophysiological evidence of altered face processing in parents of children with autism: implications for a model of abnormal development of social brain circuitry in autism. *Dev Psychopathol*. Summer 2005;17(3):679-697.
54. McCleery JP, Akshoomoff N, Dobkins KR, Carver LJ. Atypical face versus object processing and hemispheric asymmetries in 10-month-old infants at risk for autism. *Biol Psychiatry*. Nov 15 2009;66(10):950-957.

- 521 **55.** Dawson G, Webb SJ, McPartland J. Understanding the nature of face processing impairment in autism:
522 insights from behavioral and electrophysiological studies. *Dev Neuropsychol.* 2005;27(3):403-424.
- 523 **56.** Behrmann M, Thomas C, Humphreys K. Seeing it differently: visual processing in autism. *Trends Cogn*
524 *Sci.* Jun 2006;10(6):258-264.
- 525 **57.** Sasson NJ. The Development of Face Processing in Autism. *J Autism Dev Disord.* Mar 30 2006.
- 526 **58.** Grelotti DJ, Klin AJ, Gauthier I, et al. fMRI activation of the fusiform gyrus and amygdala to cartoon
527 characters but not to faces in a boy with autism. *Neuropsychologia.* 2005;43(3):373-385.
- 528 **59.** Boeschoten MA, Kenemans JL, van Engeland H, Kemner C. Face processing in Pervasive
529 Developmental Disorder (PDD): the roles of expertise and spatial frequency. *J Neural Transm.* Dec
530 2007;114(12):1619-1629.
- 531 **60.** Kanner L. Autistic disturbances of affective contact. *Nervous Child.* 1943;2:217-250.
- 532 **61.** Newman TM, Macomber D, Naples AJ, Babitz T, Volkmar F, Grigorenko EL. Hyperlexia in Children
533 with Autism Spectrum Disorders. *J Autism Dev Disord.* Sep 19 2006.
- 534 **62.** Nation K, Clarke P, Wright B, Williams C. Patterns of reading ability in children with autism spectrum
535 disorder. *J Autism Dev Disord.* Oct 2006;36(7):911-919.
- 536 **63.** Huemer SV, Mann V. A Comprehensive Profile of Decoding and Comprehension in Autism Spectrum
537 Disorders. *J Autism Dev Disord.* Nov 14 2009.
- 538 **64.** Klin A. Understanding circumscribed interests in autism spectrum disorders. Paper presented at: The
539 25th Annual TEACCH Conference 2004; Chapel Hill, NC.
- 540 **65.** Burd L, Kerbeshian J, Fisher W. Inquiry into the incidence of hyperlexia in a statewide population of
541 children with pervasive developmental disorder. *Psychol Rep.* 1985;57(1):236-238.
- 542 **66.** Grigorenko EL, Klin A, Pauls DL, Senft R, Hooper C, Volkmar F. A descriptive study of hyperlexia in a
543 clinically referred sample of children with developmental delays. *Journal of Autism & Developmental*
544 *Disorders.* Feb 2002;32(1):3-12.
- 545 **67.** Gadow K, Sprafkin J. *Child Symptom Inventories manual.* Stony Brook, NY: Checkmate Plus; 1994.

- 546 **68.** Elliott C. *The Differential Ability Scales*. 2nd ed. San Antonio, TX: Harcourt Assessment; 2007.
- 547 **69.** Wechsler D. *The Wechsler Intelligence Scale for Children - Fourth Edition*. San Antonio, TX: The
548 Psychological Corporation; 2003.
- 549 **70.** Wechsler D. *Manual for the Wechsler Adult Intelligence Scale*. 3rd ed. San Antonio, Tx: The
550 Psychological Corporation; 1997.
- 551 **71.** Lord C, Rutter M, Le Couteur A. Autism Diagnostic Interview-Revised: a revised version of a
552 diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *J*
553 *Autism Dev Disord*. 1994;24(5):659-685.
- 554 **72.** Lord C, Risi S, Lambrecht L, et al. The Autism Diagnostic Observation Schedule--Generic: A standard
555 measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism*
556 *& Developmental Disorders*. Jun 2000;30(3):205-223.
- 557 **73.** American Psychiatric Association. *Diagnostic and statistical manual of mental disorders : DSM-IV-TR*.
558 4th ed. Washington, DC: American Psychiatric Association; 2000.
- 559 **74.** Oldfield R. The assessment and analysis of handedness; The Edinburgh inventory. *Neuropsychologia*.
560 1971;9:97-113.
- 561 **75.** Psychological Corporation. *Wechsler Abbreviated Scale of Intelligence (WASI) manual*. San Antonio,
562 TX: Author; 1999.
- 563 **76.** Schneider W, Eschman A, Zuccolotto A. E-prime user's guide. Pittsburg, Psychology Software Tools
564 Inc. 2002.
- 565 **77.** Tucker DM. Spatial sampling of head electrical fields: the geodesic sensor net. *Electroencephalogr Clin*
566 *Neurophysiol*. Sep 1993;87(3):154-163.
- 567 **78.** Benton A, Sivan A, Hamsher K, Varney N, Spreen O. *Contributions to Neuropsychological Assessment*.
568 New York: Oxford University Press; 1994.
- 569 **79.** Woodcock R, McGrew K, Mather N. *Woodcock-Johnson III Tests of Achievement*. Itasca, IL: Riverside
570 Publishing; 2001.

- 571 **80.** Minschew NJ, Williams DL. The new neurobiology of autism: cortex, connectivity, and neuronal
572 organization. *Arch Neurol.* Jul 2007;64(7):945-950.
- 573 **81.** Krigolson OE, Pierce LJ, Holroyd CB, Tanaka JW. Learning to become an expert: reinforcement
574 learning and the acquisition of perceptual expertise. *J Cogn Neurosci.* Sep 2009;21(9):1834-1841.
- 575 **82.** Garibotto V, Scifo P, Gorini A, et al. Disorganization of anatomical connectivity in obsessive
576 compulsive disorder: A multi-parameter diffusion tensor imaging study in a subpopulation of patients.
577 *Neurobiol Dis.*In Press, Uncorrected Proof.
- 578 **83.** Friston KJ. Dysfunctional connectivity in schizophrenia. *World Psychiatry.* Jun 2002;1(2):66-71.
- 579 **84.** Murias M, Swanson JM, Srinivasan R. Functional connectivity of frontal cortex in healthy and ADHD
580 children reflected in EEG coherence. *Cereb Cortex.* Aug 2007;17(8):1788-1799.
- 581 **85.** Zhou Y, Dougherty JH, Jr., Hubner KF, Bai B, Cannon RL, Hutson RK. Abnormal connectivity in the
582 posterior cingulate and hippocampus in early Alzheimer's disease and mild cognitive impairment.
583 *Alzheimers Dement.* Jul 2008;4(4):265-270.
- 584 **86.** Webb SJ, Merkle K, Murias M, Richards T, Aylward E, Dawson G. ERP responses differentiate
585 inverted but not upright face processing in adults with ASD. *Soc Cogn Affect Neurosci.* May 19 2009.
- 586 **87.** Senju A, Tojo Y, Yaguchi K, Hasegawa T. Deviant gaze processing in children with autism: an ERP
587 study. *Neuropsychologia.* 2005;43(9):1297-1306.
- 588 **88.** Kemner C, Schuller AM, van Engeland H. Electrocortical reflections of face and gaze processing in
589 children with pervasive developmental disorder. *J Child Psychol Psychiatry.* Oct 2006;47(10):1063-
590 1072.
- 591 **89.** Jones W, Klin A. Heterogeneity and homogeneity across the autism spectrum: the role of development.
592 *J Am Acad Child Adolesc Psychiatry.* May 2009;48(5):471-473.
- 593 **90.** Jemel B, Mottron L, Dawson M. Impaired face processing in autism: fact or artifact? *J Autism Dev*
594 *Disord.* Jan 2006;36(1):91-106.

- 595 **91.** Dalton KM, Nacewicz BM, Alexander AL, Davidson RJ. Gaze-fixation, brain activation, and amygdala
596 volume in unaffected siblings of individuals with autism. *Biol Psychiatry*. Feb 15 2007;61(4):512-520.
- 597 **92.** Dalton KM, Nacewicz BM, Johnstone T, et al. Gaze fixation and the neural circuitry of face processing
598 in autism. *Nat Neurosci*. Apr 2005;8(4):519-526.
- 599 **93.** Taylor MJ, Batty M, Itier RJ. The faces of development: a review of early face processing over
600 childhood. *J Cogn Neurosci*. Oct 2004;16(8):1426-1442.
- 601 **94.** Martelli M, Majaj NJ, Pelli DG. Are faces processed like words? A diagnostic test for recognition by
602 parts. *J Vis*. 2005;5(1):58-70.
- 603 **95.** Scott LS, Tanaka JW, Sheinberg DL, Curran T. A reevaluation of the electrophysiological correlates of
604 expert object processing. *J Cogn Neurosci*. Sep 2006;18(9):1453-1465.
- 605
- 606

Figure Legends

Figure 1. Grand averaged waveforms across entire scalp for faces, houses, and inverted faces for typical participants and those with ASD. Electrodes of interest in right and left hemisphere are highlighted. Subpanels display the averaged waveform across the eight specified electrodes in each hemisphere for both groups.

Figure 2. Mean latency of the N170 component (in milliseconds) elicited by faces and houses for both groups in both hemispheres. Error bars represent +/- 1 S.E. Significance at the $p \leq .05$ level is indicated by *.

Figure 3. Grand averaged waveforms across entire scalp for letters and pseudoletters for typical participants and those with ASD. Electrodes of interest in right and left hemisphere are highlighted. Subpanels display the averaged waveform across the eight specified electrodes in each hemisphere for both groups.

Figure 4. Amplitude of the N170 component (in microVolts) elicited by letters and pseudoletters for both groups in both hemispheres. Error bars represent +/- 1 S.E. Significance at the $p \leq .05$ level is indicated by *, and significance at the $p \leq .01$ level is indicated by **.

Tables

Table 1

Participant Characteristics

	Typical (N=18)	ASD (N=36)
Number male (Percent)	15 (83.3)	32 (88.9)
Number White (Percent)	15 (83.3)	34 (94.4)
Number right handed (Percent)	16 (88.9)	31 (86.1)
Mean age (SD)	12.6 (2.4)	11.2 (3.4)
Mean Full Scale IQ (SD)	112.9 (13.4)	105.2 (17.3)

Table 2

N170 Latency

Hemisphere	Condition	<i>M</i> (milliseconds)	<i>SD</i>
Typical group			
Left	Faces	193.24	37.3
	Houses	215.59	43.4
	Inverted faces	194.82	35.3
	Letters	177.25	26.9
	Pseudoletters	179.56	26.8
Right	Faces	181.38	31.6
	Houses	217.50	36.4
	Inverted faces	188.85	32.7
	Letters	181.19	25.9
	Pseudoletters	176.03	28.5
ASD group			
Left	Faces	200.13	24.3
	Houses	221.67	31.2
	Inverted faces	205.42	27.7
	Letters	189.21	26.7
	Pseudoletters	178.83	19.3
Right	Faces	201.78	27.2
	Houses	219.38	27.8
	Inverted faces	204.66	31.3
	Letters	188.42	28.0
	Pseudoletters	183.51	23.6

Table 3

N170 amplitude

Hemisphere	Condition	<i>M</i> (microVolts)	<i>SD</i>
Typical group			
Left	Faces	0.39	1.7
	Houses	2.49	3.5
	Inverted faces	0.24	2.8
	Letters	- 2.27	3.0
	Pseudoletters	- 1.51	3.1
Right	Faces	- 0.16	2.0
	Houses	0.89	2.6
	Inverted faces	- 0.58	2.8
	Letters	- 3.03	4.2
	Pseudoletters	- 2.53	3.9
ASD group			
Left	Faces	- 0.32	3.4
	Houses	3.16	4.0
	Inverted faces	0.59	3.6
	Letters	- 3.09	3.6
	Pseudoletters	- 2.22	3.3
Right	Faces	0.76	3.6
	Houses	3.44	4.2
	Inverted faces	1.52	3.5
	Letters	- 3.57	3.1
	Pseudoletters	- 2.29	2.9

Table 4

Performance on behavioral measures

Measure	Typical (N = 17)		ASD (N = 36)	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Benton Face Recognition Score	41.41	3.6	37.11	4.8
Letter-Word Identification	108.41	9.9	105.67	15.0
Word Attack	101.41	9.7	103.86	11.6

Figure 1 TOP




Figure 2 TOP




Figure 3 TOP




Figure 4 TOP

