1. The probability of a leap year selected at random contain 53						
Sunday is:						
, ,	(c) 2/7 (d) 53/365					
2. A bag contains 3 red and	d 2 blue marbles. A marble is drawn at					
random. The probability of drawing a black ball is:						
(a) 3/5 (b) 2/5	(c) 0/5 (d) 1/5					
3. The probability that it wi	Il rain tomorrow is 0.85. What is the					
probability that it will not ra	in tomorrow					
(a) 0.25 (b) 0.145	(c) 3/20 (d) none of these					
4. What is the probability th	hat a number selected from the numbers					
(1, 2, 3,,15) is a multi						
	(c) 2/15 (d) 1/3					
	mes when we throw three coins?					
(a) 4 (b) 5	(c) 8 (d) 7					
` '	ime number selected at random from the					
numbers (1,2,3,35) is						
	(c) 13/35 (d) none of these					
	ity of an event and non event is :					
	(c) 0 (d) none of these.					
` '	ies are given; choose the correct answer					
for that which is not possible						
	(c) 7/5 (d) none of these.					
` '	simultaneously, than the probability of					
getting at least two heads, i						
(a) 1/4 (b) 3/8						
	ndom from the letters of the word					
	probability that the letter chosen has:					
	(c) 1 (d) none of these.					
11. A dice is thrown. Find th	ne probability of getting an even number.					
(A) 2/3 (B) 1	(C) 5/6 (D) 1/2					
(,,, _, _, _, _, _, _, _, _, _, _, _, _,	(0) 0, 0					
12. Two coins are thrown at the same time. Find the probability of						
getting both heads.						
(A) 3/4 (B) 1/4 (C) 1/	2 (D) 0					
13. Two dice are thrown sin	nultaneously. The probability of getting a					

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	9		
14. 100 cards	are numbered ne number.	from 1 to 10	0. Find the	probability of		
• • •	B) 27/50	(C) 1/4	(D)	29/100		
	blue ball is doul bag is:			f the probability n the number of		
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:						
(A) 143/150	(B) 147/1	50 (C) 1	1/25	(D) 1/50		
mixed thorou the probabilit	rked with numbers of the card with the card with the number (B) 1/10	is drawn fron per on card is	n this box r	andomly, then square.		
	h e probability o (B) 53/366					
probability of	drawn from a v getting a king (8) 3/26 (C) 7	of red suit.		cards. Find the		
equally likely	· ·	t pointing to	one of the r	number n odd number is:		
its outcome e result i.e. three probability th	onsists of tossicach time. Aryanee heads or threat at Aryan will lost 1/2 (C) 1	n wins if all the tails and lose the game.	he tosses g oses otherv			

22. Riya and Kaja same birthday is			oth will have the		
(A) 364/365			(D) 1/133225		
23. A number <i>x</i> i 2. Then the proba (A) 1/5 (B) 2/5	ability that x ² < 2	is?	ımbers -2, -1, 0 , 1,		
24. A jar contains a marble is drawing red is 2/3, then the (A) 10 (B) 6	n at random from ne number of wh	n the jar, the pro			
25. A number is 8 Then the probabi (A) 7/50 (B) 4/5	lity that it is a m	ultiple of 3 and	natural numbers. 4 is:		
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?					
a) $\frac{1}{7}$	b) $\frac{3}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$		
27. Runs scored 93, and 20. The s	•		es are 50, 70, 82,		
		c) 25.29	d) 25.69		
28. Find median a consecutive days a) 13, 15		, 4, 18, 13, 17.	eived on 9 d) 13, 16		
	sed up 4 times.		that tails turn up in		
a) ¹ / ₂ 30. X is a variate a) 8 b) 7	b) $\frac{1}{3}$ between 0 and 3		d) $\frac{1}{6}$ E(X²) is		
31.The random v	ariables X and \	have variances	0.2 and 0.5		

respectively. Let Z= 5X-2Y. The variance of Z is?

3

~. ~~~~!!!!	?		one is not possible in
probability a) P(x) = 1	75/5 NAC TOURS	P(x) = 3	
c) $P(x) = 0$.	5 d) P(x	(x) = -0.5	
33.If E(x) :	= 2 and E(z) = 4	The second secon) =?
a) 2	b) 6	c) 0	d) Insufficient data
34.The cov	ariance of two	independent	random variable is
a) 1	b) 0	c) - 1	d) Undefined
35.If Σ P(x	$(x) = k^2 - 8$ then,	the value of I	k is?
a) 0	b) 1	c) 3	d) Insufficient dat
is always?			, the sum of all probabilities
1.600		y distribution c) 1	, the sum of all probabilities d) Undefined
is always? a) 0 38.If the p	b) Infinite	c) 1	
is always? a) 0	b) Infinite robability of hit	c) 1	d) Undefined

a) 3

b) 4

c) 5

d) 7

,		s 1 b) Mean is 1 a		
c) Mean is 0	and variance is	s ∞ d) Mean is o	o and varianc	e is 0
42 Variance	of a random v	ariable X is given	hv	
a) E(X)		c) E(X2) - (E		d) (E(X))2
, -(,	,,			-, (-(-,-,-
		able X is given by		
a) E(X)	b) E(X2)	c) E(X2) - (E(X	())2	d) (E(X))2
11 Moon of	a conctant 'a' i			
a) 0	b) a	c) a/2	d) 1	
a) U	b) a	C) a/ Z	u) i	
45.Variance	of a constant '	a' is		
a) 0	b) a	c) a/2	d) 1	
4 1 -1	•	C >//O		
46.Find the r	nean and varia	ince of X?		

a) 2, 4/3	b) 3, 4/3	c) 2, 2/3	d) 3, 2/3

2/9

1/9

3/9

47. Find the expectation of a random variable X?

2/9

X

1/9

f(x)

	X	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
C).5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

5

a)

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard Deviation?

- b) \sqrt{pq} c) (np)2