

Segmentation:

Mean-shift

Dr. Tushar Sandhan

Introduction

- Number of segments?



Introduction

- Number of segments?




Introduction

- Number of segments?




Mean-shift


- Mean-shift clustering
 - iterative steepest ascent method
 - seeks peaks of probability density in feature space
 - finds modes or local maxima
 - it tries to find all possible cluster centres
 - no need of initial guess of K clusters




Mean-shift



Mean-shift




Mean-shift




Non-parametric
density estimation

Mean-shift




Non-parametric
density estimation


Mean-shift




Non-parametric
density estimation




Mean-shift




Mean-shift



Mean-shift





Mean-shift




Mean-shift

- Density estimation
 - find the underlying distribution that generates the given data
- Non-parametric density estimation
 - use data points to define distribution
 - put a small probability mass around each data-point via kernel
 - e.g. Gaussian kernel




Mean-shift




Data point density is similar to PDF value

Mean-shift



Data point density is similar to PDF value



Courtesy: T. Tappen

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

points \mathbf{x}_i , $i = 1, \dots, n$ in the d -dimensional space R^d

kernel $K(\mathbf{x})$

symmetric positive definite $d \times d$ bandwidth matrix \mathbf{H}

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

d -variate kernel $K(\mathbf{x})$ is a bounded function
compact support satisfying

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

points $\mathbf{x}_i, i = 1, \dots, n$ in the d -dimensional space R^d

kernel $K(\mathbf{x})$

symmetric positive definite $d \times d$ bandwidth matrix \mathbf{H}

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

d -variate kernel $K(\mathbf{x})$ is a bounded function
compact support satisfying

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

points \mathbf{x}_i , $i = 1, \dots, n$ in the d -dimensional space R^d

kernel $K(\mathbf{x})$

symmetric positive definite $d \times d$ bandwidth matrix \mathbf{H}

Mean-shift

- Kernel density estimation
 - find the underlying distribution
 - that generates the given data
 - in non-parametric way

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

d -variate kernel $K(\mathbf{x})$ is a bounded function
compact support satisfying

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^T K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

points \mathbf{x}_i , $i = 1, \dots, n$ in the d -dimensional space R^d

kernel $K(\mathbf{x})$

symmetric positive definite $d \times d$ bandwidth matrix \mathbf{H}

where c_K is a constant

Mean-shift

- Kernel density estimation
 - multivariate kernel can be generated from a symmetric univariate $K_1(x)$ in two different ways:

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^T K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

Mean-shift

- Kernel density estimation
 - multivariate kernel can be generated from a symmetric univariate $K_1(x)$ in two different ways:

$K^P(\mathbf{x})$ is obtained from the product of the univariate

$$K^P(\mathbf{x}) = \prod_{i=1}^d K_1(x_i)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^T K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

Mean-shift

- Kernel density estimation
 - multivariate kernel can be generated from a symmetric univariate $K_1(x)$ in two different ways:

$K^P(\mathbf{x})$ is obtained from the product of the univariate

$$K^P(\mathbf{x}) = \prod_{i=1}^d K_1(x_i)$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^T K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

Mean-shift

- Kernel density estimation
 - multivariate kernel can be generated from a symmetric univariate $K_1(x)$ in two different ways:

$K^P(\mathbf{x})$ is obtained from the product of the univariate

$$K^P(\mathbf{x}) = \prod_{i=1}^d K_1(x_i)$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^T K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

it suffices to define the function $k(x)$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

it suffices to define the function $k(x)$

constant $c_{k,d}$, which makes $K(\mathbf{x})$ integrate to one

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$
 - Epanechnikov kernel

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$
 - Epanechnikov kernel

$$k_E(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & x > 1, \end{cases}$$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- Epanechnikov kernel

$$k_E(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & x > 1, \end{cases}$$

$$K_E(\mathbf{x}) = \begin{cases} \frac{1}{2} c_d^{-1} (d+2) (1 - \|\mathbf{x}\|^2) & \|\mathbf{x}\| \leq 1 \\ 0 & \text{otherwise,} \end{cases}$$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- Epanechnikov kernel

$$k_E(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & x > 1, \end{cases}$$

$$K_E(\mathbf{x}) = \begin{cases} \frac{1}{2} c_d^{-1} (d+2) (1 - \|\mathbf{x}\|^2) & \|\mathbf{x}\| \leq 1 \\ 0 & \text{otherwise,} \end{cases}$$

c_d is the volume of the unit d -dimensional sphere

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- Epanechnikov kernel

$$k_E(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & x > 1, \end{cases}$$

$$K_E(\mathbf{x}) = \begin{cases} \frac{1}{2} c_d^{-1} (d+2) (1 - \|\mathbf{x}\|^2) & \|\mathbf{x}\| \leq 1 \\ 0 & \text{otherwise,} \end{cases}$$

c_d is the volume of the unit d -dimensional sphere

- non-differentiable at boundary

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$
 - exponential profile

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2} \mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$
 - exponential profile

$$k_N(x) = \exp\left(-\frac{1}{2}x\right) \quad x \geq 0$$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2}\mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation
 - radial symmetric kernels $K^S(\mathbf{x})$
 - profile of a kernel $k(x)$
 - exponential profile

$$k_N(x) = \exp\left(-\frac{1}{2}x\right) \quad x \geq 0$$

- yields multivariate normal kernel

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2}\mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- exponential profile

$$k_N(x) = \exp\left(-\frac{1}{2}x\right) \quad x \geq 0$$

- yields multivariate normal kernel

$$K_N(\mathbf{x}) = (2\pi)^{-d/2} \exp\left(-\frac{1}{2}\|\mathbf{x}\|^2\right)$$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2}\mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- exponential profile

$$k_N(x) = \exp\left(-\frac{1}{2}x\right) \quad x \geq 0$$

- yields multivariate normal kernel

$$K_N(\mathbf{x}) = (2\pi)^{-d/2} \exp\left(-\frac{1}{2}\|\mathbf{x}\|^2\right)$$

- final density estimator with this kernel

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2}\mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

- Kernel density estimation

- radial symmetric kernels $K^S(\mathbf{x})$
- profile of a kernel $k(x)$
- exponential profile

$$k_N(x) = \exp\left(-\frac{1}{2}x\right) \quad x \geq 0$$

- yields multivariate normal kernel

$$K_N(\mathbf{x}) = (2\pi)^{-d/2} \exp\left(-\frac{1}{2}\|\mathbf{x}\|^2\right)$$

- final density estimator with this kernel

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$$K(\mathbf{x}) = c_{k,d} k(\|\mathbf{x}\|^2)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n K_{\mathbf{H}}(\mathbf{x} - \mathbf{x}_i)$$

$$K_{\mathbf{H}}(\mathbf{x}) = |\mathbf{H}|^{-1/2} K(\mathbf{H}^{-1/2}\mathbf{x})$$

$$\begin{aligned} \int_{R^d} K(\mathbf{x}) d\mathbf{x} &= 1 & \lim_{\|\mathbf{x}\| \rightarrow \infty} \|\mathbf{x}\|^d K(\mathbf{x}) &= 0 \\ \int_{R^d} \mathbf{x} K(\mathbf{x}) d\mathbf{x} &= 0 & \int_{R^d} \mathbf{x} \mathbf{x}^\top K(\mathbf{x}) d\mathbf{x} &= c_K \mathbf{I}, \end{aligned}$$

$K^S(\mathbf{x})$ from rotating $K_1(x)$ in R^d

$$K^S(\mathbf{x}) = a_{k,d} K_1(\|\mathbf{x}\|)$$

$$a_{k,d}^{-1} = \int_{R^d} K_1(\|\mathbf{x}\|) d\mathbf{x}$$

Mean-shift

$$x_1 = x_0 + \eta \nabla f(x_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

Mean-shift

$$x_1 = x_0 + \eta \nabla f(x_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

Define a function of a kernel $g(x) = -k'(x)$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

Define a function of a kernel $g(x) = -k'(x)$

$$\begin{aligned} \hat{\nabla} f_{h,K}(\mathbf{x}) &= \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x}_i - \mathbf{x}) g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \\ &= \frac{2c_{k,d}}{nh^{d+2}} \left[\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \right] \left[\frac{\sum_{i=1}^n \mathbf{x}_i g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)}{\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)} - \mathbf{x} \right] \end{aligned}$$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

Define a function of a kernel $g(x) = -k'(x)$

$$\begin{aligned} \hat{\nabla} f_{h,K}(\mathbf{x}) &= \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x}_i - \mathbf{x}) g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \\ &= \frac{2c_{k,d}}{nh^{d+2}} \left[\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \right] \left[\frac{\sum_{i=1}^n \mathbf{x}_i g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)}{\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)} - \mathbf{x} \right] \end{aligned}$$

$$\mathbf{m}_{h,G}(\mathbf{x}) = \frac{\sum_{i=1}^n \mathbf{x}_i g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)}{\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)} - \mathbf{x}$$

Mean-shift

$$\mathbf{x}_1 = \mathbf{x}_0 + \eta \nabla f(\mathbf{x}_0)$$

$$\hat{f}(\mathbf{x}) = \frac{1}{nh^d} \sum_{i=1}^n K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$$\hat{f}_{h,K}(\mathbf{x}) = \frac{c_{k,d}}{nh^d} \sum_{i=1}^n k\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$

$c_{k,d}$ is volume of unit sphere $\in R^d$

at the modes: $\nabla f(\mathbf{x}) = \mathbf{0}$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) \equiv \nabla \hat{f}_{h,K}(\mathbf{x}) = \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x} - \mathbf{x}_i) k'\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)$$


Define a function of a kernel $g(x) = -k'(x)$

$$\begin{aligned} \hat{\nabla} f_{h,K}(\mathbf{x}) &= \frac{2c_{k,d}}{nh^{d+2}} \sum_{i=1}^n (\mathbf{x}_i - \mathbf{x}) g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \\ &= \frac{2c_{k,d}}{nh^{d+2}} \left[\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right) \right] \left[\frac{\sum_{i=1}^n \mathbf{x}_i g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)}{\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)} - \mathbf{x} \right] \end{aligned}$$


$$\mathbf{m}_{h,G}(\mathbf{x}) = \frac{\sum_{i=1}^n \mathbf{x}_i g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)}{\sum_{i=1}^n g\left(\left\|\frac{\mathbf{x} - \mathbf{x}_i}{h}\right\|^2\right)} - \mathbf{x}$$

$$\hat{\nabla} f_{h,K}(\mathbf{x}) = \hat{f}_{h,G}(\mathbf{x}) \frac{2c_{k,d}}{h^2 c_{g,d}} \mathbf{m}_{h,G}(\mathbf{x}) \quad \mathbf{m}_{h,G}(\mathbf{x}) = \frac{1}{2} h^2 c \frac{\hat{\nabla} f_{h,K}(\mathbf{x})}{\hat{f}_{h,G}(\mathbf{x})}$$


Mean-shift




Mean-shift




Mean-shift





Mean-shift



Mean-shift



Mean-shift



Mean-shift

- Algorithm

- transform image into feature space
 - initialize window at each feature point
 - for each window
 - compute mean shift vector $m(x)$
 - move density estimation window by $m(x)$
 - repeat till convergence
 - merge windows that end up near same peak
-

Mean-shift

■ Pros

- automatically finds various number of modes
- no need of initial guessing of cluster centres
- does not assume spherical clusters
- robust to outliers
- just a single para (window size w)

■ Cons

- bandwidth or windows size is an imp. Para
 - slight change in w , translates varied output
- computationally expensive
 - complexity: $O(n^2T)$
- not scalable with dimensionality

Mean-shift

input



Mean-shift

input



Mean-shift

input



mean shift segmentation



Mean-shift

input



mean shift segmentation



Mean-shift


input




mean shift segmentation




mean shift region boundaries



Mean-shift




Mean-shift



(a)


(b)





input image



output image




Mean-shift




input image




output image



Pixel Distribution Before Meanshift



Pixel Distribution After Meanshift



Mean-shift

- Surface reconstruction



- Image denoising

Mean-shift


- Surface reconstruction

- Image denoising




Mean-shift

- Surface reconstruction




- Image denoising



Mean-shift

- Surface reconstruction




- Image denoising



Mean-shift

- Surface reconstruction




- Image denoising



Mean-shift

- Surface reconstruction




- Image denoising



Conclusion

- Mean-shift




Conclusion

- Mean-shift

□ Mean-shift

- Number of cluster specification is not needed
- Mode seeking algorithm
- Computationally expensive




Conclusion

- Mean-shift

□ Mean-shift

- Number of cluster specification is not needed
- Mode seeking algorithm
- Computationally expensive



□ Mean-shift also can do

- surface reconstruction
- Image filtering