

Swap_Pricing

December 24, 2017

1 Get discount factor for JPY

- input: MoneyMarket (short term interest rate), Swap rate.
- output: discount factors for each tenor listed by MoneyMarket and Swap rate.

1.1 Pricing

1.1.1 Swap pricing formula

The value of the exchange between a float and a fixed side is given by

$$V = \sum_{i=1}^N L(t_{i-1}, t_i) \times DF(t_i) \times \delta_i - \sum_{i=1}^N SwapRate \times DF(t_i) \times \delta_i,$$

where $L(t_{i-1}, t_i)$ is the float interest rate between t_{i-1} and t_i , $DF(t_i)$ is a discount factor, δ_i is a day-count-fraction and $SwapRate$ is a Swap rate which means a par rate for a swap trade.

1.1.2 Bootstrap method for getting discount factors

Discount factors as of today can be estimated from a par swap trade which corresponds to $V = 0$ under swap pricing formula. For example, let us consider a swap trade with maturity of 1.5 year. The discount factor for 1.5 year $DF(t_{1.5Y})$ is calculated by solving the following equation:

$$\sum_{i=1}^3 L(t_{i-1}, t_i) \times DF(t_i) \times \delta = \sum_{i=1}^3 SwapRate(1.5Y) \times DF(t_i) \times \delta$$

where a quoted swap rate is used for $SwapRate(1.5Y)$, the day-count-fraction δ is assumed 6 month and the float side interest rate is assumed that a following model expressed as

$$L(t_{i-1}, t_i) = \frac{1}{\delta} \left(\frac{DF(t_{i-1})}{DF(t_i)} - 1 \right).$$

The above equation can be solved by using $DF(t_{0.5Y})$, $DF(t_{1.0Y})$ and the float interest rate which is defined as above equation. As a result, the discount factor $DF(t_{1.5Y})$ is given by

$$DF(t_{1.5Y}) = \frac{1}{(1 + \delta \times SwapRate(1.5Y))} \left(DF(t_0) - SwapRate(1.5Y) \times \delta \times (DF(t_{0.5Y}) + DF(t_{1.0Y})) \right),$$

where $DF(t_{0.5Y})$ and $DF(t_{1.0Y})$ is calculated by using a quoted LIBOR (the rate of Money Market). The short rate of Money Market means spot rate, where the cashflows is expressed as only two terms. For example, $DF(t_{0.5Y})$ is given by

$$DF(t_{0.5Y}) = \frac{1}{(1 + \delta \times L(0.0Y, 0.5Y))},$$

where $L(0.0Y, 0.5Y)$ is the LIBOR rate between today and 6 month later. Discount factors after $t_{1.5Y}$ can be calculated by the same way as the derivation of $DF(t_{1.5Y})$. This method of getting discount factors gradually is called Bootstrap method.

```
In [11]: import matplotlib.pyplot as plt
import numpy as np
import datetime

class getDF_moneymarket:
    ''' def __init__(self, libor_rate, start_day, end_day):
        self.libor_rate = libor_rate
        self.start_day = start_day
        self.end_day = end_day
        self.datetime_obj_start = datetime.datetime.strptime(start_day, '%Y/%m/%d')
        self.datetime_obj_end = datetime.datetime.strptime(end_day, '%Y/%m/%d')
        self.daycount = (self.datetime_obj_end - self.datetime_obj_start).days / 360
        self.discount_factor = 0
    '''

    def __init__(self, today, array_ccy):
        self._start_day = today

    def getDF(self, seq_moneymarket):

In [12]: DF = getDF_moneymarket(0.2, '2017/12/18', '2019/12/30')
print(DF.discount_factor)
print(DF.getDF())
print(DF.discount_factor)

0
[0.7081038552321007, '2017/12/18', '2019/12/30']
[0.7081038552321007, '2017/12/18', '2019/12/30']

In [3]: DF1 = getDF_moneymarket(0.3, '2017/12/18', '2018/3/20')
DF1.getDF()

Out[3]: 0.9287925696594427

In [66]: %matplotlib inline
import numpy as np
import csv
```

```

import time
import datetime
import matplotlib.pyplot as plt

with open('sample_moneymarket.csv', 'r') as csvfile:
    reader_obj = csv.reader(csvfile)
    # rewritten header_obj by using next method(???)
    header_obj = next(reader_obj)
    mm_list = []
    for row in reader_obj:
        mm_list.append(row)

mm_list

def get_DF(money_market_list):
    list_len = len(money_market_list)
    discount_factor = np.zeros(list_len*2).reshape(list_len, 2)
    discount_factor = [["", 0.0] for i in range(list_len)]
    day_count_fraction = np.zeros(list_len)
    # substitution the kinf of trade
    for i in range(0, list_len):
        discount_factor[i][0] = money_market_list[i][0]
    # calc daycount-fraction
    convention = 360.0
    for i in range(0, len(day_count_fraction)):
        day_count_fraction[i] = calc_daycount(money_market_list[i][1], money_market_list[i][0], convention)
    # calculate DF of 0/N
    discount_factor[0][1] = 1.0 / (1.0 + day_count_fraction[0] * float(money_market_list[0][1]))
    # calculate DF of T/N
    discount_factor[1][1] = discount_factor[0][1] / (1.0 + day_count_fraction[1]*float(money_market_list[1][1]))
    # calculate DF after 1W
    for i in range(2, list_len):
        discount_factor[i][1] = discount_factor[1][1] / (1.0 + day_count_fraction[i] * float(money_market_list[i][1]))
    return discount_factor

def calc_daycount(start_day, end_day, convention):
    datetime_obj_start = datetime.datetime.strptime(start_day, '%Y/%m/%d')
    datetime_obj_end = datetime.datetime.strptime(end_day, '%Y/%m/%d')
    daycount = (datetime_obj_end - datetime_obj_start).days / convention
    return daycount

def draw_DF(seq_discount_factor):
    list_len = len(seq_discount_factor)
    seq_DF = np.zeros(list_len)
    for i in range(0, list_len):

```

```
seq_DF[i] = seq_discount_factor[i][1]
plt.plot(seq_DF)
plt.ylim([0,1.0])
```

```
list_discountfactor = get_DF(mm_list)
draw_DF(list_discountfactor)
```

```
In [43]: [[] for i in range(5)]
```

```
Out[43]: [[], [], [], [], []]
```

```
In [55]: calc_daycount(mm_list[0][1], mm_list[0][2], 360)
         calc_daycount(mm_list[5][1], mm_list[5][2], 360)
```

```
Out[55]: 0.16666666666666666
```

```
In [56]: float(mm_list[0][3])
```

```
Out[56]: 0.00141
```

1.1.3 エラーメッセージ

1.1.4 解決策

- 数値と文字列が混ざっているのでどちらかに統一すべし