

Data Analytics using Python

Week 10 Question Bank Answer

1 Describe the Chi-square test of independence.

Ans.1) The Chi-square test of independence is a statistical test used to analyze the relationship between two categorical variables. It is used to determine whether the two variables are independent or related.

The key steps in the Chi-square test of independence are:

1. Set up the null and alternative hypotheses:
 - H_0 : The two variables are independent
 - H_1 : The two variables are not independent
2. Construct a contingency table that shows the observed frequencies for each combination of the two variables.
3. Calculate the expected frequencies under the null hypothesis that the variables are independent.
4. Calculate the Chi-square test statistic using the formula: $\chi^2 = \sum (O - E)^2 / E$ Where:
 - O = Observed frequency
 - E = Expected frequency
5. Determine the degrees of freedom, which is calculated as $(\text{number of rows} - 1) \times (\text{number of columns} - 1)$.
6. Compare the calculated Chi-square statistic to the critical value from the Chi-square distribution table at the desired significance level and degrees of freedom.
7. If the calculated Chi-square statistic is greater than the critical value, reject the null hypothesis and conclude that the two variables are not independent. Otherwise, do not reject the null hypothesis and conclude that the two variables are independent.

The Chi-square test of independence is useful in situations where you want to analyze the relationship between two categorical variables and determine whether they are independent or related. It is a widely used statistical test in various fields, such as social sciences, marketing, and healthcare.

2 What are two main types of Chi-Square tests?

Ans.2) The two main types of Chi-Square tests discussed are:

1. Chi-Square Test of Independence:

- This test is used to analyze the relationship between two categorical variables and determine whether they are independent or related.
- The null hypothesis is that the two variables are independent, while the alternative hypothesis is that they are not independent.
- The test statistic is calculated using the formula $\chi^2 = \sum (O - E)^2 / E$, where O is the observed frequency and E is the expected frequency under the null hypothesis.

2. Chi-Square Goodness-of-Fit Test:

- This test is used to compare the observed frequencies of a categorical variable to the expected frequencies based on a hypothesized probability distribution, such as the Poisson distribution.
- The null hypothesis is that the population has the hypothesized probability distribution, while the alternative hypothesis is that it does not.
- The test statistic is calculated using the formula $\chi^2 = \sum (f_o - f_e)^2 / f_e$, where f_o is the observed frequency and f_e is the expected frequency.

In summary, the Chi-Square test of independence is used to analyze the relationship between two categorical variables, while the Chi-Square goodness-of-fit test is used to compare observed frequencies to expected frequencies based on a hypothesized probability distribution.

3 What are limitations of Chi-Square Test?

Ans.3) Some key limitations of the Chi-Square test are:

1. Assumption of expected frequencies:

- The Chi-Square test requires that the expected frequencies in each cell of the contingency table be at least 5.1
- Violation of this assumption can lead to inaccurate results and invalid conclusions.

2. Sensitivity to sample size:

- The Chi-Square test is sensitive to sample size. With a large sample size, even small differences between observed and expected frequencies can lead to a significant result, rejecting the null hypothesis.1
- Conversely, with a small sample size, the test may lack the power to detect significant differences.

3. Inability to measure strength of association:

- The Chi-Square test only determines whether there is a significant relationship between the variables, but it does not provide a measure of the strength or magnitude of the association.1
- Additional measures, such as the phi coefficient or Cramer's V, are needed to quantify the strength of the relationship.

4. Limitations with ordinal variables:

- When the variables involved are ordinal, the Chi-Square test may not be the most appropriate choice, as it does not take into account the ordered nature of the data.1
- In such cases, other tests, such as the Mantel-Haenszel test or the Cochran-Armitage trend test, may be more suitable.

5. Interpretation challenges:

- Interpreting the results of the Chi-Square test can be challenging, especially when the test indicates a significant relationship but the underlying reasons for the relationship are not clear.1
- Additional analysis and interpretation may be required to understand the nature and implications of the relationship.

In summary, the Chi-Square test has several limitations, including assumptions about expected frequencies, sensitivity to sample size, inability to measure strength of association, and challenges in interpreting the results, particularly when dealing with ordinal variables.

4 What is the difference between t-test and chi-square?

Ans.4)

Characteristic	t-test	Chi-Square Test
Purpose	Used to compare the means of two groups or samples to determine if they are significantly different.	Used to analyze the relationship between two categorical variables and determine if they are independent or related.
Data Type	Requires continuous/interval data.	Requires categorical/nominal data.
Null Hypothesis	The means of the two groups are equal ($H_0: \mu_1 = \mu_2$).	The two variables are independent (H_0 : the variables are independent).
Test Statistic	The t-statistic, which follows a t-distribution.	The chi-square statistic, which follows a chi-square distribution.
Assumptions	Assumes the data follows a normal distribution and the samples are independent.	Assumes the expected frequencies in each cell of the contingency table are at least 5.
Degrees of Freedom	Calculated as $n_1 + n_2 - 2$, where n_1 and n_2 are the sample sizes.	Calculated as $(\text{number of rows} - 1) \times (\text{number of columns} - 1)$.
Interpretation	Compares the means of two groups and determines if the difference is statistically significant.	Determines if the two categorical variables are independent or related.

In summary, the t-test is used for continuous data to compare means, while the chi-square test is used for categorical data to analyze the relationship between variables. The test statistics, assumptions, and interpretations differ between the two tests.

5 What are the characteristics of chi-square?

Ans.5) The key characteristics of the chi-square test are:

1. Purpose:

- The chi-square test is used to analyze the relationship between two categorical variables and determine if they are independent or related.

2. Data Type:

- The chi-square test requires categorical or nominal data.

3. Null and Alternative Hypotheses:

- The null hypothesis (H_0) is that the two variables are independent.
- The alternative hypothesis (H_1) is that the two variables are not independent.

4. Test Statistic:

- The test statistic used in the chi-square test is the chi-square statistic, which follows a chi-square distribution.¹
- The formula for the chi-square statistic is: $\chi^2 = \sum (O - E)^2 / E$, where O is the observed frequency and E is the expected frequency under the null hypothesis.

5. Assumptions:

- The chi-square test assumes that the expected frequencies in each cell of the contingency table are at least 5.¹
- Violation of this assumption can lead to inaccurate results and invalid conclusions.

6. Degrees of Freedom:

- The degrees of freedom for the chi-square test are calculated as (number of rows - 1) $\times$ (number of columns - 1).

7. Interpretation:

- The chi-square test determines whether the two categorical variables are independent or related, but it does not provide a measure of the strength of the association.¹
- Additional measures, such as the phi coefficient or Cramer's V , are needed to quantify the strength of the relationship.

In summary, the chi-square test is a widely used statistical test for analyzing the relationship between two categorical variables, with specific characteristics related to its purpose, data type, hypotheses, test statistic, assumptions, degrees of freedom, and interpretation.

6 What do you mean by cluster analysis ? State its purpose.

Ans.6) Cluster analysis is a statistical technique used to group similar objects or observations into clusters. The purpose of cluster analysis is:

1. To find groups in data:

- Cluster analysis aims to identify natural groupings or clusters within a dataset, where objects in the same cluster are more similar to each other than to objects in different clusters.

2. To discover patterns and structure:

- By grouping similar objects together, cluster analysis can help uncover underlying patterns and structures within the data that may not be immediately apparent.

3. Exploratory data analysis:

- Cluster analysis is often used as an exploratory data analysis technique to gain insights into the data and generate hypotheses for further investigation.

4. Segmentation:

- Cluster analysis can be used to segment a population into meaningful subgroups, which is useful in applications like market segmentation, customer profiling, and targeted marketing.

5. Dimensionality reduction:

- Cluster analysis can be used to reduce the dimensionality of a dataset by representing each cluster with a single representative object or centroid.

In summary, the primary purpose of cluster analysis is to discover natural groupings or clusters within a dataset, in order to better understand the structure and patterns in the data, without any prior knowledge about the groups.

7 Differentiate between cluster analysis and discriminant analysis.

Ans.7)

Characteristic	Cluster Analysis	Discriminant Analysis
Purpose	Identifies natural groupings or clusters within a dataset based on similarity between observations.	Predicts group membership of observations based on independent variables.
Data Type	Typically used with unsupervised learning on unlabeled data to find patterns.	Supervised learning method used with labeled data to classify observations into predefined groups.
Approach	Unsupervised learning technique where groups are formed based on similarity without predefined classes.	Supervised learning technique where a model is trained on labeled data to classify new observations.
Objective	Reveals inherent structures or patterns in data without prior knowledge of group membership.	Focuses on predicting the group to which an observation belongs based on known group memberships.
Assumptions	Does not assume any prior knowledge about the groups or classes in the data.	Assumes that the independent variables are normally distributed and have equal covariance matrices.
Output	Outputs clusters or groupings of observations based on similarity measures.	Outputs a discriminant function that can be used to predict the group membership of new observations.
Application	Commonly used in market segmentation, image segmentation, and anomaly detection.	Applied in fields like biology, finance, and social sciences for classification and prediction purposes.
Example	Grouping customers based on purchasing behavior without predefined segments.	Predicting whether a patient has a specific disease based on various medical test results.

In summary, cluster analysis is an unsupervised learning technique used to identify natural groupings in data, while discriminant analysis is a supervised learning method focused on predicting group membership based on independent variables and known group labels.

8 What are different types of data? Why there is need to standardize data?

Ans.8) There are several different types of data:

1. Nominal Data:

- Categorical data that has no inherent order or ranking.
- Examples include gender, race, marital status, etc.

2. Ordinal Data:

- Categorical data that has a clear order or ranking.
- Examples include education level, income bracket, Likert scale responses, etc.
-

3. Interval Data:

- Numerical data where the difference between values is meaningful, but there is no true zero point.
- Examples include temperature (Celsius or Fahrenheit), dates, etc.

4. Ratio Data:

- Numerical data with a true zero point, allowing for meaningful comparisons of ratios.
- Examples include height, weight, income, etc.

The need to standardize data arises due to the different scales and units of measurement across different variables.

Standardization is important for several reasons:

1. Comparability:

- Standardizing the data puts all variables on a common scale, allowing for meaningful comparisons and analysis.

2. Reducing the impact of outliers:

- Standardization can help reduce the influence of outliers, which can skew the analysis.

3. Improving model performance:

- Many statistical and machine learning models perform better when the input variables are on a similar scale.

4. Interpretability:

- Standardized data can be more easily interpreted, as the units are now comparable across variables.
-

5. Addressing different measurement scales:

- Standardization allows for the integration of variables with different measurement scales, such as combining ordinal and interval data.

In summary, standardizing data is crucial to ensure comparability, reduce the impact of outliers, improve model performance, enhance interpretability, and enable the integration of variables with different measurement scales.

9 Explain the effect of standardization using example.

Ans.9) The key effects of data standardization can be explained using the following example:

Example: Gasoline Preference Versus Income Category

Before Standardization:

- The data contained variables with different scales and units, such as income categories (less than \$30,000, \$30,000 to \$49,999, etc.) and types of gasoline (regular, premium, extra premium).
- Directly comparing these variables would be challenging, as the scales are not comparable.

Effect of Standardization:

1. Comparability:

- By standardizing the data, the variables are put on a common scale, allowing for meaningful comparisons between the income categories and gasoline preferences.

2. Reducing the impact of outliers:

- Standardization can help reduce the influence of outliers in the data, which could have skewed the analysis.

3. Improving model performance:

- Many statistical and machine learning models, such as the chi-square test of independence used in the example, perform better when the input variables are on a similar scale.

4. Interpretability:

- The standardized data is more easily interpreted, as the units are now comparable across the income and gasoline preference variables.

In the example, the chi-square test of independence was performed on the standardized data, which allowed for a valid analysis of the relationship between income and gasoline preference. The test statistic and conclusions drawn from the analysis were more reliable and meaningful due to the standardization of the data.

In summary, standardizing the data in this example improved the comparability, reduced the impact of outliers, enhanced the model performance, and made the results more interpretable, leading to a more robust and reliable analysis of the relationship between income and gasoline preference.

10 Give the metrics for computation of distances between the objects.

Ans.10) The metrics commonly used for the computation of distances between objects in cluster analysis are:

1. Euclidean Distance:

- The Euclidean distance is the most common distance metric used in cluster analysis.
- It calculates the straight-line distance between two points in a multidimensional space.
- The formula for Euclidean distance between two points $P=(p_1, p_2, \dots, p_n)$ and $Q=(q_1, q_2, \dots, q_n)$ is:
- **Euclidean Distance** $= \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2 + \dots + (p_n - q_n)^2}$

2. Manhattan Distance (City Block Distance):

- The Manhattan distance is the sum of the absolute differences between the coordinates of two points.
- It is calculated as the sum of the absolute differences in the coordinates along each dimension.
- The formula for Manhattan distance between two points P and Q is:
- **Manhattan Distance** $= |p_1 - q_1| + |p_2 - q_2| + \dots + |p_n - q_n|$

3. Minkowski Distance:

- The Minkowski distance is a generalization of both the Euclidean and Manhattan distances.
- It is defined as:
- **Minkowski Distance** $= (\sum_{i=1}^n |p_i - q_i|^r)^{1/r}$
- When $r=1$, it reduces to the Manhattan distance, and when $r=2$, it becomes the Euclidean distance.

4. Cosine Similarity:

- Cosine similarity measures the cosine of the angle between two vectors in a multidimensional space.
- It is often used in text mining and recommendation systems.
- The formula for cosine similarity between two vectors A and B is:
Cosine Similarity $= \frac{A \cdot B}{\|A\| \|B\|}$

These distance metrics play a crucial role in determining the similarity or dissimilarity between objects in cluster analysis, helping to group similar objects together and separate dissimilar ones.

11 Calculate the Euclidean distance and Manhattan distance between points (2,-1) and (-2,2)

Ans.11) To calculate the Euclidean distance and Manhattan distance between points (2, -1) and (-2, 2), we use the following formulas:

1. Euclidean Distance:

- The Euclidean distance between two points (x1, y1) and (x2, y2) is given by:
- $Euclidean\ Distance = (x2 - x1)^2 + (y2 - y1)^2$

2. Manhattan Distance:

- The Manhattan distance between two points (x1, y1) and (x2, y2) is the sum of the absolute differences in their coordinates:
- $Manhattan\ Distance = |x2 - x1| + |y2 - y1|$

Given points (2, -1) and (-2, 2):

• For Euclidean Distance:

- $x1 = 2, y1 = -1$
- $x2 = -2, y2 = 2$
- Euclidean Distance
- $= \sqrt{(-2 - 2)^2 + (2 - (-1))^2}$
- $= \sqrt{(-4)^2 + (3)^2}$
- $= \sqrt{16 + 9}$
- $= \sqrt{25}$
- $= 5$

• For Manhattan Distance:

- $x1 = 2, y1 = -1$
- $x2 = -2, y2 = 2$
- Manhattan Distance
- $= |(-2 - 2)| + |2 - (-1)|$

- $=4+3=7$ $|(-2-2)|+|2-(-1)|$
- $=4+3=7$

Therefore, the Euclidean distance between points (2, -1) and (-2, 2) is 5 units, and the Manhattan distance between the same points is 7 units.