

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	ANF1	ANF2	p	test
n		510	115		
Subtype (%)	ANF1	510 (100.0)	0 (0.0)	<0.001	
	ANF2	0 (0.0)	115 (100.0)		
vital status (%)	Alive	134 (93.7)	29 (93.5)	1.000	exact
	Dead	9 (6.3)	2 (6.5)		
days to birth (median [IQR])		-21479.00 [-24882.00, -17949.50]	-20048.00 [-23620.50, -17726.00]	0.074	nonnorm
days to death (median [IQR])		1673.00 [616.00, 2417.00]	785.00 [395.00, 2271.00]	0.279	nonnorm
days to last followup (median [IQR])		757.00 [477.00, 1618.25]	730.00 [443.25, 1565.50]	0.816	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.2)	0 (0.0)	<0.001	exact
	ASIAN	27 (5.4)	5 (4.4)		
	BLACK OR AFRICAN AMERICAN	80 (16.0)	44 (38.9)		
	WHITE	391 (78.4)	64 (56.6)		
history of neoadjuvant treatment (%)	No	503 (98.8)	114 (100.0)	0.599	exact
	Yes	6 (1.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.25, 67.00]	54.00 [48.00, 64.00]	0.101	nonnorm
ethnicity (%)	HISPANIC OR LATINO	22 (4.8)	5 (4.7)	1.000	
	NOT HISPANIC OR LATINO	436 (95.2)	101 (95.3)		
primary lymph node presentation assessment (%)	NO	22 (5.4)	4 (4.3)	0.800	exact
	YES	385 (94.6)	89 (95.7)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.9)	<0.001	exact
	Infiltrating Ductal Carcinoma	341 (66.9)	95 (83.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ANF1	ANF2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	116 (22.7)	2 (1.8)	0.021	exact
	Medullary Carcinoma	1 (0.2)	4 (3.5)		
	Metaplastic Carcinoma	1 (0.2)	6 (5.3)		
	Mixed Histology (please specify)	21 (4.1)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
	Other, specify	20 (3.9)	6 (5.3)		
	Indeterminate (neither Pre or Postmenopausal)	9 (1.9)	5 (4.7)		
	Peri (6-12 months since last menstrual period)	13 (2.8)	7 (6.6)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	328 (70.1)	77 (72.6)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	118 (25.2)	17 (16.0)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.9)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	85 (17.8)	99 (91.7)	<0.001	
	Positive	392 (82.0)	8 (7.4)		
	Negative	36 (7.5)	94 (86.2)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	442 (92.5)	15 (13.8)	0.045	exact
	Equivocal	91 (22.0)	19 (20.2)		
	Indeterminate	7 (1.7)	3 (3.2)		
	Negative	250 (60.5)	66 (70.2)		
	Positive	65 (15.7)	6 (6.4)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.002	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ANF1	ANF2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	<0.001	nonnorm
er level cell percentage category (%)	<10%	14 (5.3)	21 (75.0)	<0.001	exact
	10-19%	19 (7.2)	1 (3.6)		
	20-29%	2 (0.8)	2 (7.1)		
	30-39%	4 (1.5)	1 (3.6)		
	40-49%	3 (1.1)	0 (0.0)		
	50-59%	8 (3.0)	0 (0.0)		
	60-69%	7 (2.7)	0 (0.0)		
	70-79%	40 (15.2)	0 (0.0)		
	80-89%	18 (6.8)	0 (0.0)		
	90-99%	149 (56.4)	3 (10.7)		
progesterone receptor level cell percent category (%)	<10%	55 (23.1)	24 (96.0)	<0.001	exact
	10-19%	18 (7.6)	0 (0.0)		
	20-29%	8 (3.4)	0 (0.0)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.6)	0 (0.0)		
	50-59%	10 (4.2)	0 (0.0)		
	60-69%	12 (5.0)	0 (0.0)		
	70-79%	29 (12.2)	0 (0.0)		
	80-89%	23 (9.7)	0 (0.0)		
	90-99%	64 (26.9)	1 (4.0)		
distant metastasis present ind2 (%)	NO	215 (97.7)	38 (97.4)	1.000	exact
	YES	5 (2.3)	1 (2.6)		
stage event pathologic stage (%)	Stage I	44 (8.7)	6 (5.3)	0.036	exact
	Stage IA	40 (7.9)	9 (7.9)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.9)		
	Stage IIA	147 (29.1)	55 (48.2)		
	Stage IIB	124 (24.5)	24 (21.1)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	89 (17.6)	11 (9.6)		
	Stage IIIB	12 (2.4)	2 (1.8)		
	Stage IIIC	33 (6.5)	3 (2.6)		
	Stage IV	5 (1.0)	2 (1.8)		
	Stage X	3 (0.6)	1 (0.9)		