

# Summary\_of\_clinical\_variables

## Contents

Table 1: Summarization of clinical variables stratified by subtype

|                                                             | level                               | RWR-NF1                             | RWR-NF2                             | p      | test    |
|-------------------------------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|--------|---------|
| n                                                           |                                     | 483                                 | 142                                 |        |         |
| Subtype (%)                                                 | RWR-NF1                             | 483 (100.0)                         | 0 ( 0.0)                            | <0.001 |         |
|                                                             | RWR-NF2                             | 0 ( 0.0)                            | 142 (100.0)                         |        |         |
| vital status (%)                                            | Alive                               | 130 ( 93.5)                         | 33 ( 94.3)                          | 1.000  | exact   |
|                                                             | Dead                                | 9 ( 6.5)                            | 2 ( 5.7)                            |        |         |
| days to birth<br>(median [IQR])                             |                                     | -21534.50 [-24810.50,<br>-17804.00] | -20347.50 [-24058.50,<br>-18073.25] | 0.253  | nonnorm |
| days to death<br>(median [IQR])                             |                                     | 1590.50 [606.25,<br>2430.00]        | 939.50 [473.50,<br>2357.50]         | 0.527  | nonnorm |
| days to last<br>followup<br>(median [IQR])                  |                                     | 765.00 [485.00,<br>1644.00]         | 701.00 [423.50,<br>1540.00]         | 0.398  | nonnorm |
| race list (%)                                               | AMERICAN INDIAN<br>OR ALASKA NATIVE | 0 ( 0.0)                            | 1 ( 0.7)                            | <0.001 | exact   |
|                                                             | ASIAN                               | 22 ( 4.7)                           | 10 ( 7.1)                           |        |         |
|                                                             | BLACK OR AFRICAN<br>AMERICAN        | 74 ( 15.7)                          | 50 ( 35.7)                          |        |         |
|                                                             | WHITE                               | 376 ( 79.7)                         | 79 ( 56.4)                          |        |         |
| history of<br>neoadjuvant<br>treatment (%)                  | No                                  | 476 ( 98.8)                         | 141 (100.0)                         | 0.346  | exact   |
|                                                             | Yes                                 | 6 ( 1.2)                            | 0 ( 0.0)                            |        |         |
| age at initial<br>pathologic<br>diagnosis<br>(median [IQR]) |                                     | 58.00 [48.00, 67.00]                | 55.00 [49.00, 65.00]                | 0.272  | nonnorm |
| ethnicity (%)                                               | HISPANIC OR<br>LATINO               | 20 ( 4.6)                           | 7 ( 5.4)                            | 0.897  |         |
|                                                             | NOT HISPANIC OR<br>LATINO           | 414 ( 95.4)                         | 123 ( 94.6)                         |        |         |
| primary lymph<br>node<br>presentation<br>assessment<br>(%)  | NO                                  | 18 ( 4.7)                           | 8 ( 6.8)                            | 0.501  |         |
|                                                             | YES                                 | 365 ( 95.3)                         | 109 ( 93.2)                         |        |         |
| histological<br>type (%)                                    | Infiltrating Carcinoma<br>NOS       | 0 ( 0.0)                            | 1 ( 0.7)                            | <0.001 | exact   |
|                                                             | Infiltrating Ductal<br>Carcinoma    | 320 ( 66.3)                         | 116 ( 82.3)                         |        |         |

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

|                                                             | level                                                                                        | RWR-NF1           | RWR-NF2           | p      | test    |
|-------------------------------------------------------------|----------------------------------------------------------------------------------------------|-------------------|-------------------|--------|---------|
| menopause status (%)                                        | Infiltrating Lobular Carcinoma                                                               | 116 ( 24.0)       | 2 ( 1.4)          | 0.014  | exact   |
|                                                             | Medullary Carcinoma                                                                          | 0 ( 0.0)          | 5 ( 3.5)          |        |         |
|                                                             | Metaplastic Carcinoma                                                                        | 1 ( 0.2)          | 6 ( 4.3)          |        |         |
|                                                             | Mixed Histology (please specify)                                                             | 20 ( 4.1)         | 1 ( 0.7)          |        |         |
|                                                             | Mucinous Carcinoma                                                                           | 10 ( 2.1)         | 0 ( 0.0)          |        |         |
|                                                             | Other, specify                                                                               | 16 ( 3.3)         | 10 ( 7.1)         |        |         |
|                                                             | Indeterminate (neither Pre or Postmenopausal)                                                | 9 ( 2.0)          | 5 ( 3.8)          |        |         |
|                                                             | Peri (6-12 months since last menstrual period)                                               | 12 ( 2.7)         | 8 ( 6.1)          |        |         |
|                                                             | Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)            | 307 ( 69.3)       | 98 ( 74.8)        |        |         |
|                                                             | Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement) | 115 ( 26.0)       | 20 ( 15.3)        |        |         |
| breast carcinoma progesterone receptor status (%)           | Indeterminate                                                                                | 1 ( 0.2)          | 1 ( 0.8)          | <0.001 | exact   |
| breast carcinoma estrogen receptor status (%)               | Negative                                                                                     | 70 ( 15.5)        | 114 ( 85.7)       | <0.001 |         |
|                                                             | Positive                                                                                     | 382 ( 84.3)       | 18 ( 13.5)        |        |         |
|                                                             | Negative                                                                                     | 25 ( 5.5)         | 105 ( 78.4)       |        |         |
| lab proc her2 neu immuno-histochemistry receptor status (%) | Positive                                                                                     | 428 ( 94.5)       | 29 ( 21.6)        | 0.671  | exact   |
|                                                             | Equivocal                                                                                    | 86 ( 21.9)        | 24 ( 20.9)        |        |         |
|                                                             | Indeterminate                                                                                | 7 ( 1.8)          | 3 ( 2.6)          |        |         |
|                                                             | Negative                                                                                     | 241 ( 61.5)       | 75 ( 65.2)        |        |         |
|                                                             | Positive                                                                                     | 58 ( 14.8)        | 13 ( 11.3)        |        |         |
| number of lymphnodes positive by ihc (median [IQR])         |                                                                                              | 0.00 [0.00, 0.00] | 0.00 [0.00, 0.00] | 0.014  | nonnorm |

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

|                                                       | level      | RWR-NF1           | RWR-NF2           | p      | test    |
|-------------------------------------------------------|------------|-------------------|-------------------|--------|---------|
| number of lymphnodes positive by he (median [IQR])    |            | 1.00 [0.00, 3.00] | 0.00 [0.00, 2.00] | <0.001 | nonnorm |
| er level cell percentage category (%)                 | <10%       | 8 ( 3.2)          | 27 ( 67.5)        | <0.001 | exact   |
|                                                       | 10-19%     | 19 ( 7.5)         | 1 ( 2.5)          |        |         |
|                                                       | 20-29%     | 2 ( 0.8)          | 2 ( 5.0)          |        |         |
|                                                       | 30-39%     | 3 ( 1.2)          | 2 ( 5.0)          |        |         |
|                                                       | 40-49%     | 3 ( 1.2)          | 0 ( 0.0)          |        |         |
|                                                       | 50-59%     | 8 ( 3.2)          | 0 ( 0.0)          |        |         |
|                                                       | 60-69%     | 7 ( 2.8)          | 0 ( 0.0)          |        |         |
|                                                       | 70-79%     | 37 ( 14.7)        | 3 ( 7.5)          |        |         |
|                                                       | 80-89%     | 18 ( 7.1)         | 0 ( 0.0)          |        |         |
|                                                       | 90-99%     | 147 ( 58.3)       | 5 ( 12.5)         |        |         |
| progesterone receptor level cell percent category (%) | <10%       | 48 ( 20.9)        | 31 ( 93.9)        | <0.001 | exact   |
|                                                       | 10-19%     | 18 ( 7.8)         | 0 ( 0.0)          |        |         |
|                                                       | 20-29%     | 8 ( 3.5)          | 0 ( 0.0)          |        |         |
|                                                       | 30-39%     | 8 ( 3.5)          | 0 ( 0.0)          |        |         |
|                                                       | 40-49%     | 11 ( 4.8)         | 0 ( 0.0)          |        |         |
|                                                       | 50-59%     | 10 ( 4.3)         | 0 ( 0.0)          |        |         |
|                                                       | 60-69%     | 11 ( 4.8)         | 1 ( 3.0)          |        |         |
|                                                       | 70-79%     | 29 ( 12.6)        | 0 ( 0.0)          |        |         |
|                                                       | 80-89%     | 23 ( 10.0)        | 0 ( 0.0)          |        |         |
|                                                       | 90-99%     | 64 ( 27.8)        | 1 ( 3.0)          |        |         |
| distant metastasis present ind2 (%)                   | NO         | 205 ( 97.6)       | 48 ( 98.0)        | 1.000  | exact   |
|                                                       | YES        | 5 ( 2.4)          | 1 ( 2.0)          |        |         |
| stage event pathologic stage (%)                      | Stage I    | 39 ( 8.1)         | 11 ( 7.8)         | 0.148  | exact   |
|                                                       | Stage IA   | 39 ( 8.1)         | 10 ( 7.1)         |        |         |
|                                                       | Stage IB   | 3 ( 0.6)          | 0 ( 0.0)          |        |         |
|                                                       | Stage II   | 4 ( 0.8)          | 1 ( 0.7)          |        |         |
|                                                       | Stage IIA  | 139 ( 29.0)       | 63 ( 44.7)        |        |         |
|                                                       | Stage IIB  | 117 ( 24.4)       | 31 ( 22.0)        |        |         |
|                                                       | Stage III  | 2 ( 0.4)          | 0 ( 0.0)          |        |         |
|                                                       | Stage IIIA | 85 ( 17.7)        | 15 ( 10.6)        |        |         |
|                                                       | Stage IIIB | 12 ( 2.5)         | 2 ( 1.4)          |        |         |
|                                                       | Stage IIIC | 31 ( 6.5)         | 5 ( 3.5)          |        |         |
|                                                       | Stage IV   | 5 ( 1.0)          | 2 ( 1.4)          |        |         |
|                                                       | Stage X    | 3 ( 0.6)          | 1 ( 0.7)          |        |         |