

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
n		80	143	172	85	145		
Subtype (%)	MSNE1	80 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	MSNE2	0 (0.0)	143 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	MSNE3	0 (0.0)	0 (0.0)	172 (100.0)	0 (0.0)	0 (0.0)		
	MSNE4	0 (0.0)	0 (0.0)	0 (0.0)	85 (100.0)	0 (0.0)		
	MSNE5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	145 (100.0)		
vital status (%)	Alive	18 (94.7)	42 (93.3)	42 (91.3)	22 (95.7)	39 (95.1)	0.977	exact
	Dead	1 (5.3)	3 (6.7)	4 (8.7)	1 (4.3)	2 (4.9)		
days to birth (median [IQR])		-20080.00 [- 24294.00, -18062.50]	-21685.00 [- 25558.00, -17702.00]	-22189.00 [- 25131.75, -18425.75]	-20587.00 [- 23408.50, -17988.50]	-20650.00 [- 23423.00, -17580.00]	0.300	nonnorm
days to death (median [IQR])		1363.00 [622.50, 2881.75]	1800.00 [647.75, 2445.00]	1541.00 [782.00, 2381.00]	524.00 [322.00, 785.00]	825.00 [593.50, 1660.00]		
days to last fol-lowup (median [IQR])		690.00 [443.25, 1390.00]	755.00 [477.00, 1613.75]	665.00 [463.75, 1406.50]	738.00 [449.50, 1607.50]	1013.00 [554.00, 1853.00]	0.217	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	0 (0.0)		
	ASIAN	4 (5.0)	6 (4.3)	12 (7.1)	4 (4.9)	6 (4.2)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	33 (41.2)	22 (15.7)	26 (15.5)	25 (30.5)	18 (12.7)		
	WHITE	43 (53.8)	112 (80.0)	129 (76.8)	53 (64.6)	118 (83.1)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
history of neoadjuvant treatment (%)	No	80 (100.0)	142 (100.0)	166 (96.5)	84 (100.0)	145 (100.0)	0.004	exact
	Yes	0 (0.0)	0 (0.0)	6 (3.5)	0 (0.0)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		54.00 [49.00, 66.00]	59.00 [47.50, 69.00]	60.00 [50.00, 68.00]	56.00 [49.00, 63.00]	56.00 [47.00, 64.00]	0.358	nonnorm
ethnicity (%)	HISPANIC OR LATINO	4 (5.5)	11 (8.5)	5 (3.3)	4 (5.1)	3 (2.3)	0.200	exact
	NOT HISPANIC OR LATINO	69 (94.5)	119 (91.5)	148 (96.7)	75 (94.9)	126 (97.7)		
primary lymph node presentation assessment (%)	NO	4 (5.9)	4 (3.8)	12 (8.8)	4 (6.0)	2 (1.6)	0.097	exact
	YES	64 (94.1)	100 (96.2)	124 (91.2)	63 (94.0)	123 (98.4)		
histologic type (%)	Infiltrating Carcinoma NOS	1 (1.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	64 (81.0)	103 (72.0)	130 (75.6)	61 (71.8)	78 (53.8)		
	Infiltrating Lobular Carcinoma	2 (2.5)	19 (13.3)	27 (15.7)	14 (16.5)	56 (38.6)		
	Medullary Carcinoma	5 (6.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	4 (5.1)	0 (0.0)	0 (0.0)	3 (3.5)	0 (0.0)		
	Mixed Histology (please specify)	0 (0.0)	5 (3.5)	10 (5.8)	0 (0.0)	6 (4.1)		
	Mucinous Carcinoma	0 (0.0)	7 (4.9)	1 (0.6)	1 (1.2)	1 (0.7)		
	Other, specify	3 (3.8)	9 (6.3)	4 (2.3)	6 (7.1)	4 (2.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
menopausal status (%)	Indeterm (neither Pre or Post-menopausal)	2 (2.7)	7 (5.3)	2 (1.3)	2 (2.4)	1 (0.8)	0.104	exact
	Peri (6-12 months since last menstrual period)	6 (8.2)	3 (2.3)	3 (1.9)	1 (1.2)	7 (5.4)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	51 (69.9)	90 (67.7)	118 (75.6)	61 (74.4)	85 (65.4)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	14 (19.2)	33 (24.8)	33 (21.2)	18 (22.0)	37 (28.5)		
	Indeterm	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	1 (0.7)		
breast carcinoma progesterone receptor status (%)	Negative	64 (86.5)	17 (12.6)	32 (20.3)	52 (65.0)	19 (13.7)	<0.001	exact
	Positive	10 (13.5)	118 (87.4)	126 (79.7)	27 (33.8)	119 (85.6)		
	Indeterm	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	1 (0.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
breast carci- noma estro- gen recep- tor status (%)	Negative	56 (75.7)	3 (2.2)	19 (12.0)	46 (56.8)	6 (4.3)	<0.001	
	Positive	18 (24.3)	132 (97.8)	139 (88.0)	35 (43.2)	133 (95.7)		
lab proc her2 neu im- muno- histo- chem- istry recep- tor status (%)	Equivocal	15 (23.4)	28 (24.8)	33 (23.4)	9 (12.9)	25 (21.0)	0.001	exact
	Indeterm	2 (3.1)	3 (2.7)	3 (2.1)	1 (1.4)	1 (0.8)		
	Negative	41 (64.1)	73 (64.6)	69 (48.9)	52 (74.3)	81 (68.1)		
	Positive	6 (9.4)	9 (8.0)	36 (25.5)	8 (11.4)	12 (10.1)		
number of lym- phn- odes posi- tive by ihc (me- dian [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.007	nonnorm
number of lym- phn- odes posi- tive by he (me- dian [IQR])		0.00 [0.00, 2.00]	1.00 [0.00, 3.00]	1.00 [0.00, 4.00]	0.00 [0.00, 2.00]	1.00 [0.00, 2.00]	0.061	nonnorm
er level cell per- cent- age cate- gory (%)	<10%	13 (59.1)	1 (1.4)	5 (6.4)	14 (34.1)	2 (2.6)	<0.001	exact
	10- 19%	2 (9.1)	4 (5.5)	8 (10.3)	4 (9.8)	2 (2.6)		
	20- 29%	2 (9.1)	0 (0.0)	2 (2.6)	0 (0.0)	0 (0.0)		
	30- 39%	2 (9.1)	1 (1.4)	1 (1.3)	0 (0.0)	1 (1.3)		
	40- 49%	0 (0.0)	2 (2.7)	1 (1.3)	0 (0.0)	0 (0.0)		
	50- 59%	1 (4.5)	3 (4.1)	1 (1.3)	0 (0.0)	3 (3.8)		
	60- 69%	0 (0.0)	2 (2.7)	2 (2.6)	0 (0.0)	3 (3.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
progesterone receptor level cell percent category (%)	70-79%	0 (0.0)	13 (17.8)	12 (15.4)	6 (14.6)	9 (11.5)	<0.001	exact
	80-89%	0 (0.0)	4 (5.5)	8 (10.3)	2 (4.9)	4 (5.1)		
	90-99%	2 (9.1)	43 (58.9)	38 (48.7)	15 (36.6)	54 (69.2)		
	10-19%	0 (0.0)	6 (8.7)	7 (10.8)	0 (0.0)	5 (6.3)		
	20-29%	0 (0.0)	0 (0.0)	4 (6.2)	0 (0.0)	4 (5.1)		
	30-39%	0 (0.0)	3 (4.3)	2 (3.1)	0 (0.0)	3 (3.8)		
	40-49%	0 (0.0)	5 (7.2)	3 (4.6)	0 (0.0)	3 (3.8)		
	50-59%	1 (5.6)	4 (5.8)	1 (1.5)	0 (0.0)	4 (5.1)		
	60-69%	0 (0.0)	3 (4.3)	3 (4.6)	2 (6.2)	4 (5.1)		
	70-79%	0 (0.0)	10 (14.5)	6 (9.2)	2 (6.2)	11 (13.9)		
distant metastasis present ind2 (%)	80-89%	0 (0.0)	3 (4.3)	8 (12.3)	3 (9.4)	9 (11.4)	0.029	exact
	90-99%	0 (0.0)	22 (31.9)	13 (20.0)	7 (21.9)	23 (29.1)		
	NO	30 (100.0)	57 (100.0)	72 (100.0)	30 (93.8)	64 (94.1)		
	YES	0 (0.0)	0 (0.0)	0 (0.0)	2 (6.2)	4 (5.9)		
	Stage I	5 (6.3)	13 (9.2)	12 (7.0)	6 (7.1)	14 (9.7)		
	Stage IA	4 (5.1)	18 (12.7)	8 (4.7)	7 (8.3)	12 (8.3)		
	Stage IB	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	2 (1.4)		
	Stage II	1 (1.3)	1 (0.7)	1 (0.6)	0 (0.0)	2 (1.4)		
	Stage IIA	37 (46.8)	35 (24.6)	56 (32.7)	35 (41.7)	39 (27.1)		
	Stage IIB	16 (20.3)	37 (26.1)	37 (21.6)	18 (21.4)	40 (27.8)		
stage event pathologic stage (%)	Stage III	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	1 (0.7)	0.212	exact
	Stage IIIA	9 (11.4)	23 (16.2)	33 (19.3)	11 (13.1)	24 (16.7)		
	Stage IIIB	1 (1.3)	5 (3.5)	6 (3.5)	1 (1.2)	1 (0.7)		
	Stage IIIC	3 (3.8)	6 (4.2)	15 (8.8)	4 (4.8)	8 (5.6)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
Stage IV	1 (1.3)	2 (1.4)	2 (1.2)	1 (1.2)	1 (0.7)		
Stage X	2 (2.5)	2 (1.4)	0 (0.0)	0 (0.0)	0 (0.0)		