

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MC1	MC2	MC3	MC4	p	test
n		182	308	64	71		
Subtype (%)	MC1	182 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	MC2	0 (0.0)	308 (100.0)	0 (0.0)	0 (0.0)		
	MC3	0 (0.0)	0 (0.0)	64 (100.0)	0 (0.0)		
	MC4	0 (0.0)	0 (0.0)	0 (0.0)	71 (100.0)		
vital status (%)	Alive	44 (91.7)	86 (94.5)	17 (94.4)	16 (94.1)	0.927	exact
	Dead	4 (8.3)	5 (5.5)	1 (5.6)	1 (5.9)		
days to birth (median [IQR])		-22179.00 [- 25127.50, -18519.00]	-21291.00 [- 24318.00, -17627.00]	-20249.00 [- 23873.00, -17743.50]	-20048.00 [- 24187.00, -17902.00]	0.382	nonnorm
days to death (median [IQR])		1910.50 [782.00, 2517.50]	1508.00 [577.00, 2373.00]	570.00 [372.50, 742.75]	1688.00 [606.00, 2909.50]	0.141	nonnorm
days to last followup (median [IQR])		662.00 [454.75, 1381.00]	860.00 [504.00, 1688.00]	738.00 [437.50, 1676.00]	767.00 [443.25, 1604.00]	0.254	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	ASIAN	13 (7.3)	12 (4.0)	4 (6.5)	3 (4.2)		
	BLACK OR AFRICAN AMERICAN	29 (16.3)	46 (15.3)	19 (30.6)	30 (42.3)		
	WHITE	135 (75.8)	243 (80.7)	39 (62.9)	38 (53.5)		
history of neoadjuvant treatment (%)	No	176 (96.7)	307 (100.0)	63 (100.0)	71 (100.0)	0.002	exact
	Yes	6 (3.3)	0 (0.0)	0 (0.0)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		59.50 [50.00, 68.00]	58.00 [47.75, 66.00]	55.00 [48.00, 65.00]	54.00 [48.50, 66.00]	0.482	nonnorm
ethnicity (%)	HISPANIC OR LATINO	6 (3.7)	15 (5.4)	2 (3.3)	4 (6.2)	0.789	exact
	NOT HISPANIC OR LATINO	156 (96.3)	262 (94.6)	58 (96.7)	61 (93.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MC1	MC2	MC3	MC4	p	test
primary lymph node presentation assessment (%)	NO	12 (8.3)	8 (3.3)	3 (5.6)	3 (5.1)	0.170	exact
	YES	132 (91.7)	235 (96.7)	51 (94.4)	56 (94.9)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.4)	<0.001	exact
	Infiltrating Ductal Carcinoma	139 (76.4)	195 (63.3)	46 (71.9)	56 (80.0)		
	Infiltrating Lobular Carcinoma	28 (15.4)	80 (26.0)	9 (14.1)	1 (1.4)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	5 (7.1)		
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	3 (4.7)	4 (5.7)		
	Mixed Histology (please specify)	10 (5.5)	11 (3.6)	0 (0.0)	0 (0.0)		
	Mucinous Carcinoma	1 (0.5)	9 (2.9)	0 (0.0)	0 (0.0)		
	Other, specify	4 (2.2)	13 (4.2)	6 (9.4)	3 (4.3)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	2 (1.2)	8 (2.8)	2 (3.2)	2 (3.2)	0.114	exact
	Peri (6-12 months since last menstrual period)	3 (1.8)	9 (3.2)	2 (3.2)	6 (9.5)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	125 (75.3)	189 (66.8)	48 (77.4)	43 (68.3)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	36 (21.7)	77 (27.2)	10 (16.1)	12 (19.0)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	1 (0.3)	1 (1.7)	0 (0.0)	<0.001	exact
	Negative	36 (21.6)	41 (13.9)	46 (78.0)	61 (93.8)		
	Positive	131 (78.4)	253 (85.8)	12 (20.3)	4 (6.2)		
breast carcinoma estrogen receptor status (%)	Negative	21 (12.6)	10 (3.4)	44 (73.3)	55 (84.6)	<0.001	
	Positive	146 (87.4)	285 (96.6)	16 (26.7)	10 (15.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	33 (22.1)	53 (21.0)	9 (18.0)	15 (26.8)	<0.001	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MC1	MC2	MC3	MC4	p	test
	Indeterminate	4 (2.7)	4 (1.6)	0 (0.0)	2 (3.6)		
	Negative	73 (49.0)	171 (67.9)	35 (70.0)	37 (66.1)		
	Positive	39 (26.2)	24 (9.5)	6 (12.0)	2 (3.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.035	nonnorm
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 4.00]	1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.003	nonnorm
er level cell percentage category (%)	<10%	6 (7.1)	6 (3.6)	10 (45.5)	13 (76.5)	<0.001	exact
	10-19%	9 (10.7)	8 (4.7)	2 (9.1)	1 (5.9)		
	20-29%	2 (2.4)	0 (0.0)	0 (0.0)	2 (11.8)		
	30-39%	2 (2.4)	2 (1.2)	0 (0.0)	1 (5.9)		
	40-49%	1 (1.2)	2 (1.2)	0 (0.0)	0 (0.0)		
	50-59%	2 (2.4)	6 (3.6)	0 (0.0)	0 (0.0)		
	60-69%	2 (2.4)	4 (2.4)	1 (4.5)	0 (0.0)		
	70-79%	12 (14.3)	24 (14.2)	4 (18.2)	0 (0.0)		
	80-89%	8 (9.5)	10 (5.9)	0 (0.0)	0 (0.0)		
	90-99%	40 (47.6)	107 (63.3)	5 (22.7)	0 (0.0)		
progesterone receptor level cell percent category (%)	<10%	21 (30.4)	31 (19.4)	12 (63.2)	15 (100.0)	<0.001	exact
	10-19%	7 (10.1)	11 (6.9)	0 (0.0)	0 (0.0)		
	20-29%	4 (5.8)	3 (1.9)	1 (5.3)	0 (0.0)		
	30-39%	2 (2.9)	6 (3.8)	0 (0.0)	0 (0.0)		
	40-49%	3 (4.3)	8 (5.0)	0 (0.0)	0 (0.0)		
	50-59%	2 (2.9)	8 (5.0)	0 (0.0)	0 (0.0)		
	60-69%	3 (4.3)	9 (5.6)	0 (0.0)	0 (0.0)		
	70-79%	6 (8.7)	20 (12.5)	3 (15.8)	0 (0.0)		
	80-89%	8 (11.6)	14 (8.8)	1 (5.3)	0 (0.0)		
	90-99%	13 (18.8)	50 (31.2)	2 (10.5)	0 (0.0)		
distant metastasis present ind2 (%)	NO	77 (100.0)	133 (97.8)	18 (85.7)	25 (100.0)	0.019	exact
	YES	0 (0.0)	3 (2.2)	3 (14.3)	0 (0.0)		
stage event pathologic stage (%)	Stage I	11 (6.1)	29 (9.5)	5 (7.8)	5 (7.1)	0.072	exact
	Stage IA	9 (5.0)	31 (10.2)	5 (7.8)	4 (5.7)		
	Stage IB	0 (0.0)	3 (1.0)	0 (0.0)	0 (0.0)		
	Stage II	1 (0.6)	3 (1.0)	0 (0.0)	1 (1.4)		
	Stage IIA	60 (33.1)	80 (26.2)	29 (45.3)	33 (47.1)		
	Stage IIB	38 (21.0)	81 (26.6)	14 (21.9)	15 (21.4)		
	Stage III	1 (0.6)	0 (0.0)	1 (1.6)	0 (0.0)		
	Stage IIIA	36 (19.9)	52 (17.0)	6 (9.4)	6 (8.6)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	MC1	MC2	MC3	MC4	p	test
Stage IIIB	6 (3.3)	6 (2.0)	0 (0.0)	2 (2.9)		
Stage IIIC	16 (8.8)	15 (4.9)	3 (4.7)	2 (2.9)		
Stage IV	2 (1.1)	3 (1.0)	1 (1.6)	1 (1.4)		
Stage X	1 (0.6)	2 (0.7)	0 (0.0)	1 (1.4)		