

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MDICC1	MDICC2	p	test
n		607	18		
Subtype (%)	MDICC1	607 (100.0)	0 (0.0)	<0.001	exact
	MDICC2	0 (0.0)	18 (100.0)		
vital status (%)	Alive	162 (93.6)	1 (100.0)	1.000	exact
	Dead	11 (6.4)	0 (0.0)		
days to birth (median [IQR])		-21343.50 [-24746.00, -17854.00]	-19973.50 [-22908.00, -18204.50]	0.495	nonnorm
days to death (median [IQR])		1388.00 [575.00, 2443.00]	322.00 [322.00, 322.00]	0.228	nonnorm
days to last followup (median [IQR])		766.00 [477.00, 1626.25]	621.00 [424.75, 829.50]	0.168	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.2)	0 (0.0)	0.001	exact
	ASIAN	31 (5.2)	1 (5.6)		
	BLACK OR AFRICAN AMERICAN	113 (19.0)	11 (61.1)		
	WHITE	449 (75.6)	6 (33.3)		
history of neoadjuvant treatment (%)	No	600 (99.2)	17 (94.4)	0.162	exact
	Yes	5 (0.8)	1 (5.6)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 67.00]	54.50 [49.50, 62.50]	0.570	nonnorm
ethnicity (%)	HISPANIC OR LATINO	27 (4.9)	0 (0.0)	1.000	exact
	NOT HISPANIC OR LATINO	523 (95.1)	14 (100.0)		
primary lymph node presentation assessment (%)	NO	25 (5.2)	1 (6.7)	0.556	exact
	YES	460 (94.8)	14 (93.3)		
histological type (%)	Infiltrating Carcinoma NOS	1 (0.2)	0 (0.0)	0.133	exact
	Infiltrating Ductal Carcinoma	420 (69.3)	16 (88.9)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MDICC1	MDICC2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	118 (19.5)	0 (0.0)	0.027	exact
	Medullary Carcinoma	5 (0.8)	0 (0.0)		
	Metaplastic Carcinoma	6 (1.0)	1 (5.6)		
	Mixed Histology (please specify)	21 (3.5)	0 (0.0)		
	Mucinous Carcinoma	10 (1.7)	0 (0.0)		
	Other, specify	25 (4.1)	1 (5.6)		
	Indeterminate (neither Pre or Postmenopausal)	14 (2.5)	0 (0.0)		
	Peri (6-12 months since last menstrual period)	18 (3.2)	2 (12.5)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	391 (70.1)	14 (87.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	135 (24.2)	0 (0.0)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.4)	0 (0.0)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	169 (29.8)	15 (83.3)	<0.001	exact
	Positive	397 (69.9)	3 (16.7)		
	Negative	118 (20.7)	12 (66.7)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	451 (79.3)	6 (33.3)	1.000	exact
	Equivocal	107 (21.7)	3 (21.4)		
	Indeterminate	10 (2.0)	0 (0.0)		
	Negative	307 (62.3)	9 (64.3)		
	Positive	69 (14.0)	2 (14.3)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.353	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MDICC1	MDICC2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.238	nonnorm
er level cell percentage category (%)	<10%	33 (11.5)	2 (50.0)	0.021	exact
	10-19%	19 (6.6)	1 (25.0)		
	20-29%	4 (1.4)	0 (0.0)		
	30-39%	4 (1.4)	1 (25.0)		
	40-49%	3 (1.0)	0 (0.0)		
	50-59%	8 (2.8)	0 (0.0)		
	60-69%	7 (2.4)	0 (0.0)		
	70-79%	40 (13.9)	0 (0.0)		
	80-89%	18 (6.2)	0 (0.0)		
	90-99%	152 (52.8)	0 (0.0)		
progesterone receptor level cell percent category (%)	<10%	77 (29.5)	2 (100.0)	0.847	exact
	10-19%	18 (6.9)	0 (0.0)		
	20-29%	8 (3.1)	0 (0.0)		
	30-39%	8 (3.1)	0 (0.0)		
	40-49%	11 (4.2)	0 (0.0)		
	50-59%	10 (3.8)	0 (0.0)		
	60-69%	12 (4.6)	0 (0.0)		
	70-79%	29 (11.1)	0 (0.0)		
	80-89%	23 (8.8)	0 (0.0)		
	90-99%	65 (24.9)	0 (0.0)		
distant metastasis present ind2 (%)	NO	251 (97.7)	2 (100.0)	1.000	exact
	YES	6 (2.3)	0 (0.0)		
stage event pathologic stage (%)	Stage I	48 (8.0)	2 (11.1)	0.746	exact
	Stage IA	46 (7.6)	3 (16.7)		
	Stage IB	3 (0.5)	0 (0.0)		
	Stage II	5 (0.8)	0 (0.0)		
	Stage IIA	194 (32.2)	8 (44.4)		
	Stage IIB	144 (23.9)	4 (22.2)		
	Stage III	2 (0.3)	0 (0.0)		
	Stage IIIA	99 (16.4)	1 (5.6)		
	Stage IIIB	14 (2.3)	0 (0.0)		
	Stage IIIC	36 (6.0)	0 (0.0)		
	Stage IV	7 (1.2)	0 (0.0)		
	Stage X	4 (0.7)	0 (0.0)		