

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CC1	CC2	p	test
n		495	130		
Subtype (%)	CC1	495 (100.0)	0 (0.0)	<0.001	
	CC2	0 (0.0)	130 (100.0)		
vital status (%)	Alive	131 (93.6)	32 (94.1)	1.000	exact
	Dead	9 (6.4)	2 (5.9)		
days to birth (median [IQR])		-21534.50 [-24840.50, -17959.25]	-20086.00 [-23942.75, -17727.00]	0.104	nonnorm
days to death (median [IQR])		1508.00 [616.00, 2417.00]	912.00 [423.00, 2523.00]	0.550	nonnorm
days to last followup (median [IQR])		754.50 [477.00, 1611.00]	779.00 [447.75, 1616.75]	0.972	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.8)	<0.001	exact
	ASIAN	26 (5.4)	6 (4.7)		
	BLACK OR AFRICAN AMERICAN	79 (16.3)	45 (35.2)		
	WHITE	379 (78.3)	76 (59.4)		
history of neoadjuvant treatment (%)	No	488 (98.8)	129 (100.0)	0.354	exact
	Yes	6 (1.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.50, 67.00]	54.50 [48.00, 65.00]	0.140	nonnorm
ethnicity (%)	HISPANIC OR LATINO	21 (4.7)	6 (5.0)	1.000	
	NOT HISPANIC OR LATINO	424 (95.3)	113 (95.0)		
primary lymph node presentation assessment (%)	NO	21 (5.3)	5 (4.8)	1.000	
	YES	374 (94.7)	100 (95.2)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.8)	<0.001	exact
	Infiltrating Ductal Carcinoma	330 (66.7)	106 (82.2)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CC1	CC2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	116 (23.4)	2 (1.6)	0.039	exact
	Medullary Carcinoma	0 (0.0)	5 (3.9)		
	Metaplastic Carcinoma	1 (0.2)	6 (4.7)		
	Mixed Histology (please specify)	21 (4.2)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
	Other, specify	17 (3.4)	9 (7.0)		
	Indeterminate (neither Pre or Postmenopausal)	9 (2.0)	5 (4.2)		
	Peri (6-12 months since last menstrual period)	13 (2.9)	7 (5.9)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	318 (69.7)	87 (73.7)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	116 (25.4)	19 (16.1)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.8)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	74 (15.9)	110 (90.2)	<0.001	
	Positive	389 (83.8)	11 (9.0)		
	Negative	27 (5.8)	103 (83.7)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	437 (94.2)	20 (16.3)	0.386	exact
	Equivocal	88 (21.9)	22 (21.0)		
	Indeterminate	7 (1.7)	3 (2.9)		
	Negative	246 (61.2)	70 (66.7)		
	Positive	61 (15.2)	10 (9.5)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.006	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CC1	CC2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	<0.001	nonnorm
er level cell percentage category (%)	<10%	10 (3.9)	25 (75.8)	<0.001	exact
	10-19%	19 (7.3)	1 (3.0)		
	20-29%	2 (0.8)	2 (6.1)		
	30-39%	4 (1.5)	1 (3.0)		
	40-49%	3 (1.2)	0 (0.0)		
	50-59%	8 (3.1)	0 (0.0)		
	60-69%	7 (2.7)	0 (0.0)		
	70-79%	39 (15.1)	1 (3.0)		
	80-89%	18 (6.9)	0 (0.0)		
	90-99%	149 (57.5)	3 (9.1)		
progesterone receptor level cell percent category (%)	<10%	51 (21.8)	28 (96.6)	<0.001	exact
	10-19%	18 (7.7)	0 (0.0)		
	20-29%	8 (3.4)	0 (0.0)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.7)	0 (0.0)		
	50-59%	10 (4.3)	0 (0.0)		
	60-69%	11 (4.7)	1 (3.4)		
	70-79%	29 (12.4)	0 (0.0)		
	80-89%	23 (9.8)	0 (0.0)		
	90-99%	65 (27.8)	0 (0.0)		
distant metastasis present ind2 (%)	NO	209 (97.7)	44 (97.8)	1.000	exact
	YES	5 (2.3)	1 (2.2)		
stage event pathologic stage (%)	Stage I	40 (8.1)	10 (7.8)	0.029	exact
	Stage IA	40 (8.1)	9 (7.0)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.8)		
	Stage IIA	140 (28.5)	62 (48.1)		
	Stage IIB	124 (25.3)	24 (18.6)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	86 (17.5)	14 (10.9)		
	Stage IIIB	12 (2.4)	2 (1.6)		
	Stage IIIC	32 (6.5)	4 (3.1)		
	Stage IV	5 (1.0)	2 (1.6)		
	Stage X	3 (0.6)	1 (0.8)		