

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MC1	MC2	MC3	MC4	p	test
n		179	310	63	73		
Subtype (%)	MC1	179 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	MC2	0 (0.0)	310 (100.0)	0 (0.0)	0 (0.0)		
	MC3	0 (0.0)	0 (0.0)	63 (100.0)	0 (0.0)		
	MC4	0 (0.0)	0 (0.0)	0 (0.0)	73 (100.0)		
vital status (%)	Alive	42 (91.3)	87 (94.6)	17 (94.4)	17 (94.4)	0.925	exact
	Dead	4 (8.7)	5 (5.4)	1 (5.6)	1 (5.6)		
days to birth (median [IQR])		-22189.00 [- 25149.50, -18475.25]	-21291.00 [- 24324.00, -17648.50]	-20210.00 [- 23681.00, -17713.00]	-20048.00 [- 24139.00, -18041.00]	0.330	nonnorm
days to death (median [IQR])		1910.50 [782.00, 2517.50]	1508.00 [577.00, 2373.00]	570.00 [372.50, 742.75]	1688.00 [606.00, 2909.50]	0.141	nonnorm
days to last followup (median [IQR])		659.00 [430.00, 1270.00]	862.50 [505.25, 1691.00]	738.00 [437.50, 1676.00]	839.00 [447.75, 1658.75]	0.149	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	ASIAN	13 (7.4)	12 (4.0)	4 (6.6)	3 (4.1)		
	BLACK OR AFRICAN AMERICAN	29 (16.6)	47 (15.5)	18 (29.5)	30 (41.1)		
	WHITE	132 (75.4)	244 (80.5)	39 (63.9)	40 (54.8)		
history of neoadjuvant treatment (%)	No	173 (96.6)	309 (100.0)	62 (100.0)	73 (100.0)	0.002	exact
	Yes	6 (3.4)	0 (0.0)	0 (0.0)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		60.00 [50.00, 68.00]	58.00 [48.00, 66.00]	55.00 [48.00, 64.50]	54.00 [49.00, 66.00]	0.430	nonnorm
ethnicity (%)	HISPANIC OR LATINO	6 (3.8)	15 (5.4)	2 (3.4)	4 (6.0)	0.817	exact
	NOT HISPANIC OR LATINO	153 (96.2)	264 (94.6)	57 (96.6)	63 (94.0)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MC1	MC2	MC3	MC4	p	test
primary lymph node presentation assessment (%)	NO	12 (8.5)	8 (3.3)	3 (5.7)	3 (4.9)	0.145	exact
	YES	129 (91.5)	237 (96.7)	50 (94.3)	58 (95.1)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.4)	<0.001	exact
	Infiltrating Ductal Carcinoma	137 (76.5)	196 (63.2)	45 (71.4)	58 (80.6)		
	Infiltrating Lobular Carcinoma	27 (15.1)	81 (26.1)	9 (14.3)	1 (1.4)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	5 (6.9)		
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	3 (4.8)	4 (5.6)		
	Mixed Histology (please specify)	10 (5.6)	11 (3.5)	0 (0.0)	0 (0.0)		
	Mucinous Carcinoma	1 (0.6)	9 (2.9)	0 (0.0)	0 (0.0)		
	Other, specify	4 (2.2)	13 (4.2)	6 (9.5)	3 (4.2)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	2 (1.2)	8 (2.8)	2 (3.3)	2 (3.1)	0.121	exact
	Peri (6-12 months since last menstrual period)	3 (1.8)	9 (3.2)	2 (3.3)	6 (9.2)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	123 (75.5)	190 (66.7)	47 (77.0)	45 (69.2)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	35 (21.5)	78 (27.4)	10 (16.4)	12 (18.5)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	1 (0.3)	1 (1.7)	0 (0.0)	<0.001	exact
	Negative	34 (20.7)	42 (14.1)	45 (77.6)	63 (94.0)		
	Positive	130 (79.3)	254 (85.5)	12 (20.7)	4 (6.0)		
breast carcinoma estrogen receptor status (%)	Negative	19 (11.6)	11 (3.7)	43 (72.9)	57 (85.1)	<0.001	
	Positive	145 (88.4)	286 (96.3)	16 (27.1)	10 (14.9)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	31 (21.2)	53 (20.9)	9 (18.0)	17 (29.3)	<0.001	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MC1	MC2	MC3	MC4	p	test
	Indeterminate	4 (2.7)	4 (1.6)	0 (0.0)	2 (3.4)		
	Negative	72 (49.3)	172 (68.0)	35 (70.0)	37 (63.8)		
	Positive	39 (26.7)	24 (9.5)	6 (12.0)	2 (3.4)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.038	nonnorm
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 4.00]	1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.001	nonnorm
er level cell percentage category (%)	<10%	5 (6.1)	6 (3.5)	10 (45.5)	14 (77.8)	<0.001	exact
	10-19%	9 (11.0)	8 (4.7)	2 (9.1)	1 (5.6)		
	20-29%	2 (2.4)	0 (0.0)	0 (0.0)	2 (11.1)		
	30-39%	2 (2.4)	2 (1.2)	0 (0.0)	1 (5.6)		
	40-49%	1 (1.2)	2 (1.2)	0 (0.0)	0 (0.0)		
	50-59%	1 (1.2)	7 (4.1)	0 (0.0)	0 (0.0)		
	60-69%	2 (2.4)	4 (2.4)	1 (4.5)	0 (0.0)		
	70-79%	12 (14.6)	24 (14.1)	4 (18.2)	0 (0.0)		
	80-89%	8 (9.8)	10 (5.9)	0 (0.0)	0 (0.0)		
	90-99%	40 (48.8)	107 (62.9)	5 (22.7)	0 (0.0)		
progesterone receptor level cell percent category (%)	<10%	20 (29.9)	31 (19.3)	12 (63.2)	16 (100.0)	<0.001	exact
	10-19%	7 (10.4)	11 (6.8)	0 (0.0)	0 (0.0)		
	20-29%	4 (6.0)	3 (1.9)	1 (5.3)	0 (0.0)		
	30-39%	2 (3.0)	6 (3.7)	0 (0.0)	0 (0.0)		
	40-49%	3 (4.5)	8 (5.0)	0 (0.0)	0 (0.0)		
	50-59%	1 (1.5)	9 (5.6)	0 (0.0)	0 (0.0)		
	60-69%	3 (4.5)	9 (5.6)	0 (0.0)	0 (0.0)		
	70-79%	6 (9.0)	20 (12.4)	3 (15.8)	0 (0.0)		
	80-89%	8 (11.9)	14 (8.7)	1 (5.3)	0 (0.0)		
	90-99%	13 (19.4)	50 (31.1)	2 (10.5)	0 (0.0)		
distant metastasis present ind2 (%)	NO	76 (100.0)	133 (97.8)	18 (85.7)	26 (100.0)	0.021	exact
	YES	0 (0.0)	3 (2.2)	3 (14.3)	0 (0.0)		
stage event pathologic stage (%)	Stage I	10 (5.6)	29 (9.4)	5 (7.9)	6 (8.3)	0.056	exact
	Stage IA	9 (5.1)	31 (10.1)	5 (7.9)	4 (5.6)		
	Stage IB	0 (0.0)	3 (1.0)	0 (0.0)	0 (0.0)		
	Stage II	1 (0.6)	3 (1.0)	0 (0.0)	1 (1.4)		
	Stage IIA	59 (33.1)	81 (26.4)	28 (44.4)	34 (47.2)		
	Stage IIB	38 (21.3)	81 (26.4)	14 (22.2)	15 (20.8)		
	Stage III	1 (0.6)	0 (0.0)	1 (1.6)	0 (0.0)		
	Stage IIIA	36 (20.2)	52 (16.9)	6 (9.5)	6 (8.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	MC1	MC2	MC3	MC4	p	test
Stage IIIB	6 (3.4)	6 (2.0)	0 (0.0)	2 (2.8)		
Stage IIIC	16 (9.0)	15 (4.9)	3 (4.8)	2 (2.8)		
Stage IV	2 (1.1)	3 (1.0)	1 (1.6)	1 (1.4)		
Stage X	0 (0.0)	3 (1.0)	0 (0.0)	1 (1.4)		