

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
n		56	33	83	92	49	45	76	191		
Subtype (%)	wMKL1	56 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	wMKL2	0 (0.0)	33 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL3	0 (0.0)	0 (0.0)	83 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL4	0 (0.0)	0 (0.0)	0 (0.0)	92 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	49 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	45 (100.0)	0 (0.0)	0 (0.0)		
	wMKL7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	76 (100.0)	0 (0.0)		
	wMKL8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	191 (100.0)		
vital status (%)	Alive	13 (92.9)	10 (90.9)	23 (95.8)	19 (90.5)	10 (83.3)	8 (88.9)	20 (95.2)	60 (96.8)	0.451	exact
	Dead	1 (7.1)	1 (9.1)	1 (4.2)	2 (9.5)	2 (16.7)	1 (11.1)	1 (4.8)	2 (3.2)		
days to birth (median [IQR])		-20374.00 [-23905.50, -17583.00]	-22809.00 [-25082.50, -18320.50]	-21309.00 [-25005.00, -18158.50]	-20495.50 [-24163.00, -18387.25]	-21343.00 [-24660.00, -18152.00]	-22239.50 [-25759.00, -18412.75]	-21767.50 [-25540.75, -17834.75]	-20836.50 [-23480.50, -17629.25]	0.815	nonnorm
days to death (median [IQR])		NA [NA, NA]	2535.50 [2404.25, 2666.75]	1448.00 [702.00, 2248.25]	1688.00 [770.00, 2702.50]	538.00 [269.50, 1529.00]	1745.50 [1055.25, 2435.75]	1642.50 [763.00, 2485.25]	822.50 [535.75, 1777.25]	0.551	nonnorm
days to last followup (median [IQR])		974.50 [429.00, 1757.75]	594.00 [350.00, 1270.00]	666.00 [463.00, 1417.00]	701.00 [477.00, 1631.00]	809.50 [508.00, 1920.00]	728.00 [476.00, 1224.00]	1008.00 [523.75, 1894.25]	802.00 [502.50, 1565.50]	0.601	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0.002	exact
	ASIAN BLACK OR AFRICAN AMERICAN	2 (3.6) 7 (12.7)	4 (12.9) 4 (12.9)	6 (7.3) 19 (23.2)	5 (5.4) 34 (37.0)	5 (10.9) 8 (17.4)	1 (2.3) 4 (9.1)	2 (2.7) 16 (21.3)	7 (3.7) 32 (17.1)		
	WHITE	46 (83.6)	23 (74.2)	57 (69.5)	52 (56.5)	33 (71.7)	39 (88.6)	57 (76.0)	148 (79.1)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
history of neoadjuvant treatment (%)	No	56 (100.0)	33 (100.0)	80 (96.4)	90 (97.8)	49 (100.0)	44 (97.8)	76 (100.0)	189 (100.0)	0.065	exact
	Yes	0 (0.0)	0 (0.0)	3 (3.6)	2 (2.2)	0 (0.0)	1 (2.2)	0 (0.0)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		55.00 [47.75, 65.00]	62.00 [50.00, 68.00]	58.00 [47.50, 67.50]	55.50 [50.00, 66.00]	58.00 [49.00, 67.00]	61.00 [50.00, 71.00]	59.00 [48.00, 69.25]	56.00 [48.00, 64.00]	0.757	nonnorm
ethnicity (%)	HISPANIC OR LATINO	3 (6.4)	2 (6.9)	3 (3.9)	5 (5.9)	3 (6.8)	0 (0.0)	2 (2.7)	9 (5.2)	0.751	exact
	NOT HISPANIC OR LATINO	44 (93.6)	27 (93.1)	73 (96.1)	80 (94.1)	41 (93.2)	37 (100.0)	72 (97.3)	163 (94.8)		
primary lymph node presentation assessment (%)	NO	0 (0.0)	2 (7.7)	5 (7.9)	6 (8.1)	2 (5.6)	2 (5.4)	2 (3.2)	7 (4.3)	0.539	exact
	YES	39 (100.0)	24 (92.3)	58 (92.1)	68 (91.9)	34 (94.4)	35 (94.6)	60 (96.8)	156 (95.7)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	26 (46.4)	28 (84.8)	69 (83.1)	75 (82.4)	35 (71.4)	24 (53.3)	56 (73.7)	123 (64.4)		
	Infiltrating Lobular Carcinoma	27 (48.2)	4 (12.1)	3 (3.6)	5 (5.5)	9 (18.4)	16 (35.6)	13 (17.1)	41 (21.5)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	5 (5.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	1 (1.2)	3 (3.3)	0 (0.0)	0 (0.0)	0 (0.0)	3 (1.6)		
	Mixed Histology (please specify)	1 (1.8)	1 (3.0)	4 (4.8)	0 (0.0)	4 (8.2)	5 (11.1)	2 (2.6)	4 (2.1)		
	Mucinous Carcinoma	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (2.0)	0 (0.0)	0 (0.0)	8 (4.2)		
	Other, specify	2 (3.6)	0 (0.0)	5 (6.0)	2 (2.2)	0 (0.0)	0 (0.0)	5 (6.6)	12 (6.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
menopausal status (%)	Indeterminate (neither Pre or Post-menopausal)	2 (3.8)	0 (0.0)	1 (1.3)	2 (2.5)	1 (2.2)	1 (2.4)	3 (4.2)	4 (2.2)	0.671	exact
	Peri (6-12 months since last menstrual period)	4 (7.7)	1 (3.3)	2 (2.7)	5 (6.2)	2 (4.4)	0 (0.0)	3 (4.2)	3 (1.7)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	31 (59.6)	22 (73.3)	56 (74.7)	60 (74.1)	33 (73.3)	32 (76.2)	51 (71.8)	120 (67.4)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	15 (28.8)	7 (23.3)	16 (21.3)	14 (17.3)	9 (20.0)	9 (21.4)	14 (19.7)	51 (28.7)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.4)	1 (0.6)	<0.001	exact
	Negative	9 (16.1)	2 (6.9)	25 (31.6)	67 (77.9)	7 (14.6)	3 (7.7)	13 (18.3)	58 (32.6)		
	Positive	47 (83.9)	27 (93.1)	54 (68.4)	19 (22.1)	41 (85.4)	36 (92.3)	57 (80.3)	119 (66.9)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
breast carcinoma estrogen receptor status (%)	Negative	1 (1.8)	4 (13.8)	10 (12.7)	60 (69.8)	2 (4.2)	1 (2.6)	4 (5.6)	48 (26.8)	<0.001	
	Positive	55 (98.2)	25 (86.2)	69 (87.3)	26 (30.2)	46 (95.8)	38 (97.4)	67 (94.4)	131 (73.2)		
lab proc her2 neu immuno-histo-chemistry receptor status (%)	Equivocal	7 (14.9)	6 (23.1)	15 (22.1)	19 (24.7)	6 (14.3)	9 (27.3)	20 (32.8)	28 (18.3)	0.004	exact
	Indeterminate	4 (2.1)	0 (0.0)	3 (4.4)	2 (2.6)	1 (2.4)	0 (0.0)	2 (3.3)	1 (0.7)		
number of lymph nodes positive by ihc (median [IQR])	Negative	34 (72.3)	13 (50.0)	32 (47.1)	48 (62.3)	30 (71.4)	15 (45.5)	35 (57.4)	109 (71.2)		
	Positive	5 (10.6)	7 (26.9)	18 (26.5)	8 (10.4)	5 (11.9)	9 (27.3)	4 (6.6)	15 (9.8)	0.175	nonnorm
number of lymph nodes positive by he (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]		
		1.00 [0.00, 3.00]	1.00 [0.00, 4.00]	2.00 [0.00, 5.00]	0.00 [0.00, 2.25]	1.00 [0.00, 3.00]	0.00 [0.00, 3.00]	1.00 [0.00, 2.00]	1.00 [0.00, 2.00]	0.044	nonnorm
er level cell percentage category (%)	<10%	3 (9.1)	1 (6.2)	1 (2.6)	17 (54.8)	1 (3.8)	1 (5.3)	0 (0.0)	11 (12.6)	<0.001	exact
	10-19%	0 (0.0)	1 (6.2)	6 (15.8)	2 (6.5)	1 (3.8)	1 (5.3)	5 (11.9)	4 (4.6)		
	20-29%	0 (0.0)	0 (0.0)	0 (0.0)	3 (9.7)	0 (0.0)	1 (5.3)	0 (0.0)	0 (0.0)		
	30-39%	0 (0.0)	0 (0.0)	1 (2.6)	1 (3.2)	0 (0.0)	1 (5.3)	1 (2.4)	1 (1.1)		
	40-49%	0 (0.0)	1 (6.2)	0 (0.0)	1 (3.2)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)		
	50-59%	1 (3.0)	0 (0.0)	0 (0.0)	1 (3.2)	1 (3.8)	0 (0.0)	3 (7.1)	2 (2.3)		
	60-69%	1 (3.0)	1 (6.2)	0 (0.0)	0 (0.0)	1 (3.8)	1 (5.3)	2 (4.8)	1 (1.1)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
	70-79%	5 (15.2)	3 (18.8)	7 (18.4)	2 (6.5)	3 (11.5)	1 (5.3)	8 (19.0)	11 (12.6)		
	80-89%	2 (6.1)	2 (12.5)	2 (5.3)	0 (0.0)	1 (3.8)	4 (21.1)	1 (2.4)	6 (6.9)		
	90-99%	21 (63.6)	7 (43.8)	21 (55.3)	4 (12.9)	18 (69.2)	9 (47.4)	22 (52.4)	50 (57.5)		
progesterone receptor level cell percent category (%)	<10%	8 (23.5)	4 (30.8)	11 (37.9)	22 (78.6)	7 (29.2)	1 (5.9)	10 (25.6)	16 (20.3)	0.001	exact
	10-19%	1 (2.9)	1 (7.7)	3 (10.3)	0 (0.0)	2 (8.3)	3 (17.6)	4 (10.3)	4 (5.1)		
	20-29%	1 (2.9)	0 (0.0)	1 (3.4)	0 (0.0)	0 (0.0)	3 (17.6)	0 (0.0)	3 (3.8)		
	30-39%	1 (2.9)	0 (0.0)	3 (10.3)	0 (0.0)	1 (4.2)	0 (0.0)	0 (0.0)	3 (3.8)		
	40-49%	2 (5.9)	0 (0.0)	1 (3.4)	1 (3.6)	2 (8.3)	1 (5.9)	3 (7.7)	1 (1.3)		
	50-59%	2 (5.9)	0 (0.0)	1 (3.4)	1 (3.6)	1 (4.2)	0 (0.0)	2 (5.1)	3 (3.8)		
	60-69%	3 (8.8)	1 (7.7)	0 (0.0)	1 (3.6)	0 (0.0)	0 (0.0)	2 (5.1)	5 (6.3)		
	70-79%	3 (8.8)	2 (15.4)	1 (3.4)	2 (7.1)	2 (8.3)	2 (11.8)	7 (17.9)	10 (12.7)		
	80-89%	4 (11.8)	0 (0.0)	3 (10.3)	0 (0.0)	1 (4.2)	5 (29.4)	3 (7.7)	7 (8.9)		
	90-99%	9 (26.5)	5 (38.5)	5 (17.2)	1 (3.6)	8 (33.3)	2 (11.8)	8 (20.5)	27 (34.2)		
distant metastasis present ind2 (%)	NO	22 (100.0)	14 (100.0)	33 (100.0)	30 (100.0)	19 (100.0)	21 (100.0)	28 (93.3)	86 (95.6)	0.639	exact
	YES	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (6.7)	4 (4.4)		
stage event pathologic stage (%)	Stage I	1 (1.8)	2 (6.1)	3 (3.6)	9 (9.9)	10 (20.8)	1 (2.3)	10 (13.2)	14 (7.4)	0.005	exact
	Stage IA	6 (10.7)	2 (6.1)	3 (3.6)	5 (5.5)	4 (8.3)	3 (6.8)	6 (7.9)	20 (10.6)		
	Stage IB	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.1)	0 (0.0)	0 (0.0)	2 (1.1)		
	Stage II	1 (1.8)	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	1 (2.3)	0 (0.0)	2 (1.1)		
	Stage IIA	18 (32.1)	11 (33.3)	25 (30.1)	38 (41.8)	11 (22.9)	19 (43.2)	23 (30.3)	57 (30.2)		
	Stage IIB	15 (26.8)	7 (21.2)	20 (24.1)	17 (18.7)	9 (18.8)	8 (18.2)	18 (23.7)	54 (28.6)		
	Stage III	0 (0.0)	1 (3.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.5)		
	Stage IIIA	7 (12.5)	10 (30.3)	18 (21.7)	9 (9.9)	11 (22.9)	6 (13.6)	13 (17.1)	26 (13.8)		
	Stage IIIB	1 (1.8)	0 (0.0)	5 (6.0)	3 (3.3)	1 (2.1)	0 (0.0)	0 (0.0)	4 (2.1)		
	Stage IIIC	7 (12.5)	0 (0.0)	8 (9.6)	4 (4.4)	0 (0.0)	6 (13.6)	2 (2.6)	9 (4.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
Stage IV	0 (0.0)	0 (0.0)	1 (1.2)	3 (3.3)	1 (2.1)	0 (0.0)	2 (2.6)	0 (0.0)		
Stage X	0 (0.0)	0 (0.0)	0 (0.0)	2 (2.2)	0 (0.0)	0 (0.0)	2 (2.6)	0 (0.0)		