

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	ER1	ER2	p	test
n		457	130		
Subtype (%)	ER1	457 (100.0)	0 (0.0)	<0.001	
	ER2	0 (0.0)	130 (100.0)		
vital status (%)	Alive	118 (93.7)	28 (96.6)	1.000	exact
	Dead	8 (6.3)	1 (3.4)		
days to birth (median [IQR])		-21438.00 [-24942.25, -18024.00]	-20481.00 [-23974.25, -17679.75]	0.151	nonnorm
days to death (median [IQR])		1590.50 [686.50, 2481.75]	785.00 [524.00, 1927.00]	0.210	nonnorm
days to last followup (median [IQR])		753.00 [502.50, 1667.00]	915.00 [439.00, 1836.00]	0.964	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.8)	<0.001	exact
	ASIAN	21 (4.7)	9 (7.0)		
	BLACK OR AFRICAN AMERICAN	80 (17.9)	40 (31.2)		
	WHITE	345 (77.4)	78 (60.9)		
history of neoadjuvant treatment (%)	No	450 (98.7)	129 (100.0)	0.347	exact
	Yes	6 (1.3)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [49.00, 67.00]	55.50 [48.00, 65.00]	0.162	nonnorm
ethnicity (%)	HISPANIC OR LATINO	19 (4.7)	8 (6.5)	0.580	
	NOT HISPANIC OR LATINO	384 (95.3)	115 (93.5)		
primary lymph node presentation assessment (%)	NO	17 (4.7)	6 (5.8)	0.844	
	YES	343 (95.3)	97 (94.2)		
histological type (%)	Infiltrating Carcinoma	0 (0.0)	1 (0.8)	<0.001	exact
	NOS				
	Infiltrating Ductal Carcinoma	300 (65.6)	106 (82.2)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ER1	ER2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	111 (24.3)	4 (3.1)	0.019	exact
	Medullary Carcinoma	1 (0.2)	3 (2.3)		
	Metaplastic Carcinoma	1 (0.2)	6 (4.7)		
	Mixed Histology (please specify)	17 (3.7)	1 (0.8)		
	Mucinous Carcinoma	9 (2.0)	0 (0.0)		
	Other, specify	18 (3.9)	8 (6.2)		
	Indeterminate (neither Pre or Postmenopausal)	7 (1.7)	7 (5.8)		
	Peri (6-12 months since last menstrual period)	11 (2.6)	6 (5.0)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	296 (70.8)	86 (71.7)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	104 (24.9)	21 (17.5)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.8)	<0.001	exact
	Negative	69 (15.1)	115 (89.1)		
	Positive	387 (84.7)	13 (10.1)		
	Equivocal	87 (21.8)	23 (21.3)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Indeterminate	8 (2.0)	2 (1.9)	0.991	exact
	Negative	249 (62.4)	67 (62.0)		
	Positive	55 (13.8)	16 (14.8)		
		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.058	nonnorm
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.004	nonnorm
er level cell percentage category (%)	<10%	10 (3.7)	25 (100.0)	<0.001	exact
	10-19%	20 (7.5)	0 (0.0)		
	20-29%	4 (1.5)	0 (0.0)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ER1	ER2	p	test
	30-39%	5 (1.9)	0 (0.0)		
	40-49%	3 (1.1)	0 (0.0)		
	50-59%	8 (3.0)	0 (0.0)		
	60-69%	7 (2.6)	0 (0.0)		
	70-79%	40 (15.0)	0 (0.0)		
	80-89%	18 (6.7)	0 (0.0)		
	90-99%	152 (56.9)	0 (0.0)		
progesterone receptor level cell percent category (%)	<10%	51 (21.9)	28 (93.3)	<0.001	exact
	10-19%	18 (7.7)	0 (0.0)		
	20-29%	7 (3.0)	1 (3.3)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.7)	0 (0.0)		
	50-59%	10 (4.3)	0 (0.0)		
	60-69%	12 (5.2)	0 (0.0)		
	70-79%	29 (12.4)	0 (0.0)		
	80-89%	23 (9.9)	0 (0.0)		
	90-99%	64 (27.5)	1 (3.3)		
distant metastasis present ind2 (%)	NO	179 (98.4)	48 (94.1)	0.121	exact
	YES	3 (1.6)	3 (5.9)		
stage event pathologic stage (%)	Stage I	42 (9.3)	8 (6.2)	0.039	exact
	Stage IA	33 (7.3)	11 (8.6)		
	Stage IB	3 (0.7)	0 (0.0)		
	Stage II	4 (0.9)	1 (0.8)		
	Stage IIA	129 (28.4)	59 (46.1)		
	Stage IIB	120 (26.4)	21 (16.4)		
	Stage III	1 (0.2)	1 (0.8)		
	Stage IIIA	76 (16.7)	17 (13.3)		
	Stage IIIB	10 (2.2)	2 (1.6)		
	Stage IIIC	29 (6.4)	5 (3.9)		
	Stage IV	4 (0.9)	2 (1.6)		
	Stage X	3 (0.7)	1 (0.8)		