

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	ANF1	ANF2	p	test
n		317	147		
Subtype (%)	ANF1	317 (100.0)	0 (0.0)	<0.001	
	ANF2	0 (0.0)	147 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-22687.50 [-26003.75, -18651.25]	-20339.00 [-23539.75, -17497.75]	<0.001	nonnorm
days to death (median [IQR])		1563.00 [883.00, 2573.00]	991.00 [651.00, 1624.00]	0.068	nonnorm
days to last followup (median [IQR])		865.00 [451.00, 1519.00]	968.50 [405.75, 1948.25]	0.572	nonnorm
race list (%)	ASIAN	18 (7.2)	11 (8.3)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	20 (8.0)	37 (27.8)		
	WHITE	212 (84.8)	85 (63.9)		
history of neoadjuvant treatment (%)	No	310 (97.8)	147 (100.0)	0.103	exact
	Yes	7 (2.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [50.00, 71.00]	55.00 [47.50, 64.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (3.9)	1 (0.8)	0.175	exact
	NOT HISPANIC OR LATINO	221 (96.1)	123 (99.2)		
primary lymph node presentation assessment (%)	NO	2 (1.4)	0 (0.0)	1.000	exact
	YES	145 (98.6)	67 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	210 (66.2)	130 (88.4)	<0.001	exact
	Infiltrating Lobular Carcinoma	79 (24.9)	7 (4.8)		
	Medullary Carcinoma	0 (0.0)	1 (0.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ANF1	ANF2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (2.0)		
	Mixed Histology (please specify)	6 (1.9)	2 (1.4)		
	Mucinous Carcinoma	7 (2.2)	0 (0.0)		
	Other, specify	15 (4.7)	4 (2.7)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	12 (4.0)	8 (5.8)	0.384	exact
	Peri (6-12 months since last menstrual period)	12 (4.0)	6 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	217 (72.6)	90 (64.7)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	58 (19.4)	35 (25.2)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	40 (12.9)	120 (83.3)		
	Positive	268 (86.5)	24 (16.7)		
breast carcinoma estrogen receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	5 (1.6)	106 (73.6)		
	Positive	303 (97.7)	38 (26.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	51 (18.4)	20 (15.4)	0.455	exact
	Indeterminate	2 (0.7)	0 (0.0)		
	Negative	168 (60.6)	76 (58.5)		
	Positive	56 (20.2)	34 (26.2)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.028	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.113	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ANF1	ANF2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	0.002	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 5.00]	1.00 [1.00, 2.00]	0.005	nonnorm
distant metastasis present ind2 (%)	NO	90 (96.8)	39 (95.1)	0.641	exact
	YES	3 (3.2)	2 (4.9)		
stage event pathologic stage (%)	Stage I	33 (10.5)	7 (4.9)	0.057	exact
	Stage IA	25 (8.0)	13 (9.0)		
	Stage IB	3 (1.0)	0 (0.0)		
	Stage IIA	97 (30.9)	59 (41.0)		
	Stage IIB	67 (21.3)	39 (27.1)		
	Stage IIIA	39 (12.4)	15 (10.4)		
	Stage IIIB	10 (3.2)	2 (1.4)		
	Stage IIIC	21 (6.7)	7 (4.9)		
	Stage IV	10 (3.2)	2 (1.4)		
	Stage X	9 (2.9)	0 (0.0)		