

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CIMLR1	CIMLR2	p	test
n		266	198		
Subtype (%)	CIMLR1	266 (100.0)	0 (0.0)	<0.001	
	CIMLR2	0 (0.0)	198 (100.0)		
vital status (%)	Alive	6 (100.0)	5 (83.3)	1.000	exact
	Dead	0 (0.0)	1 (16.7)		
days to birth (median [IQR])		-22778.00 [-26415.75, -18746.25]	-20691.50 [-23968.25, -17613.00]	<0.001	nonnorm
days to death (median [IQR])		1711.50 [893.50, 2813.50]	1079.50 [718.50, 1693.75]	0.140	nonnorm
days to last followup (median [IQR])		822.00 [453.50, 1367.00]	984.00 [410.25, 1911.50]	0.290	nonnorm
race list (%)	ASIAN	12 (5.6)	17 (10.1)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	18 (8.4)	39 (23.2)		
	WHITE	185 (86.0)	112 (66.7)		
history of neoadjuvant treatment (%)	No	260 (97.7)	197 (99.5)	0.247	exact
	Yes	6 (2.3)	1 (0.5)		
age at initial pathologic diagnosis (median [IQR])		62.00 [51.00, 71.75]	56.00 [48.00, 64.75]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	8 (4.0)	2 (1.3)	0.196	exact
	NOT HISPANIC OR LATINO	191 (96.0)	153 (98.7)		
primary lymph node presentation assessment (%)	NO	2 (1.6)	0 (0.0)	0.517	exact
	YES	126 (98.4)	86 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	168 (63.2)	172 (86.9)	<0.001	exact
	Infiltrating Lobular Carcinoma	70 (26.3)	16 (8.1)		
	Medullary Carcinoma	0 (0.0)	1 (0.5)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
	Metaplastic Carcinoma	1 (0.4)	2 (1.0)		
	Mixed Histology (please specify)	6 (2.3)	2 (1.0)		
	Mucinous Carcinoma	7 (2.6)	0 (0.0)		
	Other, specify	14 (5.3)	5 (2.5)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	8 (3.2)	12 (6.4)	0.264	exact
	Peri (6-12 months since last menstrual period)	10 (4.0)	8 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	184 (73.3)	123 (65.8)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	49 (19.5)	44 (23.5)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.8)	0 (0.0)	<0.001	exact
	Negative	27 (10.4)	133 (68.2)		
	Positive	230 (88.8)	62 (31.8)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (1.0)	<0.001	exact
	Negative	4 (1.5)	107 (54.9)		
	Positive	255 (98.5)	86 (44.1)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	47 (20.3)	24 (13.7)	0.003	exact
	Indeterminate	2 (0.9)	0 (0.0)		
	Negative	145 (62.5)	99 (56.6)		
	Positive	38 (16.4)	52 (29.7)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.020	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.373	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 6.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
distant metastasis present ind2 (%)	NO	78 (97.5)	51 (94.4)	0.392	exact
	YES	2 (2.5)	3 (5.6)		
stage event pathologic stage (%)	Stage I	24 (9.1)	16 (8.2)	0.556	exact
	Stage IA	25 (9.5)	13 (6.7)		
	Stage IB	3 (1.1)	0 (0.0)		
	Stage IIA	83 (31.4)	73 (37.6)		
	Stage IIB	58 (22.0)	48 (24.7)		
	Stage IIIA	33 (12.5)	21 (10.8)		
	Stage IIIB	9 (3.4)	3 (1.5)		
	Stage IIIC	16 (6.1)	12 (6.2)		
	Stage IV	6 (2.3)	6 (3.1)		
	Stage X	7 (2.7)	2 (1.0)		