

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	1	2	p	test
n		227	237		
Subtype (%)	COCA1	227 (100.0)	0 (0.0)	<0.001	
	COCA2	0 (0.0)	237 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-21488.00 [- 24290.50, -17718.50]	-22686.00 [- 25994.00, -18535.00]	0.030	nonnorm
days to death (median [IQR])		1556.00 [956.00, 2000.00]	1127.00 [792.00, 2009.00]	0.635	nonnorm
days to last followup (median [IQR])		881.00 [498.50, 1746.50]	900.50 [394.50, 1602.75]	0.500	nonnorm
race list (%)	ASIAN	11 (5.7)	18 (9.5)	0.157	exact
	BLACK OR AFRICAN AMERICAN	25 (13.0)	32 (16.8)		
	WHITE	157 (81.3)	140 (73.7)		
history of neoadjuvant treatment (%)	No	225 (99.1)	232 (97.9)	0.450	exact
	Yes	2 (0.9)	5 (2.1)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 66.00]	61.00 [50.00, 70.00]	0.054	nonnorm
ethnicity (%)	HISPANIC OR LATINO	2 (1.1)	8 (4.4)	0.105	exact
	NOT HISPANIC OR LATINO	172 (98.9)	172 (95.6)		
primary lymph node presentation assessment (%)	NO	1 (0.9)	1 (1.0)	1.000	exact
	YES	114 (99.1)	98 (99.0)		
histological type (%)	Infiltrating Ductal Carcinoma	163 (71.8)	177 (74.7)	0.174	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	1	2	p	test
	Infiltrating Lobular Carcinoma	50 (22.0)	36 (15.2)		
	Medullary Carcinoma	1 (0.4)	0 (0.0)		
	Metaplastic Carcinoma	2 (0.9)	1 (0.4)		
	Mixed Histology (please specify)	3 (1.3)	5 (2.1)		
	Mucinous Carcinoma	2 (0.9)	5 (2.1)		
	Other, specify	6 (2.6)	13 (5.5)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	6 (2.8)	14 (6.2)	0.258	exact
	Peri (6-12 months since last menstrual period)	11 (5.1)	7 (3.1)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	149 (69.6)	158 (70.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	48 (22.4)	45 (20.1)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	2 (0.9)	0.061	exact
	Negative	88 (39.3)	72 (31.3)		
	Positive	136 (60.7)	156 (67.8)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (0.9)	0.005	exact
	Negative	67 (29.9)	44 (19.1)		
	Positive	157 (70.1)	184 (80.0)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	30 (15.0)	41 (19.8)	0.011	exact
	Indeterminate	1 (0.5)	1 (0.5)		
	Negative	135 (67.5)	109 (52.7)		
	Positive	34 (17.0)	56 (27.1)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.223	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	1	2	p	test
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.955	nonnorm
er level cell percentage category (median [IQR])		1.00 [1.00, 8.00]	1.00 [1.00, 7.00]	0.622	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	0.738	nonnorm
distant metastasis present ind2 (%)	NO	76 (97.4)	53 (94.6)	0.649	exact
	YES	2 (2.6)	3 (5.4)		
stage event pathologic stage (%)	Stage I	22 (9.9)	18 (7.6)	0.571	exact
	Stage IA	22 (9.9)	16 (6.8)		
	Stage IB	1 (0.5)	2 (0.8)		
	Stage IIA	78 (35.1)	78 (33.1)		
	Stage IIB	53 (23.9)	53 (22.5)		
	Stage IIIA	19 (8.6)	35 (14.8)		
	Stage IIIB	4 (1.8)	8 (3.4)		
	Stage IIIC	14 (6.3)	14 (5.9)		
	Stage IV	5 (2.3)	7 (3.0)		
	Stage X	4 (1.8)	5 (2.1)		