

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	KLIC1	KLIC2	p	test
n		291	173		
Subtype (%)	KLIC1	291 (100.0)	0 (0.0)	<0.001	
	KLIC2	0 (0.0)	173 (100.0)		
vital status (%)	Alive	9 (100.0)	2 (66.7)	0.250	exact
	Dead	0 (0.0)	1 (33.3)		
days to birth (median [IQR])		-22629.00 [-26140.50, -18467.00]	-21014.00 [-24288.00, -17930.00]	0.016	nonnorm
days to death (median [IQR])		1602.50 [926.50, 2414.00]	1000.00 [599.50, 1658.00]	0.086	nonnorm
days to last followup (median [IQR])		923.50 [424.00, 1676.25]	867.00 [441.00, 1563.00]	0.597	nonnorm
race list (%)	ASIAN	21 (8.9)	8 (5.4)	0.065	exact
	BLACK OR AFRICAN AMERICAN	28 (11.9)	29 (19.7)		
	WHITE	187 (79.2)	110 (74.8)		
history of neoadjuvant treatment (%)	No	285 (97.9)	172 (99.4)	0.266	exact
	Yes	6 (2.1)	1 (0.6)		
age at initial pathologic diagnosis (median [IQR])		61.00 [50.00, 71.00]	57.00 [49.00, 66.00]	0.025	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (4.1)	1 (0.7)	0.096	exact
	NOT HISPANIC OR LATINO	208 (95.9)	136 (99.3)		
primary lymph node presentation assessment (%)	NO	1 (0.7)	1 (1.3)	1.000	exact
	YES	134 (99.3)	78 (98.7)		
histological type (%)	Infiltrating Ductal Carcinoma	195 (67.0)	145 (83.8)	<0.001	exact
	Infiltrating Lobular Carcinoma	73 (25.1)	13 (7.5)		
	Medullary Carcinoma	0 (0.0)	1 (0.6)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	KLIC1	KLIC2	p	test
	Metaplastic Carcinoma	1 (0.3)	2 (1.2)		
	Mixed Histology (please specify)	6 (2.1)	2 (1.2)		
	Mucinous Carcinoma	5 (1.7)	2 (1.2)		
	Other, specify	11 (3.8)	8 (4.6)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	9 (3.3)	11 (6.7)	0.430	exact
	Peri (6-12 months since last menstrual period)	11 (4.0)	7 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	196 (71.3)	111 (68.1)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	59 (21.5)	34 (20.9)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.3)	1 (0.6)	<0.001	exact
	Negative	58 (20.3)	102 (60.7)		
	Positive	227 (79.4)	65 (38.7)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (1.2)	<0.001	exact
	Negative	25 (8.7)	86 (51.2)		
	Positive	261 (91.3)	80 (47.6)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	42 (16.2)	29 (19.7)	0.271	exact
	Indeterminate	2 (0.8)	0 (0.0)		
	Negative	152 (58.5)	92 (62.6)		
	Positive	64 (24.6)	26 (17.7)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.574	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.039	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	KLIC1	KLIC2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 10.00]	1.00 [1.00, 2.00]	0.040	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 3.00]	1.00 [1.00, 2.00]	0.041	nonnorm
distant metastasis present ind2 (%)	NO	77 (96.2)	52 (96.3)	1.000	exact
	YES	3 (3.8)	2 (3.7)		
stage event pathologic stage (%)	Stage I	31 (10.8)	9 (5.3)	0.014	exact
	Stage IA	23 (8.0)	15 (8.8)		
	Stage IB	2 (0.7)	1 (0.6)		
	Stage IIA	82 (28.5)	74 (43.5)		
	Stage IIB	67 (23.3)	39 (22.9)		
	Stage IIIA	35 (12.2)	19 (11.2)		
	Stage IIIB	10 (3.5)	2 (1.2)		
	Stage IIIC	24 (8.3)	4 (2.4)		
	Stage IV	7 (2.4)	5 (2.9)		
	Stage X	7 (2.4)	2 (1.2)		