

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
n		96	165	70	133		
Subtype (%)	LRAcluster1	96 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	LRAcluster2	0 (0.0)	165 (100.0)	0 (0.0)	0 (0.0)		
	LRAcluster3	0 (0.0)	0 (0.0)	70 (100.0)	0 (0.0)		
	LRAcluster4	0 (0.0)	0 (0.0)	0 (0.0)	133 (100.0)		
vital status (%)	Alive	5 (100.0)	4 (100.0)	0 (0.0)	2 (100.0)	0.096	exact
	Dead	0 (0.0)	0 (0.0)	1 (100.0)	0 (0.0)		
days to birth (median [IQR])		-21622.00 [-, 23585.50, -17004.50]	-22710.00 [-, 26548.00, -18875.00]	-20841.50 [-, 22951.25, -17638.50]	-22430.00 [-, 25948.25, -18509.25]	0.013	nonnorm
days to death (median [IQR])		2183.00 [1173.75, 2965.00]	1642.00 [921.00, 2009.00]	1000.00 [599.50, 1425.00]	1142.00 [811.00, 1993.00]	0.273	nonnorm
days to last followup (median [IQR])		1039.00 [621.50, 2203.25]	822.00 [524.00, 1357.00]	954.00 [404.00, 1849.25]	797.00 [366.25, 1488.00]	0.037	nonnorm
race list (%)	ASIAN	5 (5.8)	5 (3.7)	3 (4.8)	16 (16.2)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	21 (24.4)	9 (6.7)	16 (25.4)	11 (11.1)		
	WHITE	60 (69.8)	121 (89.6)	44 (69.8)	72 (72.7)		
history of neoadjuvant treatment (%)	No	96 (100.0)	161 (97.6)	70 (100.0)	130 (97.7)	0.318	exact
	Yes	0 (0.0)	4 (2.4)	0 (0.0)	3 (2.3)		
age at initial pathologic diagnosis (median [IQR])		58.50 [46.00, 64.00]	62.00 [51.00, 72.00]	57.00 [48.00, 62.00]	59.00 [50.00, 70.00]	0.013	nonnorm
ethnicity (%)	HISPANIC OR LATINO	3 (3.6)	6 (4.9)	0 (0.0)	1 (1.1)	0.234	exact
	NOT HISPANIC OR LATINO	81 (96.4)	116 (95.1)	56 (100.0)	91 (98.9)		
primary lymph node presentation assessment (%)	NO	1 (1.6)	0 (0.0)	0 (0.0)	1 (2.6)	0.378	exact
	YES	60 (98.4)	84 (100.0)	30 (100.0)	38 (97.4)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
histological type (%)	Infiltrating Ductal Carcinoma	54 (56.2)	101 (61.2)	69 (98.6)	116 (87.2)	<0.001	exact
	Infiltrating Lobular Carcinoma	31 (32.3)	46 (27.9)	0 (0.0)	9 (6.8)		
	Medullary Carcinoma	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	2 (2.1)	0 (0.0)	1 (1.4)	0 (0.0)		
	Mixed Histology (please specify)	4 (4.2)	3 (1.8)	0 (0.0)	1 (0.8)		
	Mucinous Carcinoma	0 (0.0)	6 (3.6)	0 (0.0)	1 (0.8)		
menopause status (%)	Other, specify	4 (4.2)	9 (5.5)	0 (0.0)	6 (4.5)	0.368	exact
	Indeterminate (neither Pre or Postmenopausal)	5 (5.6)	2 (1.3)	5 (7.8)	8 (6.2)		
	Peri (6-12 months since last menstrual period)	2 (2.2)	8 (5.1)	3 (4.7)	5 (3.9)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	60 (67.4)	114 (73.1)	45 (70.3)	88 (68.2)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	22 (24.7)	32 (20.5)	11 (17.2)	28 (21.7)		
	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.6)		
breast carcinoma progesterone receptor status (%)						<0.001	exact
	Negative	43 (46.2)	14 (8.6)	66 (94.3)	37 (28.7)		
	Positive	50 (53.8)	148 (91.4)	4 (5.7)	90 (69.8)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.6)	<0.001	exact
	Negative	37 (39.8)	1 (0.6)	61 (87.1)	12 (9.3)		
	Positive	56 (60.2)	161 (99.4)	9 (12.9)	115 (89.1)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	8 (10.0)	30 (20.7)	13 (21.3)	20 (16.5)	<0.001	exact
	Indeterminate	0 (0.0)	2 (1.4)	0 (0.0)	0 (0.0)		
	Negative	55 (68.8)	94 (64.8)	40 (65.6)	55 (45.5)		
number of lymphnodes positive by ihc (median [IQR])	Positive	17 (21.2)	19 (13.1)	8 (13.1)	46 (38.0)	0.213	nonnorm
		0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	1.00 [0.00, 2.00]	0.085	nonnorm
er level cell percentage category (median [IQR])		1.00 [1.00, 7.25]	1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 9.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
distant metastasis present ind2 (%)	NO	33 (100.0)	54 (98.2)	15 (88.2)	27 (93.1)	0.094	exact
	YES	0 (0.0)	1 (1.8)	2 (11.8)	2 (6.9)		
stage event pathologic stage (%)	Stage I	9 (9.5)	20 (12.3)	5 (7.5)	6 (4.5)	0.004	exact
	Stage IA	10 (10.5)	20 (12.3)	4 (6.0)	4 (3.0)		
	Stage IB	0 (0.0)	1 (0.6)	0 (0.0)	2 (1.5)		
	Stage IIA	22 (23.2)	59 (36.2)	33 (49.3)	42 (31.6)		
	Stage IIB	21 (22.1)	29 (17.8)	15 (22.4)	41 (30.8)		
	Stage IIIA	11 (11.6)	17 (10.4)	7 (10.4)	19 (14.3)		
	Stage IIIB	4 (4.2)	2 (1.2)	0 (0.0)	6 (4.5)		
	Stage IIIC	13 (13.7)	7 (4.3)	2 (3.0)	6 (4.5)		
	Stage IV	2 (2.1)	4 (2.5)	1 (1.5)	5 (3.8)		
	Stage X	3 (3.2)	4 (2.5)	0 (0.0)	2 (1.5)		