

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MFA1	MFA2	MFA3	p
n		153	189	122	
Subtype (%)	MFA1	153 (100.0)	0 (0.0)	0 (0.0)	<0.001
	MFA2	0 (0.0)	189 (100.0)	0 (0.0)	
	MFA3	0 (0.0)	0 (0.0)	122 (100.0)	
vital status (%)	Alive	4 (100.0)	4 (100.0)	3 (75.0)	1.000
	Dead	0 (0.0)	0 (0.0)	1 (25.0)	
days to birth (median [IQR])		-21085.00 [-25165.00, -17901.25]	-23560.00 [-26446.00, -19763.00]	-20105.00 [-22858.00, -17569.00]	<0.001
days to death (median [IQR])		2100.00 [1517.75, 2914.50]	1009.00 [770.50, 1809.50]	991.00 [487.00, 1414.00]	0.007
days to last followup (median [IQR])		970.50 [547.00, 2103.75]	754.00 [385.00, 1299.50]	996.50 [426.50, 2033.00]	0.007
race list (%)	ASIAN	8 (5.8)	13 (9.8)	8 (7.2)	0.001
	BLACK OR AFRICAN AMERICAN	14 (10.1)	14 (10.5)	29 (26.1)	
	WHITE	117 (84.2)	106 (79.7)	74 (66.7)	
history of neoadjuvant treatment (%)	No	151 (98.7)	184 (97.4)	122 (100.0)	0.185
	Yes	2 (1.3)	5 (2.6)	0 (0.0)	
age at initial pathologic diagnosis (median [IQR])		57.00 [48.00, 68.00]	64.00 [53.00, 72.00]	54.50 [48.00, 62.00]	<0.001
ethnicity (%)	HISPANIC OR LATINO	7 (5.3)	2 (1.7)	1 (1.0)	0.127
	NOT HISPANIC OR LATINO	125 (94.7)	118 (98.3)	101 (99.0)	
primary lymph node presentation assessment (%)	NO	1 (1.0)	1 (1.7)	0 (0.0)	1.000
	YES	101 (99.0)	57 (98.3)	54 (100.0)	
histological type (%)	Infiltrating Ductal Carcinoma	75 (49.0)	155 (82.0)	110 (90.2)	<0.001
	Infiltrating Lobular Carcinoma	69 (45.1)	14 (7.4)	3 (2.5)	
	Medullary Carcinoma	0 (0.0)	0 (0.0)	1 (0.8)	

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MFA1	MFA2	MFA3	p
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	3 (2.5)	
	Mixed Histology (please specify)	4 (2.6)	2 (1.1)	2 (1.6)	
	Mucinous Carcinoma	1 (0.7)	6 (3.2)	0 (0.0)	
	Other, specify	4 (2.6)	12 (6.3)	3 (2.5)	
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	4 (2.9)	8 (4.4)	8 (7.0)	0.157
	Peri (6-12 months since last menstrual period)	8 (5.7)	4 (2.2)	6 (5.2)	
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	95 (67.9)	139 (76.0)	73 (63.5)	
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	33 (23.6)	32 (17.5)	28 (24.3)	
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	2 (1.1)	0 (0.0)	<0.001
	Negative	24 (15.7)	31 (17.0)	105 (88.2)	
	Positive	129 (84.3)	149 (81.9)	14 (11.8)	
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (1.1)	0 (0.0)	<0.001
	Negative	13 (8.5)	3 (1.6)	95 (79.8)	
	Positive	140 (91.5)	177 (97.3)	24 (20.2)	
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	18 (13.7)	35 (20.6)	18 (17.0)	0.039
	Indeterminate	2 (1.5)	0 (0.0)	0 (0.0)	
	Negative	90 (68.7)	89 (52.4)	65 (61.3)	
	Positive	21 (16.0)	46 (27.1)	23 (21.7)	
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.030
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.038

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MFA1	MFA2	MFA3	p
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 9.00]	1.00 [1.00, 2.00]	<0.001
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 6.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	<0.001
distant metastasis present ind2 (%)	NO	62 (98.4)	36 (94.7)	31 (93.9)	0.512
	YES	1 (1.6)	2 (5.3)	2 (6.1)	
stage event pathologic stage (%)	Stage I	16 (10.6)	18 (9.6)	6 (5.0)	0.006
	Stage IA	14 (9.3)	13 (6.9)	11 (9.2)	
	Stage IB	0 (0.0)	3 (1.6)	0 (0.0)	
	Stage IIA	39 (25.8)	66 (35.1)	51 (42.9)	
	Stage IIB	33 (21.9)	43 (22.9)	30 (25.2)	
	Stage IIIA	23 (15.2)	19 (10.1)	12 (10.1)	
	Stage IIIB	1 (0.7)	9 (4.8)	2 (1.7)	
	Stage IIIC	17 (11.3)	6 (3.2)	5 (4.2)	
	Stage IV	3 (2.0)	7 (3.7)	2 (1.7)	
	Stage X	5 (3.3)	4 (2.1)	0 (0.0)	