

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MONET1	MONET2	MONET3	p	test
n		136	177	151		
Subtype (%)	MONET1	136 (100.0)	0 (0.0)	0 (0.0)	<0.001	
	MONET2	0 (0.0)	177 (100.0)	0 (0.0)		
	MONET3	0 (0.0)	0 (0.0)	151 (100.0)		
vital status (%)	Alive	4 (80.0)	4 (100.0)	3 (100.0)	1.000	exact
	Dead	1 (20.0)	0 (0.0)	0 (0.0)		
days to birth (median [IQR])		-20332.00 [-23316.50, -17521.50]	-22357.00 [-26486.50, -18512.25]	-23164.00 [-25811.00, -19216.00]	<0.001	nonnorm
days to death (median [IQR])		991.00 [548.00, 1556.00]	1895.00 [1001.75, 2914.50]	1142.00 [811.00, 1993.00]	0.091	nonnorm
days to last followup (median [IQR])		984.00 [426.50, 1974.75]	849.00 [529.50, 1758.50]	864.00 [385.00, 1371.00]	0.132	nonnorm
race list (%)	ASIAN	10 (8.3)	6 (3.9)	13 (12.0)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	34 (28.1)	9 (5.8)	14 (13.0)		
	WHITE	77 (63.6)	139 (90.3)	81 (75.0)		
history of neoadjuvant treatment (%)	No	136 (100.0)	174 (98.3)	147 (97.4)	0.192	exact
	Yes	0 (0.0)	3 (1.7)	4 (2.6)		

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
age at initial pathologic diagnosis (median [IQR])		55.00 [47.75, 63.25]	61.00 [50.00, 72.00]	63.00 [51.50, 70.00]	0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	1 (0.9)	8 (5.6)	1 (1.0)	0.063	exact
	NOT HISPANIC OR LATINO	111 (99.1)	136 (94.4)	97 (99.0)		
primary lymph node pre-sentation assessment (%)	NO	0 (0.0)	0 (0.0)	2 (4.5)	0.044	exact
	YES	62 (100.0)	108 (100.0)	42 (95.5)		
histological type (%)	Infiltrating Ductal Carcinoma	121 (89.0)	88 (49.7)	131 (86.8)	<0.001	exact
	Infiltrating Lobular Carcinoma	5 (3.7)	73 (41.2)	8 (5.3)		
	Medullary Carcinoma	1 (0.7)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	3 (2.2)	0 (0.0)	0 (0.0)		
	Mixed Histology (please specify)	2 (1.5)	5 (2.8)	1 (0.7)		
	Mucinous Carcinoma	0 (0.0)	4 (2.3)	3 (2.0)		
	Other, specify	4 (2.9)	7 (4.0)	8 (5.3)		
menopause status (%)	Indeterminat (neither Pre or Post-menopausal)	8 (6.2)	3 (1.9)	9 (6.1)	0.184	exact

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
	Peri (6-12 months since last menstrual period)	5 (3.9)	9 (5.6)	4 (2.7)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	82 (64.1)	118 (72.8)	107 (72.3)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	33 (25.8)	32 (19.8)	28 (18.9)		
breast carcinoma progesterone receptor status (%)	Indeterminat	0 (0.0)	0 (0.0)	2 (1.4)	<0.001	exact
	Negative	114 (85.7)	17 (9.8)	29 (19.7)		
	Positive	19 (14.3)	157 (90.2)	116 (78.9)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	2 (1.4)	<0.001	exact
	Negative	103 (77.4)	3 (1.7)	5 (3.4)		
	Positive	30 (22.6)	171 (98.3)	140 (95.2)		
lab proc her2 neu immunohistochemistry receptor status (%)	Equivocal	19 (16.0)	25 (16.6)	27 (19.7)	0.003	exact

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
	Indeterminate	0 (0.0)	2 (1.3)	0 (0.0)		
	Negative	70 (58.8)	105 (69.5)	69 (50.4)		
	Positive	30 (25.2)	19 (12.6)	41 (29.9)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.020	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.704	nonnorm
er level cell percent-age category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 11.00]	1.00 [1.00, 8.00]	<0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 8.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
distant metastasis present ind2 (%)	NO	36 (94.7)	64 (97.0)	29 (96.7)	0.856	exact
	YES	2 (5.3)	2 (3.0)	1 (3.3)		
stage event pathologic stage (%)	Stage I	7 (5.3)	24 (13.7)	9 (6.0)	0.005	exact
	Stage IA	11 (8.3)	19 (10.9)	8 (5.3)		
	Stage IB	0 (0.0)	0 (0.0)	3 (2.0)		
	Stage IIA	54 (40.6)	54 (30.9)	48 (32.0)		
	Stage IIB	34 (25.6)	31 (17.7)	41 (27.3)		
	Stage IIIA	15 (11.3)	17 (9.7)	22 (14.7)		
	Stage IIIB	2 (1.5)	4 (2.3)	6 (4.0)		
	Stage IIIC	8 (6.0)	16 (9.1)	4 (2.7)		
	Stage IV	2 (1.5)	6 (3.4)	4 (2.7)		
	Stage X	0 (0.0)	4 (2.3)	5 (3.3)		