

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	NEMO1	NEMO2	p	test
n		315	149		
Subtype (%)	NEMO1	315 (100.0)	0 (0.0)	<0.001	
	NEMO2	0 (0.0)	149 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-22716.50 [-26100.00, -18697.75]	-20339.00 [-23435.50, -17434.00]	<0.001	nonnorm
days to death (median [IQR])		1602.50 [892.50, 2550.50]	967.00 [599.50, 1485.00]	0.045	nonnorm
days to last followup (median [IQR])		835.50 [446.25, 1510.00]	996.00 [418.00, 2018.00]	0.318	nonnorm
race list (%)	ASIAN	18 (7.2)	11 (8.2)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	20 (8.0)	37 (27.6)		
	WHITE	211 (84.7)	86 (64.2)		
history of neoadjuvant treatment (%)	No	308 (97.8)	149 (100.0)	0.102	exact
	Yes	7 (2.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [50.00, 71.00]	55.00 [47.00, 64.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (3.9)	1 (0.8)	0.175	exact
	NOT HISPANIC OR LATINO	221 (96.1)	123 (99.2)		
primary lymph node presentation assessment (%)	NO	2 (1.4)	0 (0.0)	1.000	exact
	YES	143 (98.6)	69 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	209 (66.3)	131 (87.9)	<0.001	exact
	Infiltrating Lobular Carcinoma	78 (24.8)	8 (5.4)		
	Medullary Carcinoma	0 (0.0)	1 (0.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	NEMO1	NEMO2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (2.0)		
	Mixed Histology (please specify)	6 (1.9)	2 (1.3)		
	Mucinous Carcinoma	7 (2.2)	0 (0.0)		
	Other, specify	15 (4.8)	4 (2.7)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	12 (4.0)	8 (5.7)	0.241	exact
	Peri (6-12 months since last menstrual period)	12 (4.0)	6 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	217 (73.1)	90 (63.8)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	56 (18.9)	37 (26.2)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	40 (13.0)	120 (82.2)		
	Positive	266 (86.4)	26 (17.8)		
breast carcinoma estrogen receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	5 (1.6)	106 (72.6)		
	Positive	301 (97.7)	40 (27.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	51 (18.5)	20 (15.2)	0.380	exact
	Indeterminate	2 (0.7)	0 (0.0)		
	Negative	167 (60.7)	77 (58.3)		
	Positive	55 (20.0)	35 (26.5)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.073	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.129	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	NEMO1	NEMO2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 5.00]	1.00 [1.00, 2.00]	0.003	nonnorm
distant metastasis present ind2 (%)	NO	89 (96.7)	40 (95.2)	0.648	exact
	YES	3 (3.3)	2 (4.8)		
stage event pathologic stage (%)	Stage I	31 (9.9)	9 (6.2)	0.114	exact
	Stage IA	25 (8.0)	13 (8.9)		
	Stage IB	3 (1.0)	0 (0.0)		
	Stage IIA	98 (31.4)	58 (39.7)		
	Stage IIB	66 (21.2)	40 (27.4)		
	Stage IIIA	39 (12.5)	15 (10.3)		
	Stage IIIB	10 (3.2)	2 (1.4)		
	Stage IIIC	21 (6.7)	7 (4.8)		
	Stage IV	10 (3.2)	2 (1.4)		
	Stage X	9 (2.9)	0 (0.0)		