

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	RWR-NF1	RWR-NF2	p	test
n		303	161		
Subtype (%)	RWR-NF1	303 (100.0)	0 (0.0)	<0.001	
	RWR-NF2	0 (0.0)	161 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-22723.00 [-26274.00, -18741.00]	-20332.00 [-23585.50, -17394.00]	<0.001	nonnorm
days to death (median [IQR])		1496.50 [829.00, 2715.50]	1087.00 [785.25, 1697.25]	0.221	nonnorm
days to last followup (median [IQR])		857.00 [453.25, 1519.75]	965.00 [393.00, 1871.00]	0.669	nonnorm
race list (%)	ASIAN	16 (6.7)	13 (9.1)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	18 (7.5)	39 (27.3)		
	WHITE	206 (85.8)	91 (63.6)		
history of neoadjuvant treatment (%)	No	296 (97.7)	161 (100.0)	0.102	exact
	Yes	7 (2.3)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [51.00, 71.00]	55.00 [47.00, 64.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (4.1)	1 (0.8)	0.098	exact
	NOT HISPANIC OR LATINO	213 (95.9)	131 (99.2)		
primary lymph node presentation assessment (%)	NO	2 (1.4)	0 (0.0)	1.000	exact
	YES	141 (98.6)	71 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	197 (65.0)	143 (88.8)	<0.001	exact
	Infiltrating Lobular Carcinoma	79 (26.1)	7 (4.3)		
	Medullary Carcinoma	0 (0.0)	1 (0.6)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-NF1	RWR-NF2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (1.9)		
	Mixed Histology (please specify)	5 (1.7)	3 (1.9)		
	Mucinous Carcinoma	7 (2.3)	0 (0.0)		
	Other, specify	15 (5.0)	4 (2.5)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	12 (4.2)	8 (5.3)	0.142	exact
	Peri (6-12 months since last menstrual period)	10 (3.5)	8 (5.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	211 (73.8)	96 (63.2)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	53 (18.5)	40 (26.3)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.7)	0 (0.0)	<0.001	exact
	Negative	34 (11.5)	126 (79.7)		
	Positive	260 (87.8)	32 (20.3)		
breast carcinoma estrogen receptor status (%)	Indeterminate	1 (0.3)	1 (0.6)	<0.001	exact
	Negative	2 (0.7)	109 (69.0)		
	Positive	293 (99.0)	48 (30.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	51 (19.3)	20 (14.0)	0.022	exact
	Indeterminate	2 (0.8)	0 (0.0)		
	Negative	164 (62.1)	80 (55.9)		
	Positive	47 (17.8)	43 (30.1)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.017	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.118	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-NF1	RWR-NF2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 5.00]	1.00 [1.00, 2.00]	0.002	nonnorm
distant metastasis present ind2 (%)	NO	86 (96.6)	43 (95.6)	1.000	exact
	YES	3 (3.4)	2 (4.4)		
stage event pathologic stage (%)	Stage I	31 (10.3)	9 (5.7)	0.049	exact
	Stage IA	24 (8.0)	14 (8.9)		
	Stage IB	3 (1.0)	0 (0.0)		
	Stage IIA	91 (30.3)	65 (41.1)		
	Stage IIB	65 (21.7)	41 (25.9)		
	Stage IIIA	37 (12.3)	17 (10.8)		
	Stage IIIB	10 (3.3)	2 (1.3)		
	Stage IIIC	20 (6.7)	8 (5.1)		
	Stage IV	10 (3.3)	2 (1.3)		
	Stage X	9 (3.0)	0 (0.0)		