

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
n		122	6	103	108	24	49	37	15		
Subtype (%)	wMKL1	122 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	wMKL2	0 (0.0)	6 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL3	0 (0.0)	0 (0.0)	103 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL4	0 (0.0)	0 (0.0)	0 (0.0)	108 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	24 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	49 (100.0)	0 (0.0)	0 (0.0)		
	wMKL7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	37 (100.0)	0 (0.0)		
	wMKL8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	15 (100.0)		
vital status (%)	Alive	3 (100.0)	0 (NaN)	1 (100.0)	4 (80.0)	0 (NaN)	2 (100.0)	1 (100.0)	0 (NaN)	1.000	exact
	Dead	0 (0.0)	0 (NaN)	0 (0.0)	1 (20.0)	0 (NaN)	0 (0.0)	0 (0.0)	0 (NaN)		
days to birth (median [IQR])		-21710.50 [-17518.75]	-25419.50 [-23547.00]	-22175.50 [-18170.75]	-19309.00 [-17394.00]	-23764.50 [-20551.25]	-24335.50 [-19811.00]	-23565.00 [-21655.25]	-22672.00 [-18826.50]	<0.001	nonnorm
days to death (median [IQR])		2207.00 [1012.50, 2965.00]	1877.50 [1574.75, 2180.25]	943.00 [792.00, 1563.00]	991.00 [548.00, 1556.00]	965.00 [690.75, 1167.25]	3352.00 [2574.50, 3942.00]	949.00 [390.50, 1552.50]	1793.00 [1793.00, 1793.00]	0.031	nonnorm
days to last followup (median [IQR])		1008.00 [593.25, 2192.50]	654.00 [412.75, 803.50]	651.00 [379.00, 1371.00]	996.00 [418.00, 2177.50]	921.50 [670.50, 1232.75]	791.00 [516.00, 1476.00]	958.00 [396.00, 1561.50]	683.50 [476.75, 737.00]	0.041	nonnorm
race list (%)	ASIAN	3 (2.7)	0 (0.0)	13 (17.8)	9 (9.0)	0 (0.0)	1 (3.1)	3 (11.5)	0 (0.0)	0.001	exact
	BLACK OR AFRICAN AMERICAN	10 (8.9)	1 (25.0)	8 (11.0)	27 (27.0)	4 (19.0)	2 (6.2)	4 (15.4)	1 (6.7)		
history of neoadjuvant treatment (%)	WHITE	99 (88.4)	3 (75.0)	52 (71.2)	64 (64.0)	17 (81.0)	29 (90.6)	19 (73.1)	14 (93.3)	0.219	exact
	No	120 (98.4)	6 (100.0)	102 (99.0)	108 (100.0)	24 (100.0)	47 (95.9)	35 (94.6)	15 (100.0)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
age at initial pathologic diagnosis (median [IQR])	Yes	2 (1.6)	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	2 (4.1)	2 (5.4)	0 (0.0)	<0.001	nonnorm
		59.00 [47.25, 68.00]	69.00 [63.75, 69.75]	59.00 [48.50, 71.00]	52.00 [47.00, 62.00]	64.50 [56.00, 69.00]	66.00 [54.00, 75.00]	64.00 [58.00, 69.00]	62.00 [51.50, 66.50]		
ethnicity (%)	HISPANIC OR LATINO	4 (3.8)	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	3 (9.7)	2 (8.7)	0 (0.0)	0.085	exact
	NOT HISPANIC OR LATINO	101 (96.2)	4 (100.0)	67 (100.0)	90 (98.9)	19 (100.0)	28 (90.3)	21 (91.3)	14 (100.0)		
primary lymph node presentation assessment (%)	NO	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (6.7)	0 (0.0)	0.485	exact
	YES	79 (98.8)	1 (100.0)	33 (100.0)	49 (100.0)	11 (100.0)	15 (100.0)	14 (93.3)	10 (100.0)		
histologic type (%)	Infiltrating Ductal Carcinoma	60 (49.2)	5 (83.3)	92 (89.3)	99 (91.7)	16 (66.7)	32 (65.3)	29 (78.4)	7 (46.7)	<0.001	exact
	Infiltrating Lobular Carcinoma	57 (46.7)	1 (16.7)	5 (4.9)	2 (1.9)	3 (12.5)	10 (20.4)	6 (16.2)	2 (13.3)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)	1 (6.7)		
	Mixed Histology (please specify)	2 (1.6)	0 (0.0)	1 (1.0)	2 (1.9)	0 (0.0)	2 (4.1)	1 (2.7)	0 (0.0)		
	Mucinous Carcinoma	0 (0.0)	0 (0.0)	2 (1.9)	0 (0.0)	1 (4.2)	1 (2.0)	0 (0.0)	3 (20.0)		
	Other, specify	3 (2.5)	0 (0.0)	3 (2.9)	2 (1.9)	4 (16.7)	4 (8.2)	1 (2.7)	2 (13.3)		
	Indeterminate (neither Pre or Post-menopausal)	3 (2.7)	0 (0.0)	4 (4.0)	8 (7.9)	1 (4.5)	4 (8.3)	0 (0.0)	0 (0.0)		
menopausal status (%)										0.198	exact

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
	Peri (6-12 months since last menstrual period)	5 (4.5)	0 (0.0)	3 (3.0)	5 (5.0)	2 (9.1)	2 (4.2)	0 (0.0)	1 (6.7)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	75 (67.6)	6 (100.0)	70 (69.3)	62 (61.4)	15 (68.2)	36 (75.0)	32 (94.1)	11 (73.3)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	28 (25.2)	0 (0.0)	24 (23.8)	26 (25.7)	4 (18.2)	6 (12.5)	2 (5.9)	3 (20.0)		
	Indeterminate	0 (0.0)	0 (0.0)	2 (2.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
breast carcinoma progesterone receptor status (%)	Negative	17 (13.9)	2 (33.3)	31 (30.7)	94 (89.5)	1 (4.2)	5 (11.1)	5 (13.9)	5 (33.3)		
	Positive	105 (86.1)	4 (66.7)	68 (67.3)	11 (10.5)	23 (95.8)	40 (88.9)	31 (86.1)	10 (66.7)		
breast carcinoma estrogen receptor status (%)	Indetermin	0 (0.0)	0 (0.0)	2 (2.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	6 (4.9)	0 (0.0)	9 (8.9)	91 (86.7)	0 (0.0)	0 (0.0)	0 (0.0)	5 (33.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
lab proc her2 neu im- muno- histo- chem- istry recep- tor status (%)	Positive	116 (95.1)	6 (100.0)	90 (89.1)	14 (13.3)	24 (100.0)	45 (100.0)	36 (100.0)	10 (66.7)	0.001	exact
	Equivocal	15 (14.4)	0 (0.0)	16 (16.8)	14 (15.2)	5 (23.8)	6 (15.4)	11 (31.4)	4 (26.7)		
	Indeterm	2 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Negative	74 (71.2)	4 (66.7)	40 (42.1)	62 (67.4)	12 (57.1)	25 (64.1)	17 (48.6)	10 (66.7)		
number of lym- phn- odes posi- tive by ihc (me- dian [IQR])	Positive	13 (12.5)	2 (33.3)	39 (41.1)	16 (17.4)	4 (19.0)	8 (20.5)	7 (20.0)	1 (6.7)	0.019	nonnorm
		0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]		
		0.00 [0.00, 3.00]	0.50 [0.00, 1.75]	1.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.25]	0.00 [0.00, 1.75]	0.00 [0.00, 0.00]		
er level cell per- cent- age cate- gory (me- dian [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 1.75]	1.00 [1.00, 3.50]	1.00 [1.00, 2.00]	1.00 [1.00, 11.00]	1.00 [1.00, 9.00]	1.00 [1.00, 11.00]	11.00 [1.50, 11.00]	<0.001	nonnorm
progeste recep- tor level cell per- cent cate- gory (me- dian [IQR])		1.00 [1.00, 5.00]	1.00 [1.00, 1.75]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.00 [1.00, 8.25]	1.00 [1.00, 9.00]	1.00 [1.00, 2.00]	7.00 [1.50, 10.00]	<0.001	nonnorm
distant metas- tasis present ind2 (%)	NO	48 (96.0)	0 (NaN)	24 (96.0)	27 (93.1)	5 (100.0)	9 (100.0)	8 (100.0)	8 (100.0)	0.964	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
stage event patho- logic stage (%)	YES	2 (4.0)	0 (NaN)	1 (4.0)	2 (6.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Stage I	12 (10.0)	0 (0.0)	4 (3.9)	5 (4.8)	5 (21.7)	7 (14.3)	5 (13.5)	2 (13.3)		
	Stage IA	13 (10.8)	0 (0.0)	2 (1.9)	10 (9.5)	2 (8.7)	5 (10.2)	4 (10.8)	2 (13.3)		
	Stage IB	0 (0.0)	0 (0.0)	2 (1.9)	0 (0.0)	1 (4.3)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage IIA	31 (25.8)	3 (50.0)	27 (26.2)	45 (42.9)	8 (34.8)	19 (38.8)	15 (40.5)	8 (53.3)		
	Stage IIB	23 (19.2)	1 (16.7)	36 (35.0)	27 (25.7)	3 (13.0)	8 (16.3)	5 (13.5)	3 (20.0)		
	Stage IIIA	16 (13.3)	1 (16.7)	19 (18.4)	10 (9.5)	2 (8.7)	2 (4.1)	4 (10.8)	0 (0.0)		
	Stage IIIB	1 (0.8)	1 (16.7)	2 (1.9)	2 (1.9)	1 (4.3)	3 (6.1)	2 (5.4)	0 (0.0)		
	Stage IIIC	16 (13.3)	0 (0.0)	6 (5.8)	4 (3.8)	0 (0.0)	1 (2.0)	1 (2.7)	0 (0.0)		
	Stage IV	4 (3.3)	0 (0.0)	3 (2.9)	2 (1.9)	0 (0.0)	2 (4.1)	1 (2.7)	0 (0.0)		
	Stage X	4 (3.3)	0 (0.0)	2 (1.9)	0 (0.0)	1 (4.3)	2 (4.1)	0 (0.0)	0 (0.0)		