

TCGA_holdout_Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CS1	CS2	p	test
n		227	239		
Subtype (%)	CS1	227 (100.0)	0 (0.0)	<0.001	
	CS2	0 (0.0)	239 (100.0)		
vital status (%)	Alive	4 (80.0)	7 (100.0)	0.417	exact
	Dead	1 (20.0)	0 (0.0)		
days to birth (median [IQR])		-21014.00 [-24768.00, -17894.00]	-22642.00 [-25596.50, -18564.50]	0.054	nonnorm
days to death (median [IQR])		1020.50 [752.75, 1693.75]	1787.00 [973.50, 2671.00]	0.101	nonnorm
days to last followup (median [IQR])		936.50 [384.50, 1675.50]	879.50 [536.25, 1619.00]	0.250	nonnorm
race list (%)	ASIAN	21 (11.6)	8 (3.9)	<0.001	
	BLACK OR AFRICAN AMERICAN	36 (19.9)	21 (10.3)		
	WHITE	124 (68.5)	175 (85.8)		
history of neoadjuvant treatment (%)	No	223 (98.2)	236 (98.7)	0.718	exact
	Yes	4 (1.8)	3 (1.3)		
age at initial pathologic diagnosis (median [IQR])		57.00 [48.00, 67.00]	61.00 [50.00, 69.50]	0.031	nonnorm
ethnicity (%)	HISPANIC OR LATINO	1 (0.6)	9 (4.7)	0.023	exact
	NOT HISPANIC OR LATINO	164 (99.4)	182 (95.3)		
primary lymph node presentation assessment (%)	NO	1 (1.2)	1 (0.7)	1.000	exact
	YES	79 (98.8)	135 (99.3)		
histological type (%)	Infiltrating Ductal Carcinoma	207 (91.2)	134 (56.1)	<0.001	exact
	Infiltrating Lobular Carcinoma	9 (4.0)	78 (32.6)		
	Medullary Carcinoma	1 (0.4)	0 (0.0)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CS1	CS2	p	test
	Metaplastic Carcinoma	1 (0.4)	2 (0.8)		
	Mixed Histology (please specify)	3 (1.3)	5 (2.1)		
	Mucinous Carcinoma	1 (0.4)	6 (2.5)		
	Other, specify	5 (2.2)	14 (5.9)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	14 (6.5)	6 (2.7)	0.235	
	Peri (6-12 months since last menstrual period)	9 (4.2)	10 (4.5)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	145 (67.1)	163 (72.8)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	48 (22.2)	45 (20.1)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.9)	0 (0.0)	<0.001	exact
	Negative	128 (57.7)	33 (14.1)		
	Positive	92 (41.4)	201 (85.9)		
breast carcinoma estrogen receptor status (%)	Indeterminate	2 (0.9)	0 (0.0)	<0.001	exact
	Negative	100 (45.0)	11 (4.7)		
	Positive	120 (54.1)	223 (95.3)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	30 (14.8)	41 (20.0)	<0.001	exact
	Indeterminate	0 (0.0)	2 (1.0)		
	Negative	111 (54.7)	134 (65.4)		
	Positive	62 (30.5)	28 (13.7)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.25]	0.012	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.973	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CS1	CS2	p	test
distant metastasis present ind2 (%)	NO	52 (94.5)	78 (96.3)	0.686	exact
	YES	3 (5.5)	3 (3.7)		
stage event pathologic stage (%)	Stage I	14 (6.2)	26 (11.0)	0.015	exact
	Stage IA	11 (4.9)	27 (11.4)		
	Stage IB	1 (0.4)	2 (0.8)		
	Stage IIA	80 (35.7)	76 (32.2)		
	Stage IIB	61 (27.2)	46 (19.5)		
	Stage IIIA	29 (12.9)	26 (11.0)		
	Stage IIIB	8 (3.6)	4 (1.7)		
	Stage IIIC	12 (5.4)	16 (6.8)		
	Stage IV	7 (3.1)	5 (2.1)		
	Stage X	1 (0.4)	8 (3.4)		