

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	KLIC1	KLIC2	p	test
n		599	26		
Subtype (%)	KLIC1	599 (100.0)	0 (0.0)	<0.001	exact
	KLIC2	0 (0.0)	26 (100.0)		
vital status (%)	Alive	151 (93.8)	12 (92.3)	0.586	exact
	Dead	10 (6.2)	1 (7.7)		
days to birth (median [IQR])		-21356.50 [-24650.25, -17910.50]	-20016.50 [-24876.50, -17792.50]	0.765	nonnorm
days to death (median [IQR])		1376.50 [567.75, 2430.00]	1532.00 [953.25, 2286.25]	0.525	nonnorm
days to last followup (median [IQR])		757.00 [476.25, 1611.75]	728.00 [521.50, 1362.50]	0.881	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.2)	0 (0.0)	0.763	exact
	ASIAN	31 (5.3)	1 (3.8)		
	BLACK OR AFRICAN AMERICAN	117 (20.0)	7 (26.9)		
	WHITE	437 (74.6)	18 (69.2)		
history of neoadjuvant treatment (%)	No	592 (99.2)	25 (96.2)	0.227	exact
	Yes	5 (0.8)	1 (3.8)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 67.00]	54.00 [48.25, 67.25]	0.812	nonnorm
ethnicity (%)	HISPANIC OR LATINO	25 (4.6)	2 (8.3)	0.321	exact
	NOT HISPANIC OR LATINO	515 (95.4)	22 (91.7)		
primary lymph node presentation assessment (%)	NO	25 (5.2)	1 (4.8)	1.000	exact
	YES	454 (94.8)	20 (95.2)		
histological type (%)	Infiltrating Carcinoma NOS	1 (0.2)	0 (0.0)		
	Infiltrating Ductal Carcinoma	420 (70.2)	16 (61.5)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	KLIC1	KLIC2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	115 (19.2)	3 (11.5)	0.572	exact
	Medullary Carcinoma	5 (0.8)	0 (0.0)		
	Metaplastic Carcinoma	5 (0.8)	2 (7.7)		
	Mixed Histology (please specify)	21 (3.5)	0 (0.0)		
	Mucinous Carcinoma	10 (1.7)	0 (0.0)		
	Other, specify	21 (3.5)	5 (19.2)		
	Indeterminate (neither Pre or Postmenopausal)	13 (2.4)	1 (3.8)		
	Peri (6-12 months since last menstrual period)	19 (3.5)	1 (3.8)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	385 (70.3)	20 (76.9)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	131 (23.9)	4 (15.4)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.4)	0 (0.0)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	166 (29.5)	18 (75.0)	<0.001	
	Positive	394 (70.1)	6 (25.0)		
	Negative	115 (20.5)	15 (60.0)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	447 (79.5)	10 (40.0)	0.830	exact
	Equivocal	104 (21.5)	6 (26.1)		
	Indeterminate	10 (2.1)	0 (0.0)		
	Negative	301 (62.2)	15 (65.2)		
	Positive	69 (14.3)	2 (8.7)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.116	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	KLIC1	KLIC2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 1.00]	0.020	nonnorm
er level cell percentage category (%)	<10%	32 (11.3)	3 (30.0)	0.413	exact
	10-19%	20 (7.1)	0 (0.0)		
	20-29%	3 (1.1)	1 (10.0)		
	30-39%	5 (1.8)	0 (0.0)		
	40-49%	3 (1.1)	0 (0.0)		
	50-59%	8 (2.8)	0 (0.0)		
	60-69%	7 (2.5)	0 (0.0)		
	70-79%	39 (13.8)	1 (10.0)		
	80-89%	18 (6.4)	0 (0.0)		
	90-99%	147 (52.1)	5 (50.0)		
progesterone receptor level cell percent category (%)	<10%	75 (29.3)	4 (57.1)	0.379	exact
	10-19%	18 (7.0)	0 (0.0)		
	20-29%	7 (2.7)	1 (14.3)		
	30-39%	8 (3.1)	0 (0.0)		
	40-49%	11 (4.3)	0 (0.0)		
	50-59%	9 (3.5)	1 (14.3)		
	60-69%	12 (4.7)	0 (0.0)		
	70-79%	29 (11.3)	0 (0.0)		
	80-89%	23 (9.0)	0 (0.0)		
	90-99%	64 (25.0)	1 (14.3)		
distant metastasis present ind2 (%)	NO	239 (97.6)	14 (100.0)	1.000	exact
	YES	6 (2.4)	0 (0.0)		
stage event pathologic stage (%)	Stage I	48 (8.1)	2 (7.7)	0.745	exact
	Stage IA	47 (7.9)	2 (7.7)		
	Stage IB	3 (0.5)	0 (0.0)		
	Stage II	5 (0.8)	0 (0.0)		
	Stage IIA	188 (31.6)	14 (53.8)		
	Stage IIB	142 (23.9)	6 (23.1)		
	Stage III	2 (0.3)	0 (0.0)		
	Stage IIIA	98 (16.5)	2 (7.7)		
	Stage IIIB	14 (2.4)	0 (0.0)		
	Stage IIIC	36 (6.1)	0 (0.0)		
	Stage IV	7 (1.2)	0 (0.0)		
	Stage X	4 (0.7)	0 (0.0)		